



OPEN

Neonatal head circumference by gestation reflects adaptation to maternal body size: comparison of different standards

Ruta Morkuniene¹, Janina Tutkuvienė^{1✉}, Tim J. Cole², Egle Marija Jakimaviciene¹, Jelena Isakova³, Agne Bankauskiene⁴, Nijole Drazdiene⁵ & Vytautas Basys⁶

Neonatal head circumference (HC) not only represents the brain size of *Homo sapiens*, but is also an important health risk indicator. Addressing a lack of comparative studies on head size and its variability in term and preterm neonates from different populations, we aimed to examine neonatal HC by gestation according to a regional reference and a global standard. Retrospective analysis of data on neonatal HC obtained from the Lithuanian Medical Birth Register from 2001 to 2015 (423 999 newborns of 24–42 gestational weeks). The varying distribution by gestation and sex was estimated using GAMLSS, and the results were compared with the INTERGROWTH-21st standard. Mean HC increased with gestation in both sexes, while its fractional variability fell. The 3rd percentile matched that for INTERGROWTH-21st at all gestations, while the 50th and 97th percentiles were similar up to 27 weeks, but a full channel width higher than INTERGROWTH-21st at term. INTERGROWTH-21st facilitates the evaluation of neonatal HC in early gestations, while in later gestations, the specific features of neonatal HC of a particular population tend to be more precisely represented by regional references.

Head circumference (HC) is a routine paediatric measurement that “acts as a proxy for brain size”¹. Hence, HC at birth is an indirect measure of brain growth in utero that helps, in general, to evaluate foetal growth. Although at birth the human brain is 25% of its adult weight and continues growing until the age of approximately 10 years, HC is usually of interest primarily in infancy when the head growth velocity rate is maximal¹.

Moreover, newborn HC, especially in those born preterm, is a significant health indicator. HC at birth and its postnatal growth dynamics are correlated with short-term and long-term health outcomes^{2–8}. Greater HC at birth and faster postnatal head growth are associated with better neurocognitive and intellectual abilities in adolescence and young adulthood rather than birth weight per se^{3–5}. In contrast, small HC at birth is related to the increased male risk of low intellectual performance⁶, emotional and behavioral disorders⁸, and higher arterial, especially systolic, blood pressure⁹. Considering neonatal HC an important health risk indicator for various periods of human development, an adequate HC growth assessment can facilitate not only the identification of infants at highest risk for long-term growth impairments, but also the choice of timely preventive health measures¹⁰.

Head size with its huge encephalisation ratio is the main characteristic of *Homo sapiens*. Thus it is likely to be more constant and less variable than other body size traits in a given population. On the other hand, postnatal growth is widely variable, and body size indices differ across geographic regions, populations^{11,12}, or due to socio-economic circumstances^{13–15}.

Thus tools for growth monitoring should be age- and sex-specific growth references or growth standards. Growth references describe how children from a particular region are growing, while growth standards present how children should grow under almost optimal conditions. The choice of growth reference for clinical practice is important for the evaluation of individual growth pattern. V. Neubauer et al.¹⁶ found that the interpretation

¹Department of Anatomy, Histology and Anthropology, Institute of Biomedical Sciences, Faculty of Medicine, Vilnius University, M.K. Ciurlionio str. 21, Vilnius, Lithuania. ²UCL Great Ormond Street Institute of Child Health, London, UK. ³Health Information Center, Institute of Hygiene, Didzioji str. 22, Vilnius, Lithuania. ⁴Department of Human and Medical Genetics, Institute of Biomedical Sciences, Faculty of Medicine, Vilnius University, M.K. Ciurlionio str. 21, Vilnius, Lithuania. ⁵Clinic of Children’s Diseases, Institute of Clinical Medicine, Faculty of Medicine, Vilnius University, Santariskiu str. 2, Vilnius, Lithuania. ⁶Division of Biological, Medical and Geosciences, Lithuanian Academy of Sciences, Gedimino Ave. 3, Vilnius, Lithuania. ✉email: janina.tutkuvienė@mf.vu.lt

of the postnatal growth of very preterm infants differed considerably depending on the four different references that were used: the proportion of microcephaly in very preterm infants varied from 3 to 25%. These distinct interpretations may lead to misdiagnosis and affect treatment and health monitoring strategies in clinical practice.

In 2006, the World Health Organization (WHO) published its child growth standards for children under five years. Subsequently the use of the WHO charts for particular countries or regions has been widely discussed^{14,17}. In 2008, the International Foetal and Newborn Growth Consortium for the 21st Century (INTERGROWTH-21st, IG-21) launched a multi-country project to develop similar prescriptive standards for foetal growth, neonatal size and postnatal growth of preterm infants^{18,19}.

Many recent studies^{11–13} have considered the evaluation of postnatal growth in newborns. So far, there is a lack of studies comparing head size and its variability in term and preterm neonates from different populations and geographic regions. There is no clear evidence on whether global standards for newborn HC apply to neonates from all geographical areas. Moreover, the increasing prevalence of caesarean section in clinical practice, with fewer neonates born vaginally, described by M. Oden²⁰ as a phenomenon of a sudden disappearance of the “evolutionary bottleneck”, may lead to increased variability in HC at birth.

In this context, the aim of the present study was to analyse HC in Lithuanian newborns according to their gestational age and sex and to compare the results with those provided by the IG-21 study and other countries with evolutionary insights on variability.

Methods

Study design and cohort selection. Our study examined the anonymized database from the Health Information Center of the Institute of Hygiene in Vilnius, Lithuania. The study was based on the Lithuanian Medical Data of Births registered from the year 2001 to 2015 and included all data on singleton liveborn newborns between 24 and 42 completed weeks of gestational age (GA), retrieved from medical records with the total duration of pregnancy in weeks. We excluded all cases of multiple births, stillbirth, undetermined gender, incomplete data (for sex, gestational age, birth weight, birth length, head circumference) or newborns with major congenital malformations and syndromes. The cases with the main newborn anthropometric indices (weight, length, head circumference) incompatible with gestational age (more or less than Mean (M) \pm 3 Standard Deviations (SD) following the WHO standards²¹) were removed from the analysis. In total, the cohort sample size of 423 999 newborns was derived. The sampling procedure and exclusion criteria are presented in a flow diagram (Supplementary Fig. 1).

Statistical analysis. The statistical analysis of data was performed using the standard statistical programs (SPSS 22.0, EXCEL, and R). The major parameters of descriptive statistics and percentiles of HC by GA and sex for Lithuanian newborns were calculated. The coefficient of variation (CV) was calculated and used in the comparative analysis with the foreign studies²².

GAMLSS was used to estimate the distribution applied to smooth the 3rd, 10th, 25th, 50th, 75th, 90th and 97th HC percentiles by GA and sex separately²³. The LMST method (BCT distribution) were applied to the data obtained on each sex and each measurement, respectively. The resulting main percentiles (3rd, 50th and 97th) were compared with IG-21 from 24 weeks. The analysis was carried out using the GAMLSS package (version 4.3–3) of R 4.0.3 software (www.r-project.org).

The comparison of the present data on HC of Lithuanian neonates with the data provided by the IG-21 project was conducted. Both the published standards of IG-21 project^{19,24} were presented for both sexes for every gestational week and day separately (i.e. 30 + 0, 30 + 1), while GA of the present study was recorded as complete gestational weeks (i.e. 30, 31). Therefore, the comparative analysis of the present study with the IG-21 project by GA was made by comparing the mean of HC at the specific gestational week of IG-21. The differences between the means were calculated using *t*-test. A *p*-value of < 0.05 was considered to indicate a statistically significant difference.

The data were also expressed as sex and gestation specific Z-scores using IG-21 as reference.

Ethics approval. The study was granted the approval of the governmental institution the Lithuanian Bioethics Committee (Permission No. 57, last addition—2017–02–06) and was performed in accordance with the relevant ethical guidelines and regulations.

Results

The sample size of our study (Table 1) increased dramatically with gestational age from less than 50 neonates at 24 gestational weeks to nearly 100 000 at term for each sex. The mean HC of boys was 0.5–0.8 cm greater than for girls at every gestational week. Conversely the standard deviation (SD) and the coefficient of variation (CV) of HC fell steeply with gestational age (Table 1).

The mean HC of Lithuanian preterm and term newborns was greater than for IG-21^{18,19} from 31 weeks for boys and 32 weeks for girls, the difference increasing with gestational age (Table 1). The gestational age- and sex-adjusted Z-scores of HC based on IG-21^{19,24} showed the same pattern (Supplementary Fig. 2).

The 3rd, 10th, 25th, 50th, 75th, 90th, and 97th smoothed gestational age- and sex-adjusted percentile curves for HC of Lithuanian newborns are shown in Fig. 1 and 2. The variability of HC declines with increasing gestational age, and the negative skewness in the distribution is visible as wider gaps between the lower than the upper percentiles.

Comparing the 3rd, 50th and 97th Lithuanian HC percentiles by sex and gestation with those for IG-21 confirmed the pattern seen in Table 1, of close agreement at early gestations but a widening gap with increasing gestation, though restricted to the higher percentiles (Fig. 1 and 2). On the 3rd percentile, the differences in term

GA (in weeks)	Present study				Intergrowth – 21st				Mean difference IT – IG-21
	n	M	SD	CV	n	M	SD	CV	
Boys									
24	28	23.0	1.0	0.043	3	22.7	1.6	0.070	0.3
25	71	23.6	1.4	0.059	10	23.6	1.6	0.068	0.0
26	89	24.5	1.4	0.057	13	24.5	1.6	0.065	0.0
27	124	25.5	1.5	0.059	12	25.4	1.6	0.063	0.1
28	211	26.6	1.7	0.064	19	26.3	1.6	0.061	0.3
29	190	27.5	1.7	0.062	19	27.2	1.6	0.059	0.3
30	303	28.5	1.7	0.060	25	28.1	1.6	0.057	0.4
31	306	29.7	1.7	0.057	37	28.9	1.6	0.055	0.8
32	533	30.6	1.6	0.052	52	29.8	1.6	0.054	0.8
33	744	31.5	1.6	0.051	33	31.1	1.3	0.042	0.4
34	1305	32.3	1.5	0.046	48	31.7	1.3	0.041	0.6
35	1977	33.0	1.5	0.045	127	32.2	1.3	0.040	0.8
36	3682	33.5	1.5	0.045	322	32.7	1.2	0.037	0.8
37	9651	34.3	1.5	0.044	848	33.2	1.2	0.036	1.1
38	24,745	34.9	1.4	0.040	2032	33.7	1.2	0.036	1.2
39	51,027	35.3	1.4	0.040	2985	34.1	1.1	0.032	1.2
40	93,843	35.5	1.4	0.039	2532	34.5	1.1	0.032	1.0
41	27,226	35.8	1.4	0.039	1147	34.9	1.1	0.032	0.9
42	987	35.8	1.5	0.042	204	35.2	1.1	0.031	0.6
Girls									
24	40	22.2	1.3	0.059	3	22.5	1.6	0.071	-0.3
25	65	23.1	1.3	0.056	7	23.4	1.6	0.068	-0.3
26	98	23.7	1.6	0.068	7	24.3	1.6	0.066	-0.6
27	122	25.0	1.7	0.068	11	25.1	1.6	0.064	-0.1
28	153	26.2	1.9	0.073	16	26.0	1.6	0.062	0.2
29	168	27.0	1.7	0.063	22	26.9	1.6	0.059	0.1
30	248	28.2	1.8	0.064	24	27.8	1.6	0.058	0.4
31	274	29.0	1.7	0.059	33	28.7	1.6	0.056	0.3
32	454	30.4	1.7	0.056	43	29.6	1.6	0.054	0.8
33	610	31.1	1.6	0.051	17	30.7	1.3	0.042	0.4
34	1066	31.9	1.5	0.047	65	31.3	1.2	0.038	0.6
35	1635	32.5	1.5	0.046	111	31.9	1.2	0.038	0.6
36	3183	33.1	1.5	0.045	293	32.3	1.2	0.037	0.8
37	8078	33.8	1.4	0.041	798	32.8	1.1	0.034	1.0
38	21,708	34.3	1.4	0.041	1783	33.2	1.1	0.033	1.1
39	48,487	34.8	1.4	0.040	2849	33.6	1.1	0.033	1.2
40	93,307	35.0	1.3	0.037	2486	33.9	1.1	0.032	1.1
41	26,354	35.2	1.3	0.037	1180	34.2	1.0	0.029	1.0
42	907	35.3	1.4	0.040	218	34.5	1.0	0.029	0.8

Table 1. Comparison of head circumference (HC) of Lithuanian newborns by sex and gestational age (GA) and the INTERGROWTH-21st (IG-21) reference^{19,24}. n—count, M—mean, SD—standard deviation, CV—coefficient of variation, defined as standard deviation / mean.

newborns (gestation 37–40 weeks) amounted to 0.5–0.75 cm, falling to less than 0.5 cm in the post-term period. On the 50th and 97th percentiles, the differences varied from 1–1.5 cm (Fig. 1 and 2).

Discussion

When monitoring the growth and development of neonatal HC, the primary concern is to use the best tools¹⁰. There is a lot of discussion recently concerning the choice of whether regional or global, age and sex-specific growth references or growth standards should be used for different populations^{11–13}. Our study revealed that in late preterm and term periods, with a typically smallest neonatal head circumference (HC) variability within a population, the differences between populations are the most pronounced (Table 2). The differences between the findings of the studies examined increase with the increasing GA, and particularly starting from the late preterm period, and especially, in the term newborns. In the present Lithuanian study, the variation of the mean HC in extremely, moderate to late preterm newborns HC was < 1 cm, in term newborns—> 1 cm compared to IG-21




Figure 1. The 3rd, 10th, 25th, 50th, 75th, 90th and 97th smoothed percentile curves for head circumference (cm) in Lithuanian neonate boys and the 3rd, 50th and 97th percentiles for INTERGROWTH-21st^{20,25}.

(Table 1). In extremely preterm gestations, the means of HC varies within most studies^{25–28} less than 0.5 cm compared to Lithuanian. However, according to some studies, in later gestations and in term newborns, the differences between populations in HC increase to more than 1 cm. Most of the similarities were found between Lithuanian and Finnish neonatal HC, the biggest differences – between Lithuania and Indonesia (Table 2).

Analysing the variability of HC with regard to gestational age within and between populations, the coefficient of variation (CV) was examined. According to different studies^{25–28}, the CV of HC in every population varies within a very narrow range, but is the highest in extremely preterm gestations, however, within the population, it decreases together with the increasing gestational age, same as the standard deviation (SD) (Tables 1 and 2). Hence, the closer to term, the narrower was the variability of the population's neonatal HC. The CV of HC is higher in extremely preterm periods, but HC means and extremes appear to be very similar in different populations. We presume that in early gestation there is no need to strictly set head parameters according to the mother's pelvis size, hence, greater biological variation is allowed, which is similar in most populations. On the other hand, the CV decreases with the increasing gestational age, but the means and marginal HC variants move according to a population-specific direction which is highly dependent on maternal size, particularly height and pelvic size²⁹. Here, the size of the neonatal head seems to be maximally adapted to maternal pelvic size. These considerations support the idea that head circumference is strongly anthropometrically limited by the maternal bony pelvis—“evolutionary bottleneck”, as named by M. Odent²⁰.

As the shape of the human pelvis is often interpreted as an evolutionary compromise between bipedal locomotion and childbirth of a highly encephalized neonate³⁰, HC is expected to be more strongly genetically determined and anthropometrically limited by the indices of the bony birth canal. Even though the newborn HC should be less influenced by internal or external factors than birth weight or length, many studies have raised the discussion on the complex interaction between the intrinsic and extrinsic factors in the development of neonatal HC^{31,32}. Furthermore, females with a large head, who are likely to give birth to neonates with a large head, were found to possess birth canals that are shaped to better accommodate large-headed neonates²⁹. Moreover, it is already known that variation in the shape of the female pelvis is significantly geographically structured³³.

What is more, the pelvis shape was found to be significantly associated with the stature for taller women having a more oval pelvic inlet and better accommodating a larger foetal head²⁹. In the study of R. G. Tague³⁴ femoral length/stature in females showed a significant, positive partial correlation with the anteroposterior diameter and shape of the pelvic inlet. A recent Swedish study³⁵ proved this relationship from the clinical point of view reporting decreasing risk of caesarean section (CS) with increasing maternal height after adjustment for maternal age, BMI, gestational age, parity, high birth weight and country of birth. With average Swedish women's height of 166.1 cm, maternal height of 178–179 cm was associated with the lowest risk of CS (OR=0.76, 95%

Girls




Figure 2. The 3rd, 10th, 25th, 50th, 75th, 90th and 97th smoothed percentile curves for head circumference (cm) in Lithuanian neonate girls and the 3rd, 50th and 97th percentiles for INTERGROWTH-21st^{20,25}.

GA (in weeks)	U. Sankilampi et al., FINLAND						Barbier et al., CANADA						Haksari et al., INDONESIA						Fok et al., CHINA					
	BOYS			GIRLS			BOYS			GIRLS			BOYS			GIRLS			BOYS			GIRLS		
	Mean	SD	CV	Mean	SD	CV	Mean	SD	CV	Mean	SD	CV	Mean	SD	CV	Mean	SD	CV	Mean	SD	CV	Mean	SD	CV
24	22.08	1.43	0.06	21.64	1.43	0.07	22.4	2.6	0.12	22.1	2.4	0.11	N/A	N/A	N/A	N/A	N/A	N/A	22.6	0.5	0.02	23.4	1.7	0.07
25	23.13	1.45	0.06	22.73	1.44	0.06	23.6	1.9	0.08	22.9	1.6	0.07	N/A	N/A	N/A	N/A	N/A	N/A	23.7	0.8	0.03	24.1	1.6	0.07
26	24.19	1.47	0.06	23.81	1.46	0.06	24.6	1.8	0.07	23.8	1.4	0.06	26.7	2.79	0.10	26.6	2.81	0.11	24.8	1	0.04	23.8	0.8	0.03
27	25.26	1.49	0.06	24.88	1.47	0.06	25.5	1.8	0.07	24.8	1.5	0.06	25.9	2.48	0.10	27	2.53	0.09	25.3	1.5	0.06	25	0.8	0.03
28	26.32	1.5	0.06	25.93	1.48	0.06	26.3	1.5	0.06	25.7	1.4	0.05	27.8	3.19	0.11	27.4	3.16	0.12	26.2	1.4	0.05	25.7	1.2	0.05
29	27.37	1.5	0.05	26.96	1.49	0.06	27.3	1.7	0.06	26.8	1.5	0.06	29	2.83	0.10	29.5	2.36	0.08	27.1	1.6	0.06	26.8	1.3	0.05
30	28.4	1.5	0.05	27.96	1.48	0.05	28.3	1.6	0.06	27.7	1.5	0.05	28.6	1.89	0.07	28.4	2.3	0.08	28.1	1.4	0.05	28.1	1.6	0.06
31	29.41	1.5	0.05	28.95	1.48	0.05	29.1	1.7	0.06	28.6	1.5	0.05	29.2	1.8	0.06	29.3	1.75	0.06	29	1.5	0.05	28.4	2	0.07
32	30.38	1.49	0.05	29.89	1.46	0.05	30.1	1.6	0.05	29.6	1.6	0.05	31.3	1.4	0.04	31.1	1.53	0.05	30	2	0.07	29.3	1.4	0.05
33	31.3	1.48	0.05	30.81	1.45	0.05	31.1	1.6	0.05	30.4	1.6	0.05	30.4	1.86	0.06	30.3	1.75	0.06	30.7	1.6	0.05	30.4	1.3	0.04
34	32.17	1.46	0.05	31.69	1.43	0.05	31.9	1.6	0.05	31.5	1.6	0.05	31	1.42	0.05	30.8	1.32	0.04	31.2	1.3	0.04	31.1	1.3	0.04
35	32.98	1.44	0.04	32.52	1.41	0.04	32.8	1.4	0.04	32.4	1.4	0.04	31.2	1.19	0.04	31.2	1.32	0.04	32.1	1.5	0.05	32.1	1.4	0.04
36	33.71	1.41	0.04	33.24	1.39	0.04	33.5	1.3	0.04	33.1	1.4	0.04	32.6	1.09	0.03	32.4	1.23	0.04	33.1	1.4	0.04	32.8	1.1	0.03
37	34.35	1.38	0.04	33.85	1.35	0.04	34.1	1.3	0.04	33.6	1.3	0.04	32.7	1.18	0.04	32.7	1.26	0.04	33.6	1.1	0.03	33.2	1.1	0.03
38	34.88	1.34	0.04	34.31	1.31	0.04	34.6	1.3	0.04	34	1.2	0.04	33.3	0.871	0.03	33.2	0.85	0.03	34.1	1.2	0.04	33.5	1.1	0.03
39	35.24	1.3	0.04	34.61	1.26	0.04	34.9	1.2	0.03	34.3	1.2	0.03	33.7	0.778	0.02	33.6	0.77	0.02	34.3	1.1	0.03	33.8	1.1	0.03
40	35.51	1.27	0.04	34.86	1.22	0.03	35.2	1.2	0.03	34.6	1.2	0.03	33.9	0.751	0.02	33.8	0.75	0.02	34.7	1.2	0.03	34	1.1	0.03
41	35.86	1.26	0.04	35.19	1.21	0.03	35.6	1.2	0.03	34.9	1.1	0.03	34.2	0.763	0.02	34.1	0.78	0.02	35	1.2	0.03	34.3	1.1	0.03
42	36.25	1.26	0.03	35.56	1.23	0.03	N/A	N/A	N/A	N/A	N/A	N/A	34.1	0.809	0.02	34	0.84	0.02	34.9	1.2	0.03	34.5	1.3	0.04

Table 2. The comparison of neonatal head circumference (HC) of Lithuanian newborns by sex and gestational age (GA) and its coefficient of variation (CV) and the data provided by other studies^{19,24–28}. SD—standard deviation, CV—coefficient of variation, defined as standard deviation/mean, N/A – not available.




Figure 3. Neonatal head circumference (cm) at term (40 weeks gestation) in relation to average women's height across countries^{26–29,37}.

CI 0.71–0.81), whereas height below 160 cm explained 7% of CS cases³⁵. It is worth mentioning that according to the NCD Risk Factor Collaboration³⁶, Lithuanian women are among the tallest women in the world with an average height of 167.6 cm. The comparative results of the female average height reflect the differences found between the mean neonatal HC from different populations, as shown in Table 2. Finnish women with an average height of 166.5 cm are closest to Lithuanians, followed by Canadians (164.7 cm), Chinese (163.5 cm), and finally Indonesians (154.4 cm)³⁶. This supports the previous study's findings²⁹ that perhaps maternal height is linked to pelvic size, particularly the size of the birth canal, and through that to the neonatal HC. This possible relationship between neonatal HC (cm) at term (40 weeks of gestation) and average women's height across compared countries is presented in Fig. 3 and compiled after^{26–29,35}.

With regard to these findings, scientists debate the appropriateness of growth standards vs. references, regional vs. global for proper evaluation of growth and development of neonatal HC. In our study, most of similarities with global study of IG-21 were disclosed in cut-off points for the lowest percentiles in extremely preterm newborns, apart from that, extremely preterm newborns (especially girls) had more similarities not only in the third, but also in the 50th and 97th percentiles of HC. The Brazilian study³⁷ revealed a similar pattern and found the trajectory of the third percentile parallel with the IG-21 study until the term period. The sample size of the IG-21¹⁹ reference was only modest for < 37 weeks gestation, and the later study on very preterm neonates should be interpreted with caution given the small sample size²⁴. This may explain a “wave” at 33 gestational weeks observed in the percentile curves of IG-21 (Figs. 1 and 2). Thus, although IG-21 facilitates the evaluation of the main HC percentiles for extremely preterm newborns and might serve as cut-off points for the pathological microcephaly in preterm newborns of different populations, it should be considered with caution to be confidently used as a global standard at early gestations.

As for the other extreme, the 97th percentile, above which infants would be diagnosed with macrocephaly, a large gap between the curves of both studies of more than 1 cm from late preterm to post-term was detected which could lead to an overestimation of macrocephaly in our cohort. If we compared our results with the HC curves provided by the Centers for Disease Control and Prevention (CDC)³⁸, the gap would be smaller. In line with other studies³⁹ evaluating the influence of growth curves used for the distribution of HC, our study also claims that the important consequences could have been triggered by the percentile misclassification. Therefore, from the standpoint of clinical practice, to predict the course of HC higher percentiles in moderate or late preterm periods and, especially, in Lithuanian term newborns, regional standards should be used.

Accordingly, the question has been raised by scientists whether children's growth references should be global, or specific to different populations: 'it has become apparent that a single “global” reference fails adequately to mirror the diversity in human growth¹⁷. Human growth is determined by inherited factors, and the significant variability of foetal growth in utero between ethnic groups supports this statement³¹. Therefore, the IG-21 project charts based on the idea that foetuses, infants, and children grow similarly all over the world under ideal nutritional, environmental, psychological living conditions have been widely discussed. A number of studies^{32,37,40–46} recently have compared their foetal and neonatal national growth references with the IG-21 study that was recently published, and obtained diverse results. Some studies did not find appreciable differences with IG-21 for newborn HC⁴⁰ or a statistically significant difference was observed only of female HC in the 97th percentile³². While others determined that IG-21 standards for foetuses⁴⁷ were found to be unrepresentative for regional populations leading to considerable overdiagnosis of foetal microcephaly or misclassification of infant birth size^{37,41–46} and a conclusion that regional validation was needed prior to the implementation of IG-21. We found that global standards like INTERGROWTH-21st might facilitate the evaluation of neonatal head circumference in early gestations, while in later gestations, the specific features of neonatal head circumference of a particular population tend to be more precisely represented by regional standards.

Therefore, we suggest taking into consideration the regional standards for neonatal head circumference in order to better evaluate a possible clinical pathology. It is important to stress that over the process of evolution,

neonatal body size and head circumference have adapted to the mother's body size, especially her pelvis, as a result of diverse adaptation mechanisms common to different populations in different geographical areas and under different living conditions.

Conclusions

The closer to the late preterm—term period, the greater the differences between neonatal head circumferences in different populations. This threshold is slight, but it marks the inevitable influence of the evolutionary mechanisms that operate to first concentrate on vital biological capacities for neurodevelopment and only then allow genetics, ethnicity and other complex factors to influence the variability of neonatal head circumference.

Consequently, the global standards as IG-21 may serve for the evaluation of HC in early gestations, taking into account that most countries do not have a possibility to construct their own references due to small numbers of neonates born extremely preterm. In later gestations, regional standards more precisely represent the specific features of the neonatal HC of a particular population.

Data availability

The data that support the findings of this study are available at the Health Information Center of the Institute of Hygiene of Lithuania, however restrictions apply to the availability of these data, used under a license for the current study, therefore they are not publicly available. The data are available from the corresponding author upon a reasonable request and with the permission of the Health Information Center of the Institute of Hygiene of Lithuania.

Received: 16 August 2021; Accepted: 20 June 2022

Published online: 30 June 2022

References

- Cameron, N. & Schell, L. *Human Growth and Development* (Elsevier, 2021).
- Bocca-Tjeertes, I. F. A., Reijneveld, S. A., Kerstjens, J. M., de Winter, A. F. & Bos, A. F. Growth in small-for-gestational-age preterm-born children from 0 to 4 Years: The role of both prematurity and SGA Status. *Neonatology* **103**, 293–299 (2013).
- Jensen, R. B., Juul, A., Larsen, T., Mortensen, E. L. & Greisen, G. Cognitive ability in adolescents born small for gestational age: Associations with fetal growth velocity, head circumference and postnatal growth. *Early Hum. Dev.* **91**, 755–760 (2015).
- Ranke, M. B., Krägeloh-Mann, I. & Vollmer, B. Growth, head growth, and neurocognitive outcome in children born very preterm: Methodological aspects and selected results. *Dev. Med. Child Neurol.* **57**, 23–28 (2015).
- Sammallahti, S. *et al.* Infant growth after preterm birth and neurocognitive abilities in young adulthood. *J. Pediatr.* **165**, 1109–1115 (2014).
- Bergvall, N. Birth Characteristics and risk of low intellectual performance in early adulthood: Are the associations confounded by socioeconomic factors in adolescence or familial effects?. *Pediatrics* **117**, 714–721 (2006).
- Dekhtyar, S. *et al.* Associations of head circumference at birth with early life school performance and later-life occupational prestige. *Longitud. Life Course Stud.* **6**, 26–42 (2015).
- Nomura, Y. & Gampel, S. B. Short and long-term effects of compromised birth weight, head circumference, and apgar scores on neuropsychological development. *J. Psychol. Abnorm. Child.* <https://doi.org/10.4172/2329-9525.1000127> (2014).
- Paulauskienė, S. *The Relation Between Elevated Blood Pressure in Young Men and Critical Growth Periods* (Vilnius University, 2011).
- Glass, H. C. *et al.* Outcomes for extremely premature infants. *Anesth. Analg.* **120**, 1337–1351 (2015).
- Juliussen, P. B., Roelants, M., Hoppenbrouwers, K., Hauspie, R. & Bjerknes, R. Growth of Belgian and Norwegian children compared to the WHO growth standards: Prevalence below -2 and above +2 SD and the effect of breastfeeding. *Arch. Dis. Child.* **96**, 916–921 (2011).
- Nielsen, A., Olsen, E. & Juul, A. New Danish reference values for height, weight and body mass index of children aged 0–5 years: New growth references for 0- to 5-year-old children in Denmark. *Acta Paediatr.* **99**, 268–278 (2009).
- Bertino, E. *et al.* Neonatal anthropometric charts: The Italian neonatal study compared with other European studie. *J. Pediatr. Gastroenterol. Nutr.* **51**, 353–361 (2010).
- Hui, L. L. *et al.* Are universal standards for optimal infant growth appropriate? Evidence from a Hong Kong Chinese birth cohort. *Arch. Dis. Child.* **93**, 561–565 (2008).
- Hermanussen, M., Aßmann, C. & Tutkuviene, J. Statistical agreement and cost–benefit: Comparison of methods for constructing growth reference charts. *Ann. Hum. Biol.* **37**, 57–69 (2010).
- Neubauer, V., Fuchs, T., Griesmaier, E., Pupp-Peglow, U. & Kiechl-Kohlendorfer, U. Comparing growth charts demonstrated significant deviations between the interpretation of postnatal growth patterns in very preterm infants. *Acta Paediatr.* **105**, 268–273 (2016).
- Hermanussen, M., Staub, K., Assmann, C. & van Buuren, S. Dilemmas in choosing and using growth charts. *Pediatr. Endocrinol. Rev. PER* **9**, 650–656 (2012).
- Villar, J. *et al.* Postnatal growth standards for preterm infants: The Preterm Postnatal Follow-up Study of the INTERGROWTH-21st Project. *Lancet Glob. Health* **3**, e681–e691 (2015).
- Villar, J. *et al.* International standards for newborn weight, length, and head circumference by gestational age and sex: the Newborn Cross-Sectional Study of the INTERGROWTH-21st Project. *The Lancet* **384**, 857–868 (2014).
- Odent, M. The evolution of neonatal head circumference. *SM J Public Health Epidemiol* **2**, 1029 (2016).
- WHO child growth standards: head circumference-for-age, arm circumference-for-age, triceps skinfold-for-age and subscapular skinfold-for-age: methods and development. (World Health Organization, 2007).
- Lovlie, P. Coefficient of Variation. in *Encyclopedia of Statistics in Behavioral Science* (American Cancer Society, 2005). doi:<https://doi.org/10.1002/0470013192.bsa107>.
- Cole, T. J. The LMS method for constructing normalized growth standards. *Eur. J. Clin. Nutr.* **44**, 45–60 (1990).
- Villar, J. *et al.* INTERGROWTH-21st very preterm size at birth reference charts. *The Lancet* **387**, 844–845 (2016).
- Sankilampi, U., Hannila, M.-L., Saari, A., Gissler, M. & Dunkel, L. New population-based references for birth weight, length, and head circumference in singletons and twins from 23 to 43 gestation weeks. *Ann. Med.* **45**, 446–454 (2013).
- Barbier, A. *et al.* New reference curves for head circumference at birth, by gestational age. *Pediatrics* **131**, e1158–e1167 (2013).
- Haksari, E. L., Lafeber, H. N., Hakimi, M., Pawirohartono, E. P. & Nyström, L. Reference curves of birth weight, length, and head circumference for gestational ages in Yogyakarta, Indonesia. *BMC Pediatr.* **16**, 188 (2016).
- Fok, T. F. *et al.* Updated gestational age specific birth weight, crown-heel length, and head circumference of Chinese newborns. *Arch. Dis. Child. Fetal Neonatal Ed.* **88**, F229–F236 (2003).

29. Fischer, B. & Mitteroecker, P. Covariation between human pelvis shape, stature, and head size alleviates the obstetric dilemma. *Proc. Natl. Acad. Sci.* (2015).
30. Rosenberg, K. & Trevathan, W. Bipedalism and human birth: The obstetrical dilemma revisited. *Evol. Anthropol. Issues News Rev.* **4**, 161–168 (1995).
31. Buck Louis, G. M. *et al.* Racial/ethnic standards for fetal growth: the NICHD Fetal Growth Studies. *Am. J. Obstet. Gynecol.* **213**, 1–41 (2015).
32. Villamonte-Calanche, W. *et al.* Neonatal anthropometry at 3400 m above sea level compared with INTERGROWTH 21st standards. *J. Matern. Fetal Neonatal Med.* **30**, 155–158 (2017).
33. Betti, L. & Manica, A. Human variation in the shape of the birth canal is significant and geographically structured. *Proc. R. Soc. B Biol. Sci.* **285**, 20181807 (2018).
34. Tague, R. G. Do big females have big pelvises?. *Am. J. Phys. Anthropol.* **112**, 377–393 (2000).
35. Mogren, I. *et al.* Maternal height and risk of caesarean section in singleton births in Sweden—A population-based study using data from the Swedish Pregnancy Register 2011 to 2016. *PLoS ONE* **13**, e0198124 (2018).
36. NCD Risk Factor Collaboration. Evolution of Adult Height over Time. <https://www.ncdrisc.org/data-downloads-height.html> (7 April 2021, date last accessed). <https://www.ncdrisc.org/data-downloads-height.html>.
37. Pimenta, J. R. R., Grandi, C., Aragon, D. C. & Cardoso, V. C. Comparison of birth weight, length, and head circumference between the BRISA-RP and Intergrowth-21st cohorts. *J. Pediatr. Engl. Ed.* **96**, 511–519 (2020).
38. Growth Charts—Data Table of Infant Head Circumference-for-age Charts. https://www.cdc.gov/growthcharts/html_charts/hcage_inf.htm (2019).
39. Daymont, C., Hwang, W.-T., Feudtner, C. & Rubin, D. Head-circumference distribution in a large primary care network differs from CDC and WHO curves. *Pediatrics* **126**, e836–e842 (2010).
40. Stirnemann, J. J. *et al.* Implementing the INTERGROWTH-21st fetal growth standards in France: a ‘flash study’ of the Collège Français d’Echographie Foetale (CFEF): Implementing INTERGROWTH-21st fetal size charts in France. *Ultrasound Obstet. Gynecol.* **49**, 487–492 (2017).
41. Leibovitz, Z. *et al.* Prediction of microcephaly at birth using three reference ranges for fetal head circumference: Can we improve prenatal diagnosis?: Prediction of microcephaly at birth. *Ultrasound Obstet. Gynecol.* **47**, 586–592 (2016).
42. Cheng, Y., Leung, T., Lao, T., Chan, Y. & Sahota, D. Impact of replacing Chinese ethnicity-specific fetal biometry charts with the INTERGROWTH-21st standard. *BJOG Int. J. Obstet. Gynaecol.* **123**, 48–55 (2016).
43. Sotiriadis, A. *et al.* National curves of foetal growth in singleton foetuses of Greek origin. *Eur. J. Clin. Invest.* **46**, 425–433 (2016).
44. Sletner, L., Kiserud, T., Vangen, S., Nakstad, B. & Jennum, A. K. Effects of applying universal fetal growth standards in a Scandinavian multi-ethnic population. *Acta Obstet. Gynecol. Scand.* **97**, 168–179 (2018).
45. Bendor-Samuel, O. M., Zivanovic, S., Odd, D. & Roehr, C. C. A Comparison of UK preterm anthropometric charts and intergrowth-21st: Is it time to change growth charts?. *Neonatology* **117**, 300–307 (2020).
46. Pam, V. C. *et al.* Head circumference of babies at birth in Nigeria. *J. Trop. Pediatr.* **65**, 626–633 (2019).
47. Papageorgiou, A. T. *et al.* International standards for fetal growth based on serial ultrasound measurements: The Fetal Growth Longitudinal Study of the INTERGROWTH-21st Project. *The Lancet* **384**, 869–879 (2014).

Author contributions

R.M. analysed the data, performed the calculations, and took the lead in writing the manuscript with the input from other authors. J.T. raised the main conceptual idea, designed and supervised the study, helped in data interpretation and manuscript writing, revised the final version. T.J.C. contributed in revising the manuscript, and advising on the valuable improvements to the quality and the conceptual idea of the study. E.M.J., N.D., V.B. contributed to the interpretation of the results and the final version of the manuscript. J.I. collected the database according to the required inclusion criteria. A.B. performed the statistical analysis and provided suggestions for data interpretation.

Funding

No external funding was received for this study.

Competing interests

The authors declare no competing interests.

Additional information

Supplementary Information The online version contains supplementary material available at <https://doi.org/10.1038/s41598-022-15128-3>.

Correspondence and requests for materials should be addressed to J.T.

Reprints and permissions information is available at www.nature.com/reprints.

Publisher’s note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.



Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article’s Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article’s Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

© The Author(s) 2022