



# Theory Paper: Suggesting Compassion-Based Approaches for Treating Complex Post-traumatic Stress Disorder

Annabel Rushforth<sup>1</sup> · Yasuhiro Kotera<sup>1,2</sup> · Greta Kaluzeviciute<sup>1,3</sup>

Accepted: 5 June 2022 / Published online: 11 July 2022  
© The Author(s) 2022

## Abstract

Complex post-traumatic stress disorder (CPTSD) may develop following interpersonal and cumulative traumatic events, usually during early development. In addition to the core PTSD symptom profile, CPTSD presents emotional dysregulation symptoms that can be resistant to conventional treatments. Compassion-focused therapy (CFT) may be an effective intervention for addressing the more resistant symptoms in the emotional stabilisation phase of treatment rather than the trauma-processing phase. This paper explores the diagnostic validity and prevalence of CPTSD, treatment recommendations and the role of CFT in mediating shame and stabilising emotional dysregulation. We also evaluate current evidence utilising compassion-based interventions for the components of the CPTSD symptom profile and the viability of CFT as a whole. The novelty of CPTSD as a clinical condition means there is limited evidence regarding recommended treatment. Research into the efficacy of CFT and its suitability to target CPTSD's symptom profile will contribute to the current gap in recommended treatment approaches for this condition.

**Keywords** Complex post-traumatic stress disorder · CPTSD · Compassion-focused therapy · Emotional dysregulation · Emotional stabilisation phase · Trauma-processing phase

Complex post-traumatic stress disorder (CPTSD) was initially proposed as a disorder distinct from PTSD by Herman (1992), who argued that traumatic, usually childhood experiences in CPTSD are repetitive and prolonged, with the individual feeling unable to escape (Cloitre et al., 2009). The CPTSD symptom profile includes the underdevelopment of emotional, social, cognitive and psychological competencies. CPTSD diagnostic criteria include the three core PTSD symptom clusters of re-experiencing, avoidance and a sense of threat, with three additional clusters unique to CPTSD: affect dysregulation, negative

---

✉ Annabel Rushforth  
a.rushforth1@unimail.derby.ac.uk

<sup>1</sup> College of Health, Psychology and Social Care, University of Derby, Kedleston Rd, DE22 1GB Derby, UK

<sup>2</sup> Faculty of Medicine and Health Sciences, University of Nottingham, NG7 2TU Nottingham, UK

<sup>3</sup> Institute of Psychology, Faculty of Philosophy, Vilnius University, Universiteto g. 9, LT-01513 Vilnius, Lithuania

self-concept and disturbances in relationships (World Health Organization, 2018). These three additional clusters are collectively known as ‘disturbances in self-organisation’ (DSO) symptoms (Cloitre et al., 2011; Maercker et al., 2013). The additional symptoms reflect the unique impact complex traumatic experiences have on the development of interpersonal skills and the emotional regulation system (Burns et al., 2010; Shipman et al., 2005) and hippocampal function (Chetty et al., 2014) as well as the ability to form secure attachments (Karatzias et al., 2019b). CPTSD symptoms differ from the primary risk factor for PTSD which is one incident of intense trauma (Zuj et al., 2016). CPTSD is more likely to arise following a combination of the above risk factors (Karatzias et al., 2019b) as they specifically impact complexity of symptoms (Cloitre et al., 2009; Hyland et al., 2017).

Following multiple empirical investigations for the validity of construct (Cloitre et al., 2013; Karatzias et al., 2017; Perkonig et al., 2016; Sachser et al., 2017), CPTSD was only recently recognised as distinct from PTSD in the *International Classification of Diseases*, 11th revision (ICD-11; WHO, 2018). The ICD-11 points to DSO symptoms in order to differentiate CPTSD from PTSD. Meanwhile, CPTSD is not currently recognised by the *Diagnostic and Statistical Manual of Mental Disorders*, 5th revision (DSM-5; APA, 2013), although it is possible that the next revision of the DSM will consider recognising the disorder (Cloitre et al., 2020).

The utility and discriminatory validity of separating the two disorders have been a topic of debate. Prior to the publication of ICD-11, Resick et al. (2012) argued that there is no substantial evidence to support the need for a new diagnostic category, stating that the clinical benefit had not been established and that current research lacked rigour. In a response, Herman (2012) argued that the authors had set an arbitrarily high standard for inclusion of research. Additionally, Cloitre et al. (2012) argued it would be clinically beneficial to have simplicity in the overall classification structure, which would in turn improve the conceptual organisation of symptoms and causes relevant to CPTSD.

This rebuttal has particular relevance to the DSM, given that the DSM-5 diagnostic criteria for PTSD underwent revisions recognising additional DSO symptoms as part of the core symptom profile. This revision, according to Cloitre et al. (2013), reduces efficiency, as mental health providers prefer diagnoses to have a limited number of symptoms, so as to increase diagnostic goodness of fit (accuracy of description for any one patient), and thus to increase efficacy of treatment. Finally, the distinction between PTSD and CPTSD, as Cloitre et al. (2009) argue further, allows for a better understanding of treatment options, which is especially important given the prevalence of CPTSD: 2.6% in Israel (Ben-Ezra et al., 2018) and 3.8% in the USA (Cloitre et al., 2019), compared with a 6.8% prevalence of PTSD (Gradus, 2007). However, it is possible that CPTSD may be underdiagnosed as it is a novel and emerging disorder.

The International Society for Traumatic Stress Studies (ISTSS) recommends that CPTSD treatment be delivered in a phase-based approach, initially addressing DSO symptoms through an emotional stabilisation phase, prior to addressing the core PTSD symptoms through a trauma-processing phase (Cloitre et al., 2011). One third of PTSD patients with persistent DSO symptoms are less responsive to traditional clinical interventions (Winders et al., 2020; Van der Kolk et al., 2007) such as prolonged exposure and cognitive-processing therapy (Jonas et al., 2013; Watkins et al., 2018). Gilbert and Irons (2004) and Karatzias et al. (2019a) propose that these more psychologically pervasive DSO symptoms may be more responsive to compassion-based interventions, and individuals with CPTSD who engage in compassion-based interventions to address these DSO symptoms will see greater symptom reduction than those that engage in traditional clinical interventions alone.

## Compassion-Based Interventions as a Treatment for CPTSD

Compassion-based interventions can be effective when addressing symptomology associated with shame (Karatzias et al., 2019a). Lee et al. (2001) propose that shame can mediate CPTSD symptoms and Karatzias et al. (2019a) have shown an association between the two. The Ehlers and Clark's (2000) cognitive model of PTSD focuses on mitigating fear; symptoms arise from event processing in a way that maintains a sense of threat. Treatment attempts to re-process the memories of the event, reducing maladaptive coping strategies. Lee et al.'s (2001) alternative model states that the presence of shame following a traumatic event can maintain a self-critical dialogue and in turn a sense of threat. Shame can become an obstacle to recovery, affecting the individual's ability to engage in more traditional clinical interventions such as cognitive behavioural therapy (CBT). Lee et al. (2001) propose working with compassion-based interventions, such as compassion-focused therapy (CFT), to reduce shame and self-criticism in PTSD sufferers.

CFT was initially developed as an adjunct to CBT by Gilbert (2009), aiming to balance the affiliative system and reduce threat-based processing that can contribute to psychopathology. CFT seeks to help individuals respond to their distress through a compassionate lens, enhancing the capacity to self-soothe and reduce self-critical self-talk (Gilbert, 2009, 2010), thus allowing the individual to feel safer in their environment.

There are currently no randomised controlled trials evaluating the efficacy of CFT for CPTSD. However, Karatzias et al. (2019a) found significant associations between low scores on all self-compassion subscales (self-kindness, self-judgement, common humanity, isolation, mindfulness and over-identification) and the existence of emotional hypoactivation and negative self-concept. Significant negative association between four self-compassion subscales (self-kindness, self-judgement, common humanity and isolation) and disturbances in relationships were also identified. High self-judgement significantly predicted affect dysregulation and negative self-concept, which are associated with resistance to traditional clinical interventions in CPTSD (Gilbert & Irons, 2004). The most salient issue with this study is that analyses were based on correlation, and therefore could not deduce causality. Furthermore, self-report scales may involve biased results particularly with populations that experience high levels of shame (Van de Mortel, 2008), as this may lead to dishonest responses.

Despite this, these findings are an important first step in identifying psychological factors that are associated with CPTSD, which can inform future interventional research. Specifically, Karatzias et al. (2019a) found that self-compassion was not significantly associated with core PTSD symptom clusters despite several studies showing this relationship (Winders et al., 2020). This may be partially due to previous studies using DSM-5 (APA, 2013) rather than the ICD-11 (WHO, 2018) diagnostic criteria. The ICD-11 tightened its criteria for PTSD to avoid over-diagnosing (Hansen, 2017). In contrast, the DSM-5 expanded its criteria for PTSD from primarily a fear-based disorder to include 'negative alterations in cognition and mood'. These expanded symptoms include persistent negative evaluation of self or others, elevated self-blame, negative emotional states and reckless and self-destructive behaviour (Stein et al., 2014). These symptoms are parallel with the DSO symptom criteria for CPTSD in the ICD-11. As a result, it is possible that studies using the expanded criteria will find correlates between the DSM-5 PTSD symptoms of 'negative alterations in cognitions and mood' and self-compassion subscales, which other studies using the tighter ICD-11 core PTSD symptom criteria will not. This compromises clinical utility and thus the development of treatment protocols. Uniformity regarding this would

contribute to cohesion in future research especially where CPTSD is concerned. As the ICD-11 diagnostic criteria for CPTSD could be thought of as a division between different DSM-5 symptom clusters (Stein et al., 2014), research targeting these clusters in isolation will allow differentiation between correlates of PTSD and CPTSD, further highlighting if these sibling disorders require qualitatively different treatment protocols. In the absence of data regarding the efficacy of CFT for CPTSD, research regarding compassion-based interventions for CPTSD's constituent parts, firstly the DSO symptom profile and secondly the core PTSD symptom profile, would provide valuable insight.

## Compassion-Based Interventions as a Treatment for Disturbances in Self-organisation

Prior to a trauma-processing phase of treatment, an emotional stabilisation phase of treatment is recommended to address the DSO symptoms, ensure patient safety and avoid exacerbation of symptoms during the following trauma-processing phase (Cloitre et al., 2011). To date, there have been no studies evaluating compassion-based interventions as a treatment for DSO symptoms. In the absence of these studies, it is possible to assess the available research for significant effects of compassion-based interventions on the DSO symptom indicators of affect dysregulation, negative self-concept and disturbed relationships as described by the International Trauma Questionnaire (ITQ; Shevlin et al., 2018).

Research compiled from three meta-analyses (Ferrari et al., 2019; Kirby et al., 2017; Wilson et al., 2019), one systematic review (Craig et al., 2020) and one literature review (Kirby, 2017) regarding the efficacy of compassion-based interventions yielded five interventional studies using comparison groups exploring loving kindness meditation, mindfulness-based cognitive therapy with a self-compassion element, mindful self-compassion and CFT (mindfulness-only-based interventions were excluded) as a treatment for clinical populations displaying similar symptoms to the constituent components of DSO (emotionally unstable personality disorder, depressive symptoms, mental health difficulties). The above compassion-based interventions utilise the same theoretical framework as CFT, aiming to reduce shame, balance the affiliative system and encourage self-soothing during acute stress. Of the five available studies, only two used an active control group. One of the two active control trials, Feliu-Soler et al. (2017), found significant improvements in self-concept following a 3-week loving kindness meditation relative to a mindfulness meditation intervention. This study may however lack population validity as it used a predominantly female sample. Although a lack of follow-up makes it difficult to ascertain if the results were maintained, positive outcomes were achieved in a short treatment duration. In the other active control trial, Cuppage et al. (2018) found a similar result with CFT showing significant improvements in psychopathology and a reduction in self-criticism. The transdiagnostic nature of the group suggests that CFT is addressing the underlying psychological processes giving rise to these psychopathologies, rather than just the symptoms presented. This aligns with Lee et al.'s (2001) theory of self-criticism contributing to psychopathology. Of the remaining three studies, all found significant improvements in depressive symptoms (Neff & Germer, 2013; Lee & Bang, 2010; Shahar et al., 2015).

Additional research shows significant improvements in outcome measures across domains of anxiety (Gharraee et al., 2018; Navab et al., 2019; Noorbala et al., 2013; Sommers-Spijkerman et al., 2018), substance abuse disorder (Carlyle et al., 2019) and eating disorders (Duarte et al., 2017). The available evidence suggests that CFT may be a suitable

treatment to address the range of complex symptoms in the emotional stabilisation phase. In the absence of an agreed treatment protocol, it is worthwhile investigating this with actively controlled, large-scale studies to establish efficacy relative to other treatments.

## Compassion-Based Interventions as a Treatment for PTSD

Because CPTSD can be resistant to conventional PTSD treatments, alternative approaches need to be evaluated. Much of the research regarding the efficacy of compassion-based interventions in PTSD treatment lacks statistical significance, control groups or causal considerations. Research compiled from a recent systematic review (Winders et al., 2020) yielded nine interventional studies exploring compassion-based interventions for PTSD. Of these nine, five used comparison groups, and of these five, one showed a greater significant reduction in PTSD symptoms in the compassion-based group (using cognitive-based compassion training) relative to the control group (Lang et al., 2017). Of the four remaining studies using comparison groups, one showed an improvement in PTSD symptoms in both the compassion-based condition (imagery re-scripting) and the traditional clinical intervention condition (imaginal exposure) with a slightly larger improvement in the traditional intervention condition (Hoffart et al., 2015). The remaining three showed a non-significant improvement (Beaumont et al., 2012, 2016; Held & Owens, 2015), although these used small sample sizes and were therefore potentially underpowered, giving rise to type II errors. Of the four studies without a control group, three found a significant reduction in PTSD symptoms following compassion-based interventions (Au et al., 2017; Kearney et al. 2013; Muller-Engelmann et al., 2019). The study that did not find a reduction in PTSD symptoms did however find an improvement in affect (Held et al., 2018). The absence of a control group makes it difficult to ascertain if the observed effect is due to the nature of compassion-based interventions and their effect of reducing shame and improving self-soothing capacity, or due to the effect of engaging in treatment generally. Interestingly, significant correlations between PTSD symptom severity and self-compassion have been demonstrated consistently in populations with childhood abuse or interpersonal trauma (Barlow et al., 2017; Bistricky et al., 2017; Miron et al., 2015; Thomson & Waltz's, 2008) implying the inconsistency of results in the above research may be mediated by this risk factor.

Although this research is promising, it is difficult to draw conclusions about causality from cross-sectional studies. For example, it is possible that low self-compassion is a pre-trauma risk factor for PTSD, or that self-compassion is better adopted by those with less severe symptomatology. Greater symptom reduction was found in populations with either higher self-judgement or elevated trauma-related shame (Au et al., 2017; Thomson & Waltz's, 2008). Further research is needed to clarify if this effect is only relevant to sub-populations of PTSD, such as those with complex trauma, as the research may not generalise to all PTSD populations.

Given (a) the lack of statistical significance, control groups and causal considerations in the majority of compassion-based intervention studies for PTSD and (b) the prevalence of literature reporting the efficacy of traditional clinical interventions for trauma processing such as trauma-focused cognitive-processing therapy (Jonas et al., 2013; Watkins et al., 2018), large-scale controlled studies are needed to propose CFT or other similar compassion-based interventions as a primary treatment during the trauma-processing phase of treatment for CPTSD. Future interventional and controlled research is needed to support and add causality to the

results seen in the cross-sectional data. In addition to this, future research would benefit from improved statistical power. This could be achieved through use of a repeated measures design, although this can be difficult in therapeutic research; another way to achieve this is to increase sample sizes. Additionally, improvement in symptomology can be seen as a result of engaging in treatment independent of the specific intervention (Safer & Hugo, 2006), and therefore future research could benefit from including an active control group using traditional clinical interventions to account for this effect. The addition of this research would justify further investigation into CFT as a primary treatment in the trauma-processing phase of treatment of CPTSD.

## Acceptability and Viability of Compassion-Based Interventions

Although efficacy of interventions can be assessed with outcome measures, their acceptability and viability are other important considerations (Moore et al., 2015). CFT has been found to have good attrition among a variety of populations including those with dementia (94%; Collins et al., 2018), perfectionism (96%; Rose et al., 2018) and psychosis (82%; Laithwaite et al., 2009) and those in an inpatient setting (95%; Braehler et al., 2013). Given the challenges faced regarding interpersonal domains as described above, this has particular relevance, as populations with CPTSD may find it difficult to commit to a programme of treatment given these disturbances (16–20%; Vogel et al., 2017). In addition, several studies have measured individual satisfaction with CFT, with consistently high levels reported (Duarte et al., 2017; Graser et al., 2016; Clapton et al., 2018), although it is worth noting that populations with high shame may feel unable to give critical feedback, so these results should be interpreted cautiously. Further research into satisfaction with the intervention would benefit from being obtained blind to account for this effect. These are particularly important factors in the evaluation of CFT for those who complete the therapy in groups, as it can be discouraging to peers for the group to reduce in size or be described negatively by peers. CFT has been effectively delivered in groups (McManus et al., 2018; Feliu-Soler et al., 2017; Heriot-Maitland et al., 2014), improving cost effectiveness, waitlist time and viability for treatment centres. This is particularly salient considering one implication for counsellors is the additional training they may need to undertake to effectively deliver CFT.

Further studies evaluating the extent of this for 1:1 and group delivery would inform decisions regarding the cost effectiveness versus potential changes in outcome measures. An additional issue regarding the application of CFT is an absence of implementation fidelity, as details regarding primary outcomes, who delivered the therapy, and their training were lacking (Craig et al., 2020). This can cause disparity in interpretation of results, and make it difficult for group facilitators, counsellors or psychologists to deliver CFT in the way the research has shown it to be effective. Future research would benefit from uniformity in these areas. Although CFT could benefit from this, it is considered a highly acceptable treatment option by patients and is well delivered in groups. In addition to outcome efficacy, these are important considerations when assessing its strength as an intervention.

## Conclusions

In conclusion, the recent decision by the IDC-11 to recognise CPTSD as a disorder distinct from PTSD has raised important questions regarding risk factors, validity of construct and clinical utility. However, it has been shown (Cloitre et al., 2013) that the new diagnostic criteria promote ease of use and simplicity. It has allowed scope for studies of prevalence of CPTSD, highlighting the need to assess potential treatments. The current recommendations of a phase-based approach of treatment (Cloitre et al., 2011) entail addressing DSO symptoms separately to core PTSD symptoms as this promotes ease of treatment. Evidence supports the use of CFT as a primary intervention for DSO symptoms in the emotional stabilisation phase although there are other more suitable and well-researched interventions for the core PTSD symptoms in the trauma-processing phase. As a primary treatment for DSO symptoms, CFT could benefit from further uniformity regarding delivery, although the treatment is considered highly acceptable among patient populations and can be delivered successfully in groups. The above factors imply that CFT is a highly viable option for addressing the more psychologically pervasive symptoms of CPTSD. Future research in this population would benefit from evaluating the efficacy of a CFT intervention for emotional stabilisation prior to a trauma-processing intervention, versus a trauma-processing intervention alone. Consensus between the ICD-11 and DSM-5 regarding diagnostic criteria will be a major step in understanding these research outcomes. More high-quality and large-scale research investigating CFT as a primary intervention for individuals with CPTSD is a worthwhile movement towards an agreed treatment protocol for the disorder.

## Declarations

**Conflict of Interest** The authors whose names are listed declare that they have no affiliations with or involvement in any organization or entity with any financial interest (such as honoraria; educational grants; participation in speakers' bureaus; membership, employment, consultancies, stock ownership, or other equity interest; and expert testimony or patent-licensing arrangements), and there is no conflict of interest.

**Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

## References

- American Psychiatric Association. (2013). Anxiety Disorders. In *Diagnostic and statistical manual of mental disorders* (5th ed.).
- Au, T. M., Sauer-Zavala, S., King, M. W., Petrocchi, N., Barlow, D. H., & Litz, B. T. (2017). Compassion-based therapy for trauma-related shame and posttraumatic stress: Initial evaluation using a multiple baseline design. *Behavior Therapy, 48*(2), 207–221. <https://doi.org/10.1016/j.beth.2016.11.012>

- Barlow, M. R., Turow, R. E. G., & Gerhart, J. (2017). Trauma appraisals, emotion regulation difficulties, and self-compassion predict posttraumatic stress symptoms following childhood abuse. *Child Abuse & Neglect*, 65, 37–47. <https://doi.org/10.1016/j.chiabu.2017.01.006>
- Beaumont, E. A., Durkin, M., McAndrew, S. L., & Martin, C. (2016). Using compassion focused therapy as an adjunct to trauma-focused CBT for fire service personnel suffering with symptoms of trauma. *Cognitive Behaviour Therapist*, 9, 34. <https://doi.org/10.1017/S1754470X16000209>
- Beaumont, E., Galpin, A., & Jenkins, P. (2012). 'Being kinder to myself': A prospective comparative study, exploring post-trauma therapy outcome measures, for two groups of clients, receiving either Cognitive Behaviour Therapy or Cognitive Behaviour Therapy and Compassionate Mind Training. *Counselling Psychology Review*, 27(1), 31–43. Retrieved from <https://www.bps.org.uk/publications/counselling-psychology-review>
- Ben-Ezra, M., Karatzias, T., Hyland, P., Brewin, C. R., Cloitre, M., Bisson, J. I., & Shevlin, M. (2018). Post-traumatic stress disorder (PTSD) and complex PTSD (CPTSD) as per ICD-11 proposals: A population study in Israel. *Depression and Anxiety*, 35(3), 264–274. <https://doi.org/10.1002/da.22723>
- Bistricky, S. L., Gallagher, M. W., Roberts, C. M., Ferris, L., Gonzalez, A. J., & Wetterneck, C. T. (2017). Frequency of interpersonal trauma types, avoidant attachment, self-compassion, and interpersonal competence: A model of persisting posttraumatic symptoms. *Journal of Aggression Maltreatment & Trauma*, 26(6), 608–625. <https://doi.org/10.1080/10926771.2017.1322657>
- Braehler, C., Gumley, A., Harper, J., Wallace, S., Norrie, J., & Gilbert, P. (2013). Exploring change processes in compassion focused therapy in psychosis: Results of a feasibility randomized controlled trial. *British Journal of Clinical Psychology*, 52(2), 199–214. <https://doi.org/10.1111/bjc.12009>
- Burns, E. E., Jackson, J. L., & Harding, H. G. (2010). Child maltreatment, emotion regulation, and posttraumatic stress: The impact of emotional abuse. *Journal of Aggression Maltreatment & Trauma*, 19(8), 801–819. <https://doi.org/10.1080/10926771.2010.522947>
- Carlyle, M., Rockliff, H., Edwards, R., Ene, C., Karl, A., Marsh, B. ... Morgan, C. J. (2019). Investigating the feasibility of brief compassion focused therapy in individuals in treatment for opioid use disorder. *Substance Abuse: Research and Treatment*, 13, 1–9. <https://doi.org/10.1177/1178221819836726>
- Chetty, S., Friedman, A. R., Taravosh-Lahn, K., Kirby, E. D., Mirescu, C., Guo, F. ... Tsai, M. K. (2014). Stress and glucocorticoids promote oligodendrogenesis in the adult hippocampus. *Molecular Psychiatry*, 19(12), 1275–1283. Retrieved from <https://www.nature.com/articles/mp2013190>. Accessed 01/10/2020.
- Clapton, N. E., Williams, J., Griffith, G. M., & Jones, R. S. (2018). 'Finding the person you really are... on the inside': compassion focused therapy for adults with intellectual disabilities. *Journal of Intellectual Disabilities*, 22(2), 135–153. <https://doi.org/10.1177/1744629516688581>
- Cloitre, M., Brewin, C. R., Bisson, J. I., Hyland, P., Karatzias, T., Lueger-Schuster, B., ... , Shevlin, M. (2020). Evidence for the coherence and integrity of the complex PTSD (CPTSD) diagnosis: response to Achterhof et al., (2019) and Ford (2020). <https://doi.org/10.1080/20008198.2020.1739873>
- Cloitre, M., Courtois, C. A., Charuvastra, A., Carapezza, R., Stolbach, B. C., & Green, B. L. (2011). Treatment of complex PTSD: Results of the ISTSS expert clinician survey on best practices. *Journal of Traumatic Stress*, 24(6), 615–627. <https://doi.org/10.1002/jts.20697>
- Cloitre, M., Courtois, C. A., Ford, J. D., Green, B. L., Alexander, P., Briere, J., & Van der Hart, O. (2012). The ISTSS expert consensus treatment guidelines for complex PTSD in adults. Retrieved from <https://www.researchgate.net/profile/Stephen-Cheung-4/post/What-are-some-practical-ideas-for-providing-trauma-informed-services/attachment/59d63b37c49f478072ea716b/AS%3A273739588014080%401442276011973/download/ComplexPTSD.pdf>. Accessed 01/10/2020.
- Cloitre, M., Garvert, D. W., Brewin, C. R., Bryant, R. A., & Maercker, A. (2013). Evidence for proposed ICD-11 PTSD and complex PTSD: A latent profile analysis. *European Journal of Psychotraumatology*, 4(1), 20706. <https://doi.org/10.3402/ejpt.v4i0.20706>
- Cloitre, M., Hyland, P., Bisson, J. I., Brewin, C. R., Roberts, N. P., Karatzias, T., & Shevlin, M. (2019). ICD-11 posttraumatic stress disorder and complex posttraumatic stress disorder in the United States: A population-based study. *Journal of Traumatic Stress*, 32(6), 833–842. <https://doi.org/10.1002/jts.22454>
- Cloitre, M., Stolbach, B. C., Herman, J. L., Kolk, B. V. D., Pynoos, R., Wang, J., & Petkova, E. (2009). A developmental approach to complex PTSD: Childhood and adult cumulative trauma as predictors of symptom complexity. *Journal of Traumatic Stress*, 22(5), 399–408. <https://doi.org/10.1002/jts.20444>
- Collins, R. N., Gilligan, L. J., & Poz, R. (2018). The evaluation of a compassion-focused therapy group for couples experiencing a dementia diagnosis. *Clinical Gerontologist*, 41(5), 474–486. <https://doi.org/10.1080/07317115.2017.1397830>
- Craig, C., Hiskey, S., & Spector, A. (2020). Compassion Focused Therapy: a systematic review of its effectiveness and acceptability in clinical populations. *Expert Review of Neurotherapeutics*, 20(4), 385–400. <https://doi.org/10.1080/14737175.2020.1746184>



- Cuppige, J., Baird, K., Gibson, J., Booth, R., & Hevey, D. (2018). Compassion focused therapy: Exploring the effectiveness with a transdiagnostic group and potential processes of change. *British Journal of Clinical Psychology, 57*(2), 240–254. <https://doi.org/10.1111/bjc.12162>
- Duarte, C., Pinto-Gouveia, J., & Stubbs, R. J. (2017). Compassionate Attention and Regulation of Eating Behaviour: A pilot study of a brief low-intensity intervention for binge eating. *Clinical Psychology & Psychotherapy, 24*(6), 1437–1447. <https://doi.org/10.1002/cpp.2094>
- Ehlers, A., & Clark, D. M. (2000). A cognitive model of posttraumatic stress disorder. *Behaviour Research and Therapy, 38*(4), 319–345. [https://doi.org/10.1016/S0005-7967\(99\)00123-0](https://doi.org/10.1016/S0005-7967(99)00123-0)
- Feliu-Soler, A., Pascual, J. C., Elices, M., Martín-Blanco, A., Carmona, C., Cebolla, A., & Soler, J. (2017). Fostering self-compassion and loving-kindness in patients with borderline personality disorder: A randomized pilot study. *Clinical Psychology & Psychotherapy, 24*(1), 278–286. <https://doi.org/10.1002/cpp.2000>
- Ferrari, M., Hunt, C., Harrysunker, A., Abbott, M. J., Beath, A. P., & Einstein, D. A. (2019). Self-compassion interventions and psychological outcomes: A meta-analysis of RCTs. *Mindfulness, 10*(8), 1455–1473. <https://doi.org/10.1007/s12671-019-01134-6>
- Gharraee, B., Tajrishi, K. Z., Farani, A. R., Bolhari, J., & Farahani, H. (2018). A randomized controlled trial of compassion focused therapy for social anxiety disorder. *Iranian Journal of Psychiatry and Behavioral Sciences, 12*(4). <https://doi.org/10.5812/ijpbs.80945>
- Gilbert, P. (2009). Introducing compassion-focused therapy. *Advances in Psychiatric Treatment, 15*(3), 199–208. <https://doi.org/10.1192/apt.bp.107.005264>
- Gilbert, P. (2010). An introduction to compassion focused therapy in cognitive behaviour therapy. *International Journal of Cognitive Therapy, 3*(2), 97–112. <https://doi.org/10.1521/ijct.2010.3.2.97>
- Gilbert, P., & Irons, C. (2004). A pilot exploration of the use of compassionate images in a group of self-critical people. *Memory (Hove, England), 12*(4), 507–516. <https://doi.org/10.1080/09658210444000115>
- Gradus, J. L. (2007). Epidemiology of PTSD. *National Center for PTSD (United States Department of Veterans Affairs)*. Retrieved from [https://www.mentalhealth.va.gov/coe/cih-visn2/Documents/Provider\\_Education\\_Handouts/Epidemiology\\_of\\_PSTD\\_Version\\_3.pdf](https://www.mentalhealth.va.gov/coe/cih-visn2/Documents/Provider_Education_Handouts/Epidemiology_of_PSTD_Version_3.pdf). Accessed 01/10/2020.
- Graser, J., Höfling, V., Weßlau, C., Mendes, A., & Stangier, U. (2016). Effects of a 12-week mindfulness, compassion, and loving kindness program on chronic depression: A pilot within-subjects wait-list controlled trial. *Journal of Cognitive Psychotherapy, 30*(1), 35–49. <https://doi.org/10.1891/0889-8391.30.1.35>
- Hansen, M., Hyland, P., Karstoft, K. I., Vaegter, H. B., Bramsen, R. H., Nielsen, A. B., & Andersen, T. E. (2017). Does size really matter? A multisite study assessing the latent structure of the proposed ICD-11 and DSM-5 diagnostic criteria for PTSD. *European Journal of Psychotraumatology, 8*(sup7), 1398002. <https://doi.org/10.1080/20008198.2017.1398002>
- Held, P., & Owens, G. P. (2015). Effects of self-compassion workbook training on trauma-related guilt in a sample of homeless veterans: A pilot study. *Journal of Clinical Psychology, 71*(6), 513–526. <https://doi.org/10.1002/jclp.22170>
- Held, P., Owens, G. P., Thomas, E. A., White, B. A., & Anderson, S. E. (2018). A pilot study of brief self-compassion training with individuals in substance use disorder treatment. *Traumatology, 24*(3), 219. <https://doi.org/10.1037/trm0000146>
- Heriot-Maitland, C., Vidal, J. B., Ball, S., & Irons, C. (2014). A compassionate-focused therapy group approach for acute inpatients: Feasibility, initial pilot outcome data, and recommendations. *British Journal of Clinical Psychology, 53*(1), 78–94. <https://doi.org/10.1111/bjc.12040>
- Herman, J. (2012). CPTSD is a distinct entity: Comment on Resick et al.(2012). *Journal of Traumatic Stress, 25*(3), 256–257. <https://doi.org/10.1002/jts.21697>
- Herman, J. L. (1992). Complex PTSD: A syndrome in survivors of prolonged and repeated trauma. *Journal of Traumatic Stress, 5*(3), 377–391. <https://doi.org/10.1002/jts.2490050305>
- Hoffart, A., Øktedalen, T., & Langkaas, T. F. (2015). Self-compassion influences PTSD symptoms in the process of change in trauma-focused cognitive-behavioral therapies: a study of within-person processes. *Frontiers in Psychology, 6*, 1273. <https://doi.org/10.3389/fpsyg.2015.01273>
- Hyland, P., Murphy, J., Shevlin, M., Vallières, F., McElroy, E., & Cloitre, M. (2017). Variation in post-traumatic response: The role of trauma type in predicting ICD-11 PTSD and CPTSD symptoms. *Social Psychiatry and Psychiatric Epidemiology, 52*(6), 727–736. <https://doi.org/10.1007/s00127-017-1350-8>
- Jonas, D. E., Cusack, K., Forneris, C. A., Wilkins, T. M., Sonis, J., Middleton, J. C., ... , Gaynes, B. N. (2013). Psychological and pharmacological treatments for adults with posttraumatic stress disorder (PTSD). Retrieved from <https://pubmed.ncbi.nlm.nih.gov/23658937/>. Accessed 01/10/2020.

- Karatzias, T., Hyland, P., Bradley, A., Fyvie, C., Logan, K., Easton, P., & Cloitre, M. (2019a). Is self-compassion a worthwhile therapeutic target for ICD-11 Complex PTSD (CPTSD)? *Behavioural and Cognitive Psychotherapy*, 47(3), 257–269. <https://doi.org/10.1017/S1352465818000577>
- Karatzias, T., Murphy, P., Cloitre, M., Bisson, J., Shevlin, M., Hyland, P., & Bradley, A. (2019b). Psychological interventions for ICD-11 complex PTSD symptoms: systematic review and meta-analysis. *Psychological Medicine*, 49(11), 1761–1775. <https://doi.org/10.1017/S0033291719000436>
- Karatzias, T., Shevlin, M., Fyvie, C., Hyland, P., Efthymiadou, E., Wilson, D., & Cloitre, M. (2017). Evidence of distinct profiles of posttraumatic stress disorder (PTSD) and complex posttraumatic stress disorder (CPTSD) based on the new ICD-11 trauma questionnaire (ICD-TQ). *Journal of Affective Disorders*, 207, 181–187. <https://doi.org/10.1016/j.jad.2016.09.032>
- Kearney, D. J., Malte, C. A., McManus, C., Martinez, M. E., Felleman, B., & Simpson, T. L. (2013). Loving-kindness meditation for posttraumatic stress disorder: A pilot study. *Journal of Traumatic Stress*, 26(4), 426–434. <https://doi.org/10.1002/jts.21832>
- Kirby, J. N. (2017). Compassion interventions: The programmes, the evidence, and implications for research and practice. *Psychology and Psychotherapy: Theory Research and Practice*, 90(3), 432–455. <https://doi.org/10.1111/papt.12104>
- Kirby, J. N., Tellegen, C. L., & Steindl, S. R. (2017). A meta-analysis of compassion-based interventions: Current state of knowledge and future directions. *Behavior Therapy*, 48(6), 778–792. <https://doi.org/10.1016/j.beth.2017.06.003>
- Laithwaite, H., O'Hanlon, M., Collins, P., Doyle, P., Abraham, L., Porter, S., & Gumley, A. (2009). Recovery after psychosis (RAP): A compassion focused programme for individuals residing in high security settings. *Behavioural and Cognitive Psychotherapy*, 37(5), 511. <https://doi.org/10.1017/S1352465809990233>
- Lang, A. J., Casmar, P., Hurst, S., Harrison, T., Golshan, S., Good, R., & Negi, L. (2017). Compassion meditation for veterans with posttraumatic stress disorder (PTSD): A nonrandomized study. *Mindfulness*, 11(63), 74. <https://doi.org/10.1007/s12671-017-0866-z>
- Lee, D. A., Scragg, P., & Turner, S. (2001). The role of shame and guilt in traumatic events: A clinical model of shame-based and guilt-based PTSD. *British Journal of Medical Psychology*, 74(4), 451–466. <https://doi.org/10.1348/000711201161109>
- Lee, W. K., & Bang, H. J. (2010). The effects of mindfulness-based group intervention on the mental health of middle-aged Korean women in community. *Stress and Health*, 26(4), 341–348. <https://doi.org/10.1002/smi.1303>
- Maercker, A., Brewin, C. R., Bryant, R. A., Cloitre, M., van Ommeren, M., Jones, L. M., & Somasundaram, D. J. (2013). Diagnosis and classification of disorders specifically associated with stress: proposals for ICD-11. *World Psychiatry*, 12(3), 198–206. <https://doi.org/10.1002/wps.20057>
- McManus, J., Tsvivos, Z., Woodward, S., Fraser, J., & Hartwell, R. (2018). Compassion focused therapy groups: evidence from routine clinical practice. *Behaviour Change*, 35(3), 167–173. <https://doi.org/10.1017/bec.2018.16>
- Miron, L. R., Sherrill, A. M., & Orcutt, H. K. (2015). Fear of self-compassion and psychological inflexibility interact to predict PTSD symptom severity. *Journal of Contextual Behavioral Science*, 4(1), 37–41. <https://doi.org/10.1016/j.jcbs.2014.10.003>
- Moore, G. F., Audrey, S., Barker, M., Bond, L., Bonell, C., Hardeman, W., ... , Baird, J. (2015). Process evaluation of complex interventions: Medical Research Council guidance. *BMJ*, 350. <https://doi.org/10.1136/bmj.h1258>
- Müller-Engelmann, M., Schreiber, C., Kümmerle, S., Heidenreich, T., Stangier, U., & Steil, R. (2019). A trauma-adapted mindfulness and loving-kindness intervention for patients with PTSD after interpersonal violence: a multiple-baseline study. *Mindfulness*, 10(6), 1105–1123. <https://doi.org/10.1007/s12671-018-1068-z>
- Navab, M., Dehghani, A., & Salehi, M. (2019). Effect of compassion-focused group therapy on psychological symptoms in mothers of attention-deficit hyperactivity disorder children: A pilot study. *Counselling and Psychotherapy Research*, 19(2), 149–157. <https://doi.org/10.1002/capr.12212>
- Neff, K. D., & Germer, C. K. (2013). A pilot study and randomized controlled trial of the mindful self-compassion program. *Journal of Clinical Psychology*, 69(1), 28–44. <https://doi.org/10.1002/jclp.21923>
- Noorbala, F., Borjali, A., Ahmadian-Attari, M. M., & Noorbala, A. A. (2013). Effectiveness of compassionate mind training on depression, anxiety, and self-criticism in a group of Iranian depressed patients. *Iranian Journal of Psychiatry*, 8(3), 113. Retrieved from <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3887227/>. Accessed 01/10/2020.
- Perkonig, A., Höfler, M., Cloitre, M., Wittchen, H. U., Trautmann, S., & Maercker, A. (2016). Evidence for two different ICD-11 posttraumatic stress disorders in a community sample of adolescents

- and young adults. *European Archives of Psychiatry and Clinical Neuroscience*, 266(4), 317–328. <https://doi.org/10.1007/s00406-015-0639-4>
- Readings, J. (2016). *Evaluation of the Effectiveness and Acceptability of a Phase-Based Treatment for Complex Post-Traumatic Stress Disorder*. Doctoral dissertation, Royal Holloway, University of London. Retrieved from <https://pure.royalholloway.ac.uk/portal/files/26909178/2016JenniferReadingsDClinPsy.pdf>
- Resick, P. A., Bovin, M. J., Calloway, A. L., Dick, A. M., King, M. W., Mitchell, K. S., & Wolf, E. J. (2012). A critical evaluation of the complex PTSD literature: Implications for DSM-5. *Journal of Traumatic Stress*, 25(3), 241–251. <https://doi.org/10.1002/jts.21699>
- Rose, A., McIntyre, R., & Rimes, K. A. (2018). Compassion-focused intervention for highly self-critical individuals: Pilot study. *Behavioural and Cognitive Psychotherapy*, 46(5), 583–600. <https://doi.org/10.1017/S135246581800036X>
- Sachser, C., Keller, F., & Goldbeck, L. (2017). Complex PTSD as proposed for ICD-11: Validation of a new disorder in children and adolescents and their response to Trauma-Focused Cognitive Behavioral Therapy. *Journal of Child Psychology and Psychiatry*, 58(2), 160–168. <https://doi.org/10.1111/jcpp.12640>
- Safer, D. L., & Hugo, E. M. (2006). Designing a control for a behavioural group therapy. *Behaviour Therapy*, 37(2), 120–130. <https://doi.org/10.1016/j.beth.2005.06.001>
- Shahar, B., Szepeswol, O., Zilcha-Mano, S., Haim, N., Zamir, O., Levi-Yeshuvi, S., & Levit-Binnun, N. (2015). A wait-list randomized controlled trial of loving-kindness meditation programme for self-criticism. *Clinical Psychology & Psychotherapy*, 22(4), 346–356. <https://doi.org/10.1002/cpp.1893>
- Shevlin, M., Hyland, P., Roberts, N. P., Bisson, J. I., Brewin, C. R., & Cloitre, M. (2018). A psychometric assessment of disturbances in self-organization symptom indicators for ICD-11 complex PTSD using the International Trauma Questionnaire. *European Journal of Psychotraumatology*, 9(1), 1419749. <https://doi.org/10.1080/20008198.2017.1419749>
- Shipman, K., Edwards, A., Brown, A., Swisher, L., & Jennings, E. (2005). Managing emotion in a maltreating context: A pilot study examining child neglect. *Child Abuse & Neglect*, 29(9), 1015–1029. <https://doi.org/10.1016/j.chiabu.2005.01.006>
- Sommers-Spijkerman, M. P. J., Trompetter, H. R., Schreurs, K. M. G., & Bohlmeijer, E. T. (2018). Compassion-focused therapy as guided self-help for enhancing public mental health: A randomized controlled trial. *Journal of Consulting and Clinical Psychology*, 86(2), 101. <https://doi.org/10.1037/ccp0000268>
- Stein, D. J., McLaughlin, K. A., Koenen, K. C., Atwoli, L., Friedman, M. J., Hill, E. D., & Alonso, J. (2014). DSM-5 and ICD-11 definitions of posttraumatic stress disorder: Investigating “narrow” and “broad” approaches. *Depression and Anxiety*, 31(6), 494–505.
- Thompson, B. L., & Waltz, J. (2008). Self-compassion and PTSD symptom severity. *Journal of Traumatic Stress: Official Publication of The International Society for Traumatic Stress Studies*, 21(6), 556–558. <https://doi.org/10.1002/da.22279>
- Van de Mortel, T. F. (2008). Faking it: social desirability response bias in self-report research. *Australian Journal of Advanced Nursing*, 25(4), 40. Retrieved from <https://doi.org/10.3316/informit.210155003844269>
- Van der Kolk, B. A., Spinazzola, J., Blaustein, M. E., Hopper, J. W., Hopper, E. K., Korn, D. L., & Simpson, W. B. (2007). A randomized clinical trial of eye movement desensitization and reprocessing (EMDR), fluoxetine, and pill placebo in the treatment of posttraumatic stress disorder: treatment effects and long-term maintenance. *Journal of Clinical Psychiatry*, 68(1), 37. Retrieved from <http://citeseerx.ist.psu.edu/viewdoc/download?doi=10.1.1.476.4096&rep=rep1&type=pdf>
- Vogel, M., Braungardt, T., Kaul, S., & Schneider, W. (2017). Attrition and outcome in group psychotherapy among traumatized and non-traumatized inpatients. *Journal of Psychology & Psychotherapy*, 7(3), 2161–0487. <https://doi.org/10.4172/2161-0487.1000301>
- Watkins, L. E., Sprang, K. R., & Rothbaum, B. O. (2018). Treating PTSD: A review of evidence-based psychotherapy interventions. *Frontiers in Behavioral Neuroscience*, 12, 258. <https://doi.org/10.3389/fnbeh.2018.00258>
- Wilson, A. C., Mackintosh, K., Power, K., & Chan, S. W. (2019). Effectiveness of self-compassion related therapies: A systematic review and meta-analysis. *Mindfulness*, 10(6), 979–995. <https://doi.org/10.1007/s12671-018-1037-6>
- Winders, S. J., Murphy, O., Looney, K., & O'Reilly, G. (2020). Self-compassion, trauma, and posttraumatic stress disorder: A systematic review. *Clinical Psychology & Psychotherapy*, 27(3), 300–329. <https://doi.org/10.1002/cpp.2429>
- World Health Organization. (2018). *International Statistical Classification of Diseases and Related Health Problems* (11th ed, ICD-11), Retrieved from <https://www.who.int/classifications/classification-of-diseases>. Accessed 01/10/2020.

Zuj, D. V., Palmer, M. A., Lommen, M. J., & Felmingham, K. L. (2016). The centrality of fear extinction in linking risk factors to PTSD: a narrative review. *Neuroscience & Biobehavioral Reviews*, *69*, 15–35. <https://doi.org/10.1016/j.neubiorev.2016.07.014>

**Publisher's Note** Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.