RESEARCH Open Access

Check for updates

The gender gap in adolescents' emotional and behavioural problems in Georgia: a cross-sectional study using Achenbach's Youth Self Report

Khatia Antia^{1,2*}, Justina Račaitė³, Genė Šurkienė³ and Volker Winkler¹

Abstract

Background Adolescents, particularly girls, are vulnerable to mental health disorders. Knowledge about young people's mental health in Eastern European countries is limited. This study is the first to investigate adolescents' self-reported emotional and behavioural problems in Georgia from a public mental health perspective.

Methods This study utilized Achenbach's Youth Self-Reported syndrome scales among 933 adolescents studying from grades 7–12 in 18 public schools in Georgia. We compared the gender-specific results with each other and with the Achenbach's Normative Sample using two-sample t-tests. Linear regression was used to assess associations between internalizing and externalizing problems and individual as well as demographic characteristics such as parental migration experience (being 'left-behind' or 'staying behind').

Results The study found that girls obtained higher scores than boys in Youth Self-Reported empirical syndrome scales and the internalizing broadband scale. Rule-breaking behaviour was the only syndrome scale where boys scored higher. Compared to Achenbach's Normative Sample, adolescents in Georgia scored higher on all scales. Regression analyses showed that illnesses, having fewer than three close friends, problems at school, and worse relationships with peers/siblings/parents (compared to peers) were associated with higher internalizing and externalizing problems scores in both genders. Performing household chores, living with only one parent, or having a migrant parent showed no associations in either gender.

Conclusion Emotional and behavioural difficulties of adolescents in Georgia, especially girls, require attention. Having close friends, developing strong relationships with family members, and a supportive school environment could help mitigate emotional and behavioural problems among adolescents in Georgia.

Keywords Adolescents, Gender gap, Mental health, Emotional difficulties, Problem behaviour, Self-report, Left-behind, Georgia



^{*}Correspondence: Khatia Antia khatia.antia@uni-heidelberg.de

¹Heidelberg Institute of Global Health, University Hospital Heidelberg, Im Neuenheimer Feld 130.3, 69120 Heidelberg, Germany

²School of Health Sciences, The University of Georgia, 77a Kostava Street, Tbilisi 0175, Georgia

³Department of Public Health, Institute of Health Sciences, Faculty of Medicine, Vilnius University, M. K. Čiurlionio str. 21, Vilnius LT-03101, Lithuania

Background

Adolescence is a transitional period into adulthood, a unique time essential for human development [1, 2]. Increased stress, anxiety, and behavioural changes make this period challenging for youth [2, 3]. Available literature suggests that around 50% of lifetime mental illnesses develop during the teenage years [4]. Adolescents with poor mental health may experience personal, academic and social challenges [5].

Various outcome measures are applied to examine the mental health of children and adolescents [6], among which internalizing and externalizing are the most commonly used broadband self-reported measures [7]. When investigating internalizing and externalizing problems, scholars observe a universal gender gap, specifically, worse internalizing outcomes among girls and worse externalizing outcomes among boys [8–14]. Moreover, the risk of developing depression and anxiety is twice as high in adolescent girls as in boys [7, 15–17]. Evidence from European studies suggest that older adolescents are worse off than younger ones [8–14].

Georgia is an Eastern European, post-Soviet country and one of the migrant-sending countries in the region [18]. In the aftermath of the Soviet Union's collapse, the population rapidly decreased, from 5.4 million in 1989 to 3.7 million in 2022, mainly due to international labour migration [19, 20]. As a result, more than 55% of migrants are now comprised of women, and approximately 40% are children and adolescents who may be considered as/ called 'left-behind' or 'staying behind', denoting that they live with caregivers while their parent(s) work abroad [18]. Existing literature suggests that left-behind children and adolescents are particularly vulnerable to mental illnesses [21-24]. Following the rise of parental out-migration in Georgia, the percentage of youth and adults with mental health disorders has increased five times (from 4.0 to 2010 to 20.0 per 100,000 in 2021). This is while in children below the age of fourteen, the prevalence increased from 0.4 to 2010 to 3.0 per 100,000 children in 2021, which is nearly an 8-fold increase [25].

The study aims to examine the emotional and behavioural difficulties of adolescents in Georgia from a public mental health perspective. To our knowledge, this is the first study using the Achenbach System of Empirically Based Assessment (ASEBA) Youth Self-Report (YSR) in Georgia. A better understanding of problem behaviours in the context of gender, parental migration and other demographic factors of adolescents in Georgia would in turn facilitate identifying mental health needs and developing more targeted interventions.

Methods

Study design and data

In this school-based cross-sectional study, we collected data from October 2021 to January 2022. We purposively invited 22 public schools from urban and rural areas in the Samegrelo-Zemo Svaneti region to participate in the study. All except one school principal agreed to partake in the study and three schools were closed during the data collection phase due to increased COVID-19 cases. We selected the Samegrelo-Zemo Svaneti region as it is located in the Central, North-Western part of Georgia and borders the Black Sea, Abkhazia and Russia [26]. With 10.8% of the country's territory and 301 thousand residents, Samegrelo-Zemo Svaneti is the second-largest region in Georgia [26].

In each school, we invited all students from grades 7 to 12 (ages 12–18) to participate in our study. We arranged meetings with class tutors, asking for their assistance before and during the data collection process. The tutors assisted with arranging meetings and collecting consent forms from students and parents/caregivers. After explaining the questionnaire and the scoring method/guidelines, the data collection team (the first author and a research assistant) surveyed all students whose parents had given consent, in classrooms during/after regular school hours. Each survey took around 30–45 minutes; Data collection was anonymous; such that neither the school administration nor the tutors had access to the responses/results.

The questionnaire consisted of the Youth Self Report (YSR) of the ASEBA System supplemented with a few additional questions (listed in Appendix A1) to obtain socio-demographic information, such as the number of siblings and household members, as well as parental migration experience [27]. The YSR is a widely used child-report measure for adolescents aged 11–18 that assesses problem behaviours along two broadband scales: internalizing and externalizing, with 117 items classified as problem items or socially desirable items (listed in Appendix A2), each scored: 0=not true, 1=somewhat true, or 2=very true/often true [28].

The ASEBA manual compares multicultural norm scores with borderline and clinical cut-off scores for children living in different societies [28]. For societies/countries without a norm score, such as Georgia, it is recommended to use the ASEBA Standard that equals Group 2 (societies with medium problem scores) [28, 29].

The questionnaire (ASEBA Licence N2186-10-06-20) as well as consent forms were in Georgian language and contained the contact details of the first author in case of questions/ queries. The study was approved by the Ethics Committee of the Medical Faculty of Heidelberg University N- S-160/2021 and by the Ethics Committee of the University of Georgia N-11-12401.

Statistical analysis

Following descriptive analysis, we compared the YSR scores of the empirical syndrome items, the broadband scales between boys and girls, and the ASEBA Standard separated by gender, using two-sample t-tests.

We then, modelled the broadband scales using linear and multiple regression to identify associated variables for girls and boys, separately. The following independent variables were available for analysis: age groups (12–13, 14-15, 16-18 years); performing in any sports; number of siblings; doing household chores; having an illness (in general); the number of close friends; having school problems; living with mother, father or both parents; relationship with peers/siblings/parents; and the parents' migration status. The variables assessing the 'relationship with peers, siblings and parents' asks for 'worse, likewise or better' relationships compared to peers. Regarding parental migration, we grouped adolescents into nonleft-behind (not having a migrant parent) and left-behind (having at least one migrant parent). After bivariable analysis and collinearity check, we included all independent variables in the multiple linear regression analysis. For the statistical analysis, we used Stata version IC 15 and considered p<0.05 as significant.

Results

All 2,495 students between grades 7 and 12 from 18 Public schools (11 rural, 7 urban) were invited to partake in the study, of which 933 (37.9%) agreed to participate and filled out the questionnaire. The responses varied considerably across schools, ranging from 10.7 to 70.7%.

Table 1 School characteristics (numbers of students, location, and participation)

School N°	N of Students (grade 7 to 12)	Location	Participation (%)
1	56	rural	64.3
2	72	rural	41.7
3	256	urban	70.7
4	61	rural	21.3
5	79	rural	35.4
6	112	urban	22.3
7	94	rural	39.4
8	36	rural	19.4
9	203	urban	54.2
10	148	urban	33.8
11	58	rural	46.6
12	297	urban	48.5
13	63	rural	42.9
14	89	rural	38.2
15	86	urban	68.1
16	81	rural	39.5
17	527	urban	14.6
18	177	rural	10.7
Total	2 495	-	37.4

Table 1 describes the detailed characteristics of participant schools.

Among the participants, 50.4% were girls, and 45.4% were boys, with the remaining 4.2% not answering the question about their gender. Most participants reported having no illness (90%), performing sports (89.3%) and stated having at least three close friends (86.7%). Team sports such as football, basketball, volleyball, and table tennis were the most reported sports activities (87.2%). Household chores such as cleaning the house and looking after siblings were done by 22.2%. Almost 90% of adolescents acknowledged having hobbies, such as reading, dancing, listening to and making music, playing cards and computer games, and making handicrafts. About a quarter admitted facing problems at school. Only 4.4% of the adolescents stated having worse relationships with their peers than other adolescents, while 67.9% claimed better relationships with their parents, and 52.2% stated having better relations with their siblings than their peers. In our sample, 76.4% of the adolescents lived with both parents, while 27.7% indicated having at least one migrant parent. Table 2 presents a detailed description of participant characteristics.

Table 3 shows the comparison of empirical syndrome scales and the broadband scales internalizing and externalizing among boys and girls within our sample and the comparison to the ASEBA Standard. The analysis revealed higher scores among girls for all syndromes except for rule-breaking behaviour, in which boys scored higher. The internalizing score was also higher among girls, while no difference in externalizing score could be observed. Adolescents who did not indicate their gender obtained similar mean scores as boys (internalizing 9.5, externalizing 10.7).

Compared to the ASEBA Normative Sample, the girls and boys in our sample obtained higher scores for all scales, except for the empirical syndromes such as 'somatic complains' and 'thoughts problems' among boys. In broadband internalizing, we observed only a minor difference among boys. Generally, differences in mean scores were noted to be larger for girls than for boys.

All scales remained within the normal range of the ASEBA standard group 2.

Tables 4 and 5 present the results of the multiple regression analyses among girls and boys, respectively. Results of bivariable models are similar and shown in Appendix A4 and A5.

Multiple regression analysis showed associations between girls with higher internalizing scores and not performing any sports, having less than three close friends, experiencing school problems, and reporting worse relationships with peers, siblings, and parents. While having better relationships with peers and parents and not having an illness were associated with lower

Table 2 Characteristics of the participants (n = 933)

Table 2 Characteristics of the participant		
Variable	N	%
Gender		
Girl	470	50.4
Boy	424	45.4
Missing	39	4.2
Age group		
12–13	359	38.5
14–15	336	36.0
16–18	238	25.5
Performing any sport		
Yes	833	89.3
No	96	10.3
Missing	4	0.4
Number of siblings		
0	159	17.0
1–3	399	42.8
4–5	25	2.7
≥6	5	0.5
Missing	345	37.0
Doing chores		
Yes	207	22.2
No	717	76.8
Missing	9	1.0
Having an illness		
Yes	64	6.9
No	840	90.0
Missing	29	3.1
Number of close friends		
0	24	2.6
1–3	353	37.9
≥4	522	55.9
Missing	34	3.6
Having school problems		
Yes	237	25.4
No	643	68.9
Missing	53	5.7
Living with	33	5.7
Mother only	84	9.0
Father only	37	13.0
Both	712	76.3
Missing	100	10.7
Relationship with peers (compared to peers)	100	10.7
Worse	41	4.4
Likewise	396	42.4
Better	416	44.6
	80	
Missing Relationship with siblings (compared to page)	00	8.6
Relationship with siblings (compared to peers)	27	4.0
Worse	37	4.0
Likewise	268	28.7
Better Do not have siblings	487	52.2
Do not have siblings	97	10.4
Missing	44	4.7
Relationship with parents (compared to peers)	2.1	2.2
Worse	21	2.2

Table 2 (continued)

Variable	N	%
Likewise	192	20.6
Better	633	67.9
Missing	87	9.3
Having at least one migrant parent		
Yes	258	27.7
No	615	65.9
Missing	60	6.4

internalizing scores. Performing in any sport and having relationships with siblings had no association with externalizing scores.

Multiple linear regression analysis for boys revealed associations between higher internalizing scores, having less than three close friends, experiencing school problems, and having worse relationships with peers. In contrast, not having an illness and better relationships with peers were associated with lower internalizing scores. School problems and age bracket of 16–18 years was associated with higher externalizing scores. Additionally, relationships with siblings appeared to be associated with externalizing; however, better/worse relationships and not having siblings showed an association with lower externalizing scores (Table 5).

Discussion

This study examined the emotional and behavioural problems of Georgian schoolchildren using Achenbach's Youth Self Report. While boys expressed more rule-breaking behaviour in our study, girls displayed more problem behaviours and internalizing symptoms, indicating the gender gap. Compared to the ASEBA normative standard, Georgian adolescents in both genders showed more emotional and behavioural problems, with a gap that was even wider among girls. Importantly, factors such as the relationship with peers/siblings/parents, number of close friends and school problems might influence the results.

Our findings on internalizing problems are consistent with evidence from previous studies [2, 30–33]. For example, a comparative study of Campbell, Bann [2] from 73 countries suggested worse internalizing outcomes among girls. Remarkably, the wealthiest countries in Europe with the highest gender equality index show the most significant gender gaps [2]. Irrespective of socioeconomic conditions, teenage girls perceive their health as poorer [30, 31] and express higher levels of anxiety and psychological distress [32, 34–36]. Research, including longitudinal data, demonstrates an increasing trend of emotional problems among girls [33], while a minor increase is observed among European boys [37]. Various theories have been proposed to explain this observation, mostly suggesting individual [34], biological

(puberty-related hormonal changes) [38, 39], social relationships [40] and reporting differences [41]. Some academics have also suggested that shifting educational paradigms towards female education have led teenage girls to experience increased educational pressure and psychological distress [34].

Furthermore, non-differing externalizing problems among Georgian girls and boys in our study align with

tend to be universal, it seems more inconsistent with externalizing.

Similar to our results, other scholars have found that a lack of friendships and relationship difficulties with parents are associated with emotional and behavioural problems among European youths [42]. Conflicts within families were also found to increase the risk of developing mental health disorders among youth in Lithuania

Table 3 Youth Self-Report Emotional and behavioural problems - descriptive statistics for syndrome and broadband scales

						ASEBA Normativ	ve Sample ¹		
Outcome	Girls			Boys		Girls		Boys	
Empirically based syndromes	N	mean (95% CI)	N	mean (95% CI)	p-value ²	mean (normal range ³)	p-value ⁴	mean (normal range)	p-value ⁴
Anxious/ Depressed	370	7.2 (6.7–7.7)	338	3.9 (3.6–4.3)	< 0.001	5.1 (< 12.0)	< 0.001	3.4 (< 9.0)	0.016
Withdrawn/ Depressed	424	5.4 (5.1–5.8)	372	4.0 (3.7–4.3)	< 0.001	3.1 (< 8.0)	< 0.001	2.7 (< 7.0)	< 0.001
Somatic Complains	384	4.7 (4.3-5.0)	362	2.2 (2.0-2.5)	< 0.001	3.4 (< 8.0)	< 0.001	2.3 (< 7.0)	0.658
Social Problems	420	4.1 (3.8–4.4)	359	3.4 (3.1–3.7)	0.002	3.2 (< 8.0)	< 0.001	2.9 (< 8.0)	0.006
Thought Problems	321	5.1 (4.6–5.7)	256	3.2 (2.8–3.6)	< 0.001	3.7 (< 10.0)	< 0.001	3.1 (< 9.0)	0.715
Attention Problems	429	6.3 (6.0-6.6)	367	5.4 (5.1–5.8)	< 0.001	4.6 (< 10.0)	< 0.001	4.9 (< 10.0)	0.014
Rule-Breaking Behaviour	409	3.8 (3.4–4.2)	347	4.7 (4.3–5.1)	0.001	3.5 (< 9.0)	0.228	3.8 (< 10.0)	< 0.001
Aggressive behaviour	424	9.2 (8.7–9.7)	352	7.3 (6.8–7.8)	< 0.001	6.5 (< 14.0)	< 0.001	6.0 (< 13.0)	< 0.001
Other problems	422	4.7 (4.5-5.0)	362	4.0 (3.8–4.3)	< 0.001	-	-	-	-
Broadband scales									
Internalizing	305	16.9 (15.7–18.1)	305	9.8 (9.0-10.6)	< 0.001	11.6 (< 19.0)	< 0.001	8.3 (< 14.0)	0.001
Externalizing	391	13.0 (12.2–13.9)	318	12.0 (11.2–12.8)	0.081	9.9 (< 16.0)	< 0.001	9.8 (< 17.0)	< 0.001

Number of observations, mean score, and p-values for comparisons between girls and boys, as well as to the ASEBA Normative Sample

the American normative standard [29]; however, it contradicts the leading international literature on this issue [8, 34, 36]. For example, a study of more than 27,000 adolescents from 24 countries showed worse externalizing problems among boys [8]. Even though stressful life events seem to intensify externalizing problems in both genders, the increase is higher in boys [36]. To explain this difference, some scholars have suggested that boys are more vulnerable to self-criticism [36]. Another possible explanation is that generally, in self-reports, girls focus more on internalizing, and boys focus on externalizing problems [34]. While gender effects on internalizing

[43]. In contrast, close ties and relationships with one's family members correlates with better mental health and reduced risky behaviour during adolescence [42, 44]. Moreover, close friendships may even prevent and mitigate aggression and antisocial behaviour among adolescent boys [45]. Given that boys in Georgia display higher rule-breaking behaviour, developing friendships and social connections are believed to bear an important impact on the mitigation of this behaviour in this population.

Consistent with our study's findings, other scholars suggest an association between school environment

¹ASEBA Standard – multicultural normative samples group 2 adopted from the Achenbach and Rescorla (2007)

²Comparison between girls and boys from the study sample

³Normal range – given by the ASEBA for group 2

⁴Comparison between ASEBA Normative Sample and the study sample

Table 4 Multiple linear regression using the broadband internalizing and externalizing scales as the outcomes for girls

	Internalizing		Externalizing	
Independent Variable	Coefficient (95% CI)	p-value	Coefficient (95% CI)	p-value
Age group		0.335		0.268
2–13	Ref.		Ref.	
14–15	1.61 (-2.16–5.37)		2.81 (0.05–5.57)	
16–18	-1.52 (-5.33–2.29)		0.19 (-2.71–3.09)	
Performing any Sport		0.008		0.197
res	Ref.		Ref.	
No	3.19 (-1.46–7.85)		-2.64 (-6.04–0.76)	
Number of siblings		0.740		0.329
)	Ref.		Ref.	
1–3	5.00		7.92	
	(-18.41-28.42)		(-4.86-20.71)	
4–5	9.85 (-14.34–34.04)		6.12 (-7.40–19.65)	
≥6	3.19 (-23.44–29.83)		0.22 (-16.55 -16.99)	
Doing Chores		0.126		0.329
Yes	0.12		-0.25	
	(-3.30-3.55)		(-2.79-2.30)	
No	Ref.		Ref.	
Having an illness		< 0.001		0.001
Yes	Ref.		Ref.	
No	-6.17 (-12.79–0.46)		-2.85 (-6.96–1.26)	
Number of Close Friends		< 0.001		0.015
0	10.20 (1.85–18.56)		1.51 (-5.61–8.63)	
1–3	0.74		-0.10	
	(-2.60-4.08)		(-2.63–2.43)	
≥4	Ref.		Ref.	
Having School problems		< 0.001		< 0.001
Yes	4.69		1.24	
	(1.30-8.07)		(-1.26-3.74)	
No	Ref.		Ref.	
Living with		0.075		0.063
Mother only	-2.90 (-7.85–2.06)		0.68 (-2.76–4.13)	
Father only	0.08		-4.45	
	(-8.48–8.64)		(-10.70–1.80)	
Both parents	Ref.		Ref.	
Relationship with peers (compared to peers)		< 0.001		0.007
Worse	6.90 (0.98-14.79)		1.49 (-3.68–6.66)	
Likewise	Ref.		Ref.	
Better	-4.01		-2.16	
Relationship with siblings (compared to peers)	(-7.390.63)	< 0.001	(-4.70–0.37)	0.051
Worse	6.36	\ 0.001	9.74	0.031
AAOI 2C	6.36 (-15.40–28.12)		9.74 (-1.70–21.17)	
Likewise	(-13.40–28.12) Ref.		(-1.70–21.17) Ref.	
Better	0.39		-0.88	
Dette.	(-3.48–4.27)		(-3.68–1.92)	

Table 4 (continued)

	Internalizing		Externalizing	
Independent Variable	Coefficient (95% CI)	p-value	Coefficient (95% CI)	p-value
Do not have siblings	6.03		6.71	
	(-17.48–29.54)		(-6.40–19.82)	
Relationship with parents (compared to peers)		< 0.001		< 0.001
Worse	5.32		10.41	
	(-8.18-18.81)		(1.89-18.93)	
Likewise	Ref.		Ref.	
Better	-7.15		-3.23	
	(-11.412.89)		(-6.290.19)	
Having at least one migrant parent		0.445		0.107
Yes	1.72		2.77	
	(-1.71-5.16)		(0.18-5.37)	
No	Ref.		4.61	
			(1.73-7.49)	
Const.	21.07	0.095	9.65	0.183
	(-3.75-45.90)		(-4.61-23.91)	
Adj. R ²	0.3371		0.1978	

and psychological distress among adolescents [34, 46-48]. School stress is mainly discussed in the context of increased schoolwork pressure, conflicts with schoolteachers, and peer bullying [46]. School problems may increase psychosomatic complaints, anxiety, and psychological distress [34] and even lead adolescents to 'school burnout' [47]. Girls seem more responsible for performing well at school, which may cause an imbalance between increased pressure and available resources (e.g. needed support), eventually leading to stress and anxiety [34, 49]. In contrast, a multi-country study by Bradshaw, Martorano [48] suggests that a supportive school environment and good relationships with classmates are crucial determinants of adolescents' well-being and happiness. Similar to youths from other countries, adolescents in our study could experience problems at school because of schoolwork pressure and conflicts with teachers and classmates. School factors could also explain more significant internalizing problems that was observed among teenage girls in our sample.

In general, the gender gap is described as a cross-cultural phenomenon [3, 16, 50, 51]. In addition to the socio-economic conditions, its size seems to be affected by the area of residence (rural/urban) [50], migration background and ethnicity [3, 16, 50, 51]. Considering the migration profile of Georgia, parental migration was another factor we investigated in this study. We found that adolescents with or without parental migration face similar emotional and behavioural difficulties. Our findings only partially reflect evidence from previous studies in this field from Georgia, where some scholars argue that children of migrant parents are vulnerable to poor mental health [52]; while others suggest that parents' migration is associated with children's improved mental

health and well-being [53–55]. Worse mental health outcomes are indicated among left-behind children in other Eastern European countries [56–58] and internationally [21, 24, 59–62]. Several factors could explain this inconsistency. Our study involves adolescents above 11, while most existing studies include children of all ages (0–18). Younger children could be more vulnerable to migratory separation. Informants, other than adolescents may evaluate parental migration impact more negatively. As Lee, Park [63] suggested, self-reported emotional problems of adolescents are more reliable than caregiver reports. Other factors, such as closeness with a migrant parent, good relationships with family members and social interactions, may mitigate the adverse effects of parental absence and, consequently, explain non-differing problem behaviours among left-behind and non-left-behind adolescents in Georgia and elsewhere.

Less than 40% of targeted youths participated in our study. We believe the possible reasons are threefold: (1) There might have been a COVID-19 pandemic-related fear among the public, as many students were absent in the classroom; (2) In Georgia, completing informed consent forms as part of an official request to participate in a research project is rare - it could be seen as "dangerous"; (3) The YSR questionnaire was anonymous, however, many adolescents and their families might still find the subject too sensitive. Our study is the first attempt to encourage young people, their families, schools and communities to collaborate with scientists on similar research projects in the future.

Several elements in our research provide a different perspective/viewpoint and more insightful conclusions to the body of existing literature. We investigated adolescents' emotional and behavioural problems in Georgia,

Table 5 Multiple linear regression using the broadband internalizing and externalizing scales as the outcomes for boys

	Internalizing		Externalizing	
Independent Variable	Coefficient (95% CI)	p-value	Coefficient (95% CI)	p-value
Age group		0.549		0.003
12–13	Ref.		Ref.	
14–15	-0.99		1.51	
	(-3.49–1.50)		(-1.29-4.32)	
16–18	0.95		2.96	
	(-1.85–3.75)		(-0.33-6.24)	
Performing any Sport		0.093		0.438
Yes	Ref.		Ref.	
No	1.43		-0.83	
	(-2.67-5.54)		(-5.60-3.93)	
Number of Siblings		0.163		0.600
0	Ref.		Ref.	
1–3	-0.26		0.70	
	(-7.10–6.56)		(-8.33-9.72)	
4–5	4.00		7.62	
	(-4.72–12.72)		(-3.11–18.35)	
≥6	-1.84		-2.75	
	(-11.89–8.20)		(-14.94–9.43)	
Doing chores		0.393		0.780
Yes	0.26		-1.90	
163	(-2.44–2.96)		(-5.05 -1.25)	
No	Ref.		Ref.	
Having an illness	nei.	< 0.001	ivei.	0.081
Yes	Ref.	\0.001	Ref.	0.001
No	-4.09 (-8.31–0.13)		-2.26 (-7.23–2.72)	
Number of Close Friends	(-0.51-0.15)	< 0.001	(-7.23-2.72)	0.311
	25.04	< 0.001	20.70	0.511
0	25.91		-20.78	
1 2	(11.17–40.66)		(-37.673.88) -0.36	
1–3	0.40		-0.36 (-3.00–2.27)	
× 4	(-1.92–2.71)			
≥4	Ref.	2 2 2 7	Ref.	0.045
Having school problems		0.007		0.015
Yes	3.64		3.58	
	(1.05–6.23)		(0.69–6.47)	
No	Ref.		Ref.	
Living with		0.395		0.794
Mother only	2.75		-2.83	
	(-1.14–6.64)		(-7.19–1.53)	
Father only	0.10		-1.83	
	(-4.53–4.33)		(-6.99–3.32)	
Both parents	Ref.		Ref.	
Relationship with peers (compared to peers)		< 0.001		0.672
Worse	7.08		7.35	
	(0.01–14.15)		(-1.83–16.53)	
Likewise	Ref.		Ref.	
Better	-1.89		-0.72	
Dettel	(-4.15–0.36)		-0.72 (-3.27–1.83)	
Relationship with siblings	(1.15 0.50)	0.126	(3.27 1.03)	0.024
(compared to peers)		0.120		0.024
Worse	-4.99		-6.59	
	(-16.81–6.84)		(-17.40–4.22)	
Likewise	Ref.		Ref.	

Table 5 (continued)

	Internalizing		Externalizing	
Independent Variable	Coefficient (95% CI)	p-value	Coefficient (95% CI)	p-value
Better	-1.88 (-4.43–0.66)		-1.76 (-4.70–1.17)	
Do not have siblings	-2.50 (-9.62–4.62)		-2.36 (-11.42–6.70)	
Relationship with parents (compared to peers)		0.068		0.050
Worse	-3.61 (-14.59–7.36)		3.99 (-7.21 -15.18)	
Likewise	Ref.		Ref.	
Better	-0.47 (-3.48–2.53)		-1.66 (-5.04–1.72)	
Having at least one migrant parent		0.932		0.449
Yes	0.27 (-2.27–2.80)		4.61 (1.73–7.49)	
No	Ref.		Ref.	
Constant	15.63 (7.07–24.20)	< 0.001	13.15 (2.09–24.22)	0.020
Adj. R ²	0.3177		0.1195	

addressing a major research gap in Eastern Europe. Instead of adults, we asked adolescents to report on their problem behaviours, because they are indispensable informants of their mental health. We applied ASEBA tool, the YSR, which is identified among the 11 most reliable measures in child psychopathology [6] and is increasingly used in research worldwide [8, 9, 28, 29]. The YSR had only been applied in clinical practice in Georgia up until now. Finally, we considered parental migration status, which is a critical sociodemographic factor for Georgia.

This study had several limitations. We collected data during the COVID-19 pandemic when the government of Georgia allowed schools to re-open after three semesters of school closure. Schools were given the autonomy to decide between online or in-classroom teaching. Our initial plan of random sample selection was not possible. The COVID-19 pandemic and school closures could have also influenced the results. All included schools were in the Samegrelo-Zemo Svaneti region; thus, our findings are not representative of other areas or the entire population of Georgia. As stated above, the overall response rate in our study is low and varies across schools. Responses of adolescents who missed school on the day of data collection could differ. Not all adolescents who participated in the study answered all questions. We excluded all observations with missing YSR problem items from the analysis. These study findings are based on information that was self-reported by adolescents. Including other informants such as teachers, parents or caregivers and comparing results could provide a broader picture/ more insights. Lastly, the cross-sectional design of the study does not allow us to conclude on possible causes of emotional and behavioural problems of adolescents in Georgia.

Conclusion

The findings of this study highlight the need to address the emotional and behavioural difficulties that Georgian adolescents face, particularly internalizing problems in girls and rule-breaking behaviour in boys. Our results reflect the increasing trend of child and adolescent mental health disorders in Georgia. Relationships within family, social connections and school environment appear as important contributors. Our findings indicate that social support interventions may protect adolescents from experiencing emotional and behavioural difficulties in Georgia.

Finally, we emphasize the necessity of raising public awareness for child and adolescent mental health issues. It is essential that adolescents, schools and parents understand the potential benefits of participating in similar research projects to ours. We further encourage collaborations between state and non-governmental organizations, child and adolescent mental health professionals, as well as global health scientists in developing/promoting mental health interventions involving adolescents, their families, and communities in Georgia.

List of Abbreviations

YSR Youth Self-Report

ASEBA The Achenbach System of Empirically Based Assessment

SDG Sustainable Development Goals
UNICEF The United Nations Children's Fund
COVID-19 Coronavirus Disease 2019

Supplementary Information

The online version contains supplementary material available at https://doi.org/10.1186/s13034-023-00592-0.

Supplementary Material 1

Acknowledgements

We thank school principals, school personnel and a research assistant Marika Antia, for their support during data collection. We thank Zahra Karimian and Priya Shreedhar for their assistance with editing the manuscript. We are also grateful to all adolescents and parents/caregivers for their consent and participation in the study.

Authors' Contributions

KA, JR, GS, and VW conceptualised and designed the study; KA collected the data; KA and VW analysed the data; KA wrote the main manuscript text; VW supervised the project; JR, GS and VW reviewed and edited the manuscript; KA obtained funding. All authors read and approved the final manuscript.

Funding

This study was funded by Deutscher Akademischer Austauschdienst—German Academic Exchange Service (DAAD) Research Grants—Doctoral Programmes in Germany, 2019/20; grant number: 57440921. For the publication fee, we acknowledge financial support by the Else Kröner-Fresenius-Stiftung within the Heidelberg Graduate School of Global Health, by Deutsche Forschungsgemeinschaft within the funding programme Open Access Publishing, and by Ruprecht-Karls-Universität Heidelberg. Open Access funding enabled and organized by Projekt DEAL.

Data Availability

Collected data is stored in a password-protected hard drive at Heidelberg Institute of Global Health. Datasets generated and/or analysed during the current study are available from the corresponding author on reasonable request.

Declarations

Ethics approval and consent to participate

This study was approved by the Ethics Committee of the Medical Faculty of Heidelberg University N-S-160/2021 and by the Ethics Committee of the University of Georgia N-11-12401. All participants as well as their parents/ caregivers provided written informed consent to participate in the study. Georgian translation of the Youth Self Report (YSR) questionnaire was obtained from the ASEBA - Licence N2186-10-06-20. All study procedures and methods were performed in accordance with the relevant guidelines and regulations of the Declaration of Helsinki.

Consent for publication

Not applicable.

Competing interests

The authors declare that they have no competing interests.

Received: 24 January 2023 / Accepted: 15 March 2023 Published online: 30 March 2023

References

- Stehlik T. Educational philosophy for 21st century teachers. Switzerland: Palgrave Macmillan; 2018.
- Campbell OLK, Bann D, Patalay P. The gender gap in adolescent mental health: a cross-national investigation of 566,829 adolescents across 73 countries. SSM - Population Health. 2021;13:100742.
- Murad SD, Joung IM, van Lenthe FJ, Bengi-Arslan L, Crijnen AA. Predictors of self - reported problem behaviours in turkish immigrant and dutch adolescents in the Netherlands. J Child Psychol Psychiatry. 2003;44(3):412–23.

- Kessler RC, Amminger GP, Aguilar-Gaxiola S, Alonso J, Lee S, Ustün TB. Age of onset of mental disorders: a review of recent literature. Curr Opin Psychiatry. 2007;20(4):359–64.
- European Network of Ombudspersons for Children. Child and Adolescent Mental Health in Europe: ENOC; 2018 [cited 2022 19 Dec]. Available from: https://enoc.eu/wp-content/uploads/2018/09/ENOC-2018-report-on-Child-Mental-Health-ENG-FV.pdf.
- Deighton J, Croudace T, Fonagy P, Brown J, Patalay P, Wolpert M. Measuring mental health and wellbeing outcomes for children and adolescents to inform practice and policy: a review of child self-report measures. Child Adolesc Psychiatry Mental Health. 2014;8(1):14.
- Zahn-Waxler C, Klimes-Dougan B, Slattery MJ. Internalizing problems of childhood and adolescence: prospects, pitfalls, and progress in understanding the development of anxiety and depression. Dev Psychopathol. 2000;12(3):443–66.
- Rescorla L, Achenbach TM, Ivanova MY, Dumenci L, Almqvist F, Bilenberg N, et al. Epidemiological comparisons of problems and positive qualities reported by adolescents in 24 countries. J Consult Clin Psychol. 2007;75(2):351–8.
- Shahini M, Rescorla L, Wancata J, Ahmeti A. Mental health problems in Kosovar adolescents: results from a national mental health survey. Neuropsychiatr. 2015;29(3):125–32.
- Sandoval M, Lemos S, Vallejo G. Self-reported competences and problems in spanish adolescents: a normative study of the YSR. Psicothema. 2006;18(4):804–9.
- Heyerdahl S, Kvernmo S, Wichstrøm L. Self-reported behavioural/emotional problems in norwegian adolescents from multiethnic areas. Eur Child Adolesc Psychiatry. 2004;13(2):64–72.
- 12. Fitzpatrick C, Deehan A. Competencies and problems of irish children and adolescents. Eur Child Adolesc Psychiatry. 1999;8(1):17–23.
- Helstelä L, Sourander A. Self-reported competence and emotional and behavioral problems in a sample of finnish adolescents. Nord J Psychiatry. 2001;55(6):381–5.
- Roussos A, Francis K, Zoubou V, Kiprianos S, Prokopiou A, Richardson C. The standardization of Achenbach's Youth Self-Report in Greece in a national sample of high school students. Eur Child Adolesc Psychiatry. 2001;10(1):47–53.
- Lewinsohn PM, Hops H, Roberts RE, Seeley JR, Andrews JA. Adolescent psychopathology: I. Prevalence and incidence of depression and other DSM-III-R disorders in high school students. J Abnorm Psychol. 1993;102(1):133–44.
- Alves DE, Roysamb E, Oppedal B, Zachrisson HD. Emotional problems in preadolescents in Norway: the role of gender, ethnic minority status, and home- and school-related hassles. Child Adolesc Psychiatry Ment Health. 2011;5(1):37.
- 17. Nolen-Hoeksema S, Girgus JS. The emergence of gender differences in depression during adolescence. Psychol Bull. 1994;115(3):424–43.
- The State Commission on Migration Issues. Migration Profile of Georgia 2021 [cited 2022 12. Dec]. Available from: https://migration.commission.ge/files/mmp21_eng_web3c.pdf.
- SCMI. Migration Profile of Georgia: The State Commission on Migration Issues.
 ; 2015 [cited 2022 29 Nov]. Available from: https://migration.commission.ge/files/migration_profile_2015_30.11.15.pdf.
- GEOSTAT. Population and Demography: National Statistics Office of Georgia.
 ; 2022 [cited 2022 05 Dec]. Available from: https://www.geostat.ge/en/modules/categories/316/population-and-demography.
- Antia K, Boucsein J, Deckert A, Dambach P, Račaitė J, Šurkienė G, et al. Effects of International Labour Migration on the Mental Health and well-being of left-behind children: a systematic literature review. Int J Environ Res Public Health. 2020;17(12):4335.
- 22. Cheng J, Sun YH. Depression and anxiety among left-behind children in China: a systematic review. Child Care Health Dev. 2015;41(4):515–23.
- 23. Ding L, Yuen LW, Buhs ES, Newman IM. Depression among Chinese Left-Behind children: a systematic review and meta-analysis. Child Care Health Dev. 2019;45(2):189–97.
- Fellmeth G, Rose-Clarke K, Zhao C, Busert LK, Zheng Y, Massazza A, et al. Health impacts of parental migration on left-behind children and adolescents: a systematic review and meta-analysis. The Lancet. 2018;392(10164):2567–82.
- 25. GEOSTAT. Morbidity with acute and chronic diseases by main disease group: National Statistics office of Georgia; 2022 [cited 2022 05 Dec]. Available from: https://www.geostat.ge/en/modules/categories/54/healthcare.

- 26. Administration of the State Representative. Regions: Classification of urbanrural population of Samegrelo-Zemo Svaneti region Online: Georgia; 2023 [cited 2023 19 Jan]. Available from: https://szs.gov.ge/geo/static/39.
- Achenbach TM. Integrative guide for the 1991 CBCL/4–18. YSR, and TRF profiles: Department of Psychiatry, University of Vermont; 1991.
- Achenback T, Rescorla LA. Manual for the ASEBA school-age forms & profiles. Burling: University of Vermont (Research center for children, youth and families); 2001.
- Ivanova MY, Achenbach TM, Rescorla LA, Dumenci L, Almqvist F, Bilenberg N, et al. The generalizability of the Youth Self-Report syndrome structure in 23 societies. J Consult Clin Psychol. 2007;75(5):729–38.
- Cavallo F, Zambon A, Borraccino A, Raven-Sieberer U, Torsheim T, Lemma P, et al. Girls growing through adolescence have a higher risk of poor health. Qual Life Res. 2006;15(10):1577–85.
- Kaye-Tzadok A, Kim SS, Main G. Children's subjective well-being in relation to gender — what can we learn from dissatisfied children? Child Youth Serv Rev. 2017;80:96–104.
- West P, Sweeting H. Fifteen, female and stressed: changing patterns of psychological distress over time. J Child Psychol Psychiatry. 2003;44(3):399–411.
- Duinhof EL, Stevens GW, van Dorsselaer S, Monshouwer K, Vollebergh WA. Ten-year trends in adolescents' self-reported emotional and behavioral problems in the Netherlands. Eur Child Adolesc Psychiatry. 2015;24(9):1119–28.
- Wiklund M, Malmgren-Olsson EB, Ohman A, Bergström E, Fjellman-Wiklund A. Subjective health complaints in older adolescents are related to perceived stress, anxiety and gender - a cross-sectional school study in Northern Sweden. BMC Public Health. 2012;12:993.
- Schraml K, Perski A, Grossi G, Simonsson-Sarnecki M. Stress symptoms among adolescents: the role of subjective psychosocial conditions, lifestyle, and selfesteem. J Adolesc. 2011;34(5):987–96.
- Leadbeater BJ, Kuperminc GP, Blatt SJ, Hertzog C. A multivariate model of gender differences in adolescents' internalizing and externalizing problems. Dev Psychol. 1999;35(5):1268–82.
- van Vuuren CL, Uitenbroek DG, van der Wal MF, Chinapaw MJM. Sociodemographic differences in 10-year time trends of emotional and behavioural problems among adolescents attending secondary schools in Amsterdam, the Netherlands. Eur Child Adolesc Psychiatry. 2018;27(12):1621–31.
- Hyde JS, Mezulis AH, Abramson LY. The ABCs of depression: integrating
 affective, biological, and cognitive models to explain the emergence of the
 gender difference in depression. Psychol Rev. 2008;115(2):291–313.
- Lien L, Dalgard F, Heyerdahl S, Thoresen M, Bjertness E. The relationship between age of menarche and mental distress in norwegian adolescent girls and girls from different immigrant groups in Norway: results from an urban city cross-sectional survey. Soc Sci Med. 2006;63(2):285–95.
- Oldehinkel AJ, Bouma EM. Sensitivity to the depressogenic effect of stress and HPA-axis reactivity in adolescence: a review of gender differences. Neurosci Biobehav Rev. 2011;35(8):1757–70.
- Hagquist C. Discrepant trends in mental health complaints among younger and older adolescents in Sweden: an analysis of WHO data 1985–2005. J Adolesc Health. 2010;46(3):258–64.
- 42. Kapi A, Veltsista A, Kavadias G, Lekea V, Bakoula C. Social determinants of self-reported emotional and behavioral problems in greek adolescents. Soc Psychiatry Psychiatr Epidemiol. 2007;42(7):594–8.
- 43. Lesinskiene S, Girdzijauskiene S, Gintiliene G, Butkiene D, Puras D, Goodman R, et al. Epidemiological study of child and adolescent psychiatric disorders in Lithuania. BMC Public Health. 2018;18(1):548.
- 44. Griesbach D, Amos A, Currie C. Adolescent smoking and family structure in Europe. Soc Sci Med. 2003;56(1):41–52.
- Bagwell CL, Coie JD. The best friendships of aggressive boys: relationship quality, conflict management, and rule-breaking behavior. J Exp Child Psychol. 2004;88(1):5–24.
- Hjern A, Alfven G, Ostberg V. School stressors, psychological complaints and psychosomatic pain. Acta Paediatr. 2008;97(1):112–7.
- Salmela-Aro K, Tynkkynen L. Gendered pathways in school burnout among adolescents. J Adolesc. 2012;35(4):929–39.
- Bradshaw J, Martorano B, Natali L, de Neubourg C. Children's Subjective Well-Being in Rich Countries. Child Indic Res. 2013;6(4):619–35.
- Wiklund M, Bengs C, Malmgren-Olsson EB, Ohman A. Young women facing multiple and intersecting stressors of modernity, gender orders and youth. Soc Sci Med. 2010;71(9):1567–75.
- Broberg AG, Ekeroth K, Gustafsson PA, Hansson K, Hägglöf B, Ivarsson T, et al. Self-reported competencies and problems among swedish adolescents: a

- normative study of the YSR. Youth Self Report. Eur Child Adolesc Psychiatry. 2001:10(3):186–93
- Torsheim T, Ravens-Sieberer U, Hetland J, Välimaa R, Danielson M, Overpeck M. Cross-national variation of gender differences in adolescent subjective health in Europe and North America. Soc Sci Med. 2006;62(4):815–27.
- Antia K, Rodoreda AB, Winkler V. Parental migration and left-behind children in Georgia – school teachers' experience and perception: a qualitative study. BMC Public Health. 2022;22(1):2077.
- Gassmann F, Siegel M, Vanore M, Waidler J. Unpacking the relationship between parental Migration and Child well-Being: evidence from Moldova and Georgia. Child Indic Res. 2018;11(2):423–40.
- Cebotari V, Siegel M, Mazzucato V. Migration and child health in Moldova and Georgia. Comp Migration Stud. 2018;6(1):1–22.
- 55. Vanore M. All in the family: family-member migration and the psychosocial health of children in Georgia. Migration Stud. 2016;4(3):309–30.
- Botezat A, Pfeiffer F. The impact of parental labour migration on left-behind children's educational and psychosocial outcomes: evidence from Romania. Popul Space Place. 2020;26(2):e2277.
- 57. Tomşa R, Jenaro C. Children left behind in romania: anxiety and predictor variables. Psychol Rep. 2015;116(2):485–512.
- Leskauskas D, Adomaitienė V, Šeškevičienė G, Čėsnaitė E, Šmigelskas K.
 Self-reported emotional and behavioral problems of left-behind children in Lithuania. Child Indic Res. 2020;13(4):1203–16.
- Zhao C, Wang F, Zhou X, Jiang M, Hesketh T. Impact of parental migration on psychosocial well-being of children left behind: a qualitative study in rural China. Int J Equity Health. 2018;17(1):80.
- Aryal N, Regmi PR, van Teijlingen E, Simkhada P, Mahat P. Adolescents left behind by migrant workers: a call for community-based mental health interventions in Nepal. WHO South East Asia J Public Health. 2019;8(1):38–41.
- Hu H, Gao J, Jiang H, Jiang H, Guo S, Chen K et al. A Comparative Study of Behavior Problems among Left-Behind Children, Migrant Children and Local Children.Int J Environ Res Public Health. 2018;15(4).
- 62. Umami R, Turnip SS. Emotional and behavioral problems among left-behind children in Indonesia. Indian J Psychol Med. 2019;41(3):240–5.
- 63. Lee SJ, Park SH, Cloninger CR, Chae H. Behavior problems and personality in korean high school students. PeerJ. 2018;6:e6106.

Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

- **K.A.** is a PhD candidate at the Heidelberg Institute of Global Health (HIGH) and holds a Medical Doctor's degree and a master's degree in Health Management. She is a Georgian citizen and collaborates with the School of Health Sciences, the University of Georgia. Her research focuses on migration and health, child and adolescent mental health and well-being.
- **J.R.** is a PhD candidate and junior assistant at the Institute of Health Sciences, Faculty of Medicine, Vilnius University. She holds a master's degree in Public Health. She is a lecturer for child and adolescent health safety, and her research focuses on child health, particularly on left-behind children of migrant parents.
- **G. Š** is a professor at the Institute of Health Sciences, Faculty of Medicine, Vilnius University. She has more than 45 years of experience in child health research, is the author of several textbooks for child health education and safety and is an editorial board member of several scientific journals. She is a former Director of the Public Health Institute at Vilnius University and was recognized for honoured healthcare worker status by the Ministry of Heath of The Republic of Lithuania.
- **V.W.** is the research group leader for "Epidemiology of Transition" at the Heidelberg Institute of Global Health (HIGH). Considerations of health and its determinants in population groups facing major changes are his main interests. He focuses on migrants, non-communicable diseases and the analysis of longitudinal data, as well as systematic reviews.