

Universality Theorems for Some Composite Functions

Kęstutis Janulis^a, Donatas Jurgaitis^b,
Antanas Laurinćikas^a and Renata Macaitienė^c

^a*Faculty of Mathematics and Informatics, Vilnius University*
Naugarduko str. 24, LT-03225 Vilnius, Lithuania

^b*Institute of Informatics, Mathematics and E-Studies, Šiauliai University*
P. Višinskio str. 19, LT-77156 Šiauliai, Lithuania

^c*Research Institute, Šiauliai University*
Vilniaus str. 88, LT-76285 Šiauliai, Lithuania
E-mail: kestutis.janulis@gmail.com
E-mail: d.jurgaitis@cr.su.lt
E-mail: antanas.laurincikas@mif.vu.lt
E-mail(*corresp.*): renata.macaitiene@mi.su.lt

Received June 15, 2015; revised November 17, 2015; published online January 15, 2016

Abstract. In [5], it was proved that a collection consisting from Dirichlet L -functions and periodic Hurwitz zeta-functions is universal in the sense that the shifts of those functions approximate simultaneously a given collection of analytic functions. In the paper, we prove theorems on the universality of composite functions of the above collection.

Keywords: Dirichlet L -function, Hurwitz zeta-function, mixed joint universality, periodic Hurwitz zeta-function, universality.

AMS Subject Classification: 11M06; 11M41.

1 Introduction

In [16], Voronin discovered the universality property of the Riemann zeta-function $\zeta(s)$, $s = \sigma + it$, on the approximation of analytic functions from a wide class by shifts $\zeta(s + i\tau)$, $\tau \in \mathbb{R}$. At the moment, it is known that the majority of zeta and L -functions are universal in the above sense. Also, some zeta and L -functions are jointly universal: their shifts approximate simultaneously a given collection of analytic functions. A series of works are devoted to mixed joint universality when a collection of analytic functions are approximated simultaneously by shifts of zeta-functions with Euler product and without Euler product. The first result in this direction belongs to H. Mishou who proved [14] the joint universality of the function $\zeta(s)$ and the Hurwitz zeta-function $\zeta(s, \alpha)$

with transcendental parameter α . This result has been generalized in [6] for a periodic zeta and a periodic Hurwitz zeta-functions. In [7], the mixed joint universality has been obtained for a wide collection consisting from periodic zeta and periodic Hurwitz zeta-functions. We remind that the periodic Hurwitz zeta-function $\zeta(s, \alpha; \mathbf{a})$, where α , $0 < \alpha \leq 1$, is a fixed parameter and $\mathbf{a} = \{a_m : m \in \mathbb{N}_0 = \mathbb{N} \cup \{0\}\}$ is a periodic sequence of complex numbers, is a generalization of the classical Hurwitz zeta-function $\zeta(s, \alpha)$ when $a_m \equiv 1$, and is defined, for $\sigma > 1$, by the series

$$\zeta(s, \alpha; \mathbf{a}) = \sum_{m=0}^{\infty} \frac{a_m}{(m + \alpha)^s},$$

and by analytic continuation elsewhere. In [2], the mixed joint universality for a system of functions

$$\zeta(s), \zeta(s, \alpha_1, \mathbf{a}_{1l_1}), \dots, \zeta(s, \alpha_1, \mathbf{a}_{1l_1}), \dots, \zeta(s, \alpha_r, \mathbf{a}_{rl_1}), \dots, \zeta(s, \alpha_r, \mathbf{a}_{rl_r})$$

has been considered. In a series of papers [11, 12, 15], the Riemann zeta-function has been replaced by zeta-functions of certain cusp forms. In [5], in place of the function $\zeta(s)$ a collection of Dirichlet L -functions $L(s, \chi)$ has been put. We will state the latter result.

Let $D = \{s \in \mathbb{C} : \frac{1}{2} < \sigma < 1\}$. Denote by \mathcal{K} the class of compact subsets of the strip D with connected complements, and by $H_0(K)$ and $H(K)$, $K \in \mathcal{K}$, the classes of continuous non-vanishing and continuous on K functions, respectively, which are analytic in the interior of K . Let $\text{meas}A$ be the Lebesgue measure of a measurable set $A \subset \mathbb{R}$. Suppose that $\mathbf{a}_{jl} = \{a_{mj} : m \in \mathbb{N}_0\}$ is a periodic sequence of complex numbers with minimal period $k_{jl} \in \mathbb{N}$, $j = 1, \dots, r$, $l = 1, \dots, l_j$. Let k_j be the least common multiple of the periods k_{j1}, \dots, k_{jl_j} , and

$$A_j = \begin{pmatrix} a_{1j1} & a_{1j2} & \dots & a_{1jl_j} \\ a_{2j1} & a_{2j2} & \dots & a_{2jl_j} \\ \dots & \dots & \dots & \dots \\ a_{k_j j1} & a_{k_j j2} & \dots & a_{k_j jl_j} \end{pmatrix}, \quad j = 1, \dots, r.$$

Then, in [5], the following theorem has been proved.

Theorem 1. *Suppose that χ_1, \dots, χ_d are pairwise non-equivalent Dirichlet characters, the numbers $\alpha_1, \dots, \alpha_r$ are algebraically independent over the field of rational numbers \mathbb{Q} , and that $\text{rank}(A_j) = l_j$, $j = 1, \dots, r$. For $j = 1, \dots, d$, let $K_j \in \mathcal{K}$ and $f_j \in H_0(K_j)$, and, for $j = 1, \dots, r$, $l = 1, \dots, l_j$, let $K_{jl} \in \mathcal{K}$ and $f_{jl} \in H(K_{jl})$. Then, for every $\varepsilon > 0$,*

$$\liminf_{T \rightarrow \infty} \frac{1}{T} \text{meas} \left\{ \tau \in [0, T] : \sup_{1 \leq j \leq d} \sup_{s \in K_j} |L(s + i\tau, \chi_j) - f_j(s)| < \varepsilon, \right. \\ \left. \sup_{1 \leq j \leq r} \sup_{1 \leq l \leq l_j} \sup_{s \in K_{jl}} |\zeta(s + i\tau, \alpha_j; \mathbf{a}_{jl}) - f_{jl}(s)| < \varepsilon \right\} > 0.$$

Denote by $H(D)$ the space of analytic on D functions equipped with the topology of uniform convergence on compacta. In [8] and [10], the Voronin theorem has been generalized for $F(\zeta(s))$ with certain operators $F : H(D) \rightarrow H(D)$, in [9], the universality of $F(\zeta(s), \zeta(s, \alpha))$ has been studied with operators $F : H^2(D) \rightarrow H(D)$. The papers [3] and [4] are devoted to the universality of the functions $F(L(s, \chi_1), \dots, L(s, \chi_{r_1}), \zeta(s, \alpha_1), \dots, \zeta(s, \alpha_{r_2}))$ for some operators $F : H^{r_1+r_2}(D) \rightarrow H(D)$. The aim of the present paper is the universality of composite functions of a collection of L and zeta-functions in Theorem 1, i.e., we consider the universality of the functions

$$F(L(s, \chi_1), \dots, L(s, \chi_d), \zeta(s, \alpha_1; \mathbf{a}_{11}), \dots, \zeta(s, \alpha_1; \mathbf{a}_{1l_1}), \dots, \zeta(s, \alpha_r; \mathbf{a}_{r1}), \dots, \zeta(s, \alpha_r; \mathbf{a}_{rl_r}))$$

for some operators F .

First we deal with approximation of functions from the class $H(K)$, $K \in \mathcal{K}$. Let, for brevity, $v = d + l_1 + \dots + l_r$. We say that the operator $F : H^v(D) \rightarrow H(D)$ belongs to the class $Lip(\beta_1, \dots, \beta_v)$, $\beta_1 > 0, \dots, \beta_v > 0$, if the following hypotheses are satisfied:

1° For every polynomial $p = p(s)$ and all sets $K_1, \dots, K_d \in \mathcal{K}$, there exists an element $g = (g_1, \dots, g_d, g_{11}, \dots, g_{1l_1}, \dots, g_{r1}, \dots, g_{rl_r}) \in F^{-1}\{p\} \subset H^v(D)$ such that $g_j \neq 0$ on K_j , $j = 1, \dots, d$;

2° For all $K \in \mathcal{K}$, there exist a constant $c > 0$ and sets $K_1, \dots, K_v \in \mathcal{K}$ such that, for all $(g_{j1}, \dots, g_{jv}) \in H^v(D)$, $j = 1, 2$,

$$\begin{aligned} \sup_{s \in K} |F(g_{11}(s), \dots, g_{1v}(s)) - F(g_{21}(s), \dots, g_{2v}(s))| \\ \leq c \sup_{1 \leq j \leq v} \sup_{s \in K_j} |g_{1j}(s) - g_{2j}(s)|^{\beta_j}. \end{aligned}$$

Theorem 2. *Suppose that χ_1, \dots, χ_d are pairwise non-equivalent Dirichlet characters, the numbers $\alpha_1, \dots, \alpha_r$ are algebraically independent over the field of rational numbers \mathbb{Q} , $\text{rank}(A_j) = l_j$, $j = 1, \dots, r$, and that $F \in Lip(\beta_1, \dots, \beta_v)$. Let $K \in \mathcal{K}$ and $f(s) \in H(K)$. Then, for every $\varepsilon > 0$,*

$$\begin{aligned} \liminf_{T \rightarrow \infty} \frac{1}{T} \text{meas} \left\{ \tau \in [0, T] : \sup_{s \in K} |F(L(s + i\tau, \chi_1), \dots, L(s + i\tau, \chi_d), \zeta(s + i\tau, \alpha_1; \mathbf{a}_{11}), \dots, \zeta(s + i\tau, \alpha_1; \mathbf{a}_{1l_1}), \dots, \zeta(s + i\tau, \alpha_r; \mathbf{a}_{r1}), \dots, \zeta(s + i\tau, \alpha_r; \mathbf{a}_{rl_r})) - f(s)| < \varepsilon \right\} > 0. \end{aligned}$$

We give an example of the operator $F \in Lip(\beta_1, \dots, \beta_v)$. Let, for $(g_1, \dots, g_d, g_{11}, \dots, g_{1l_1}, \dots, g_{r1}, \dots, g_{rl_r}) \in H^v(D)$,

$$\begin{aligned} F(g_1, \dots, g_d, g_{11}, \dots, g_{1l_1}, \dots, g_{r1}, \dots, g_{rl_r}) = c_1 g_1^{(n_1)} + \dots + c_d g_d^{(n_d)} \\ + c_{11} g_{11}^{(n_{11})} + \dots + c_{1l_1} g_{1l_1}^{(n_{1l_1})} + \dots + c_{r1} g_{r1}^{(n_{r1})} + \dots + c_{rl_r} g_{rl_r}^{(n_{rl_r})}, \end{aligned}$$

where $c_1, \dots, c_d, c_{11}, \dots, c_{1l_1}, \dots, c_{r1}, \dots, c_{rl_r} \in \mathbb{C} \setminus \{0\}$, $n_1, \dots, n_d, n_{11}, \dots, n_{1l_1}, \dots, n_{r1}, \dots, n_{rl_r} \in \mathbb{N}$, and $f^{(n)}$ denotes the n th derivative of f . It is not

difficult to see that, for each polynomial $p = p(s)$ and all sets $K_1, \dots, K_d \in \mathcal{K}$, there exists an element $\underline{g} \in F^{-1}\{p\}$ such that $g_j(s) \neq 0$ on K_j , $j = 1, \dots, d$. For example, if

$$p(s) = a_k s^k + a_{k-1} s^{k-1} + \dots + a_0, \quad a_k \neq 0,$$

we can take $\underline{g} = (1, \dots, 1, 1, \dots, 1, 1, \dots, g_{rl_r})$, where

$$g_{rl_r}(s) = \frac{1}{c_{rl_r}} \left(\frac{a_n s^{k+n_{rl_r}}}{(k+1) \cdots (k+n_{rl_r})} + \dots + \frac{a_0 s^{n_{rl_r}}}{1 \cdots n_{rl_r}} \right).$$

Thus, hypothesis 1° of the class $Lip(\beta_1, \dots, \beta_v)$ is satisfied.

Hypothesis 2° of the class $Lip(\beta_1, \dots, \beta_v)$ follows from the Cauchy integral formula. We write F in a more convenient form

$$F(g_1, \dots, g_v) = \sum_{j=1}^v c_j g_j^{(n_j)}.$$

Let $K \in \mathcal{K}$, and $K \subset G \subset K_1$, where G is an open set and $K_1 \in \mathcal{K}$. Moreover, let L be a simple closed contour lying in $K_1 \setminus G$ and containing inside the K . Then the Cauchy integral formula shows that, for $(g_{j1}, \dots, g_{jv}) \in H^v(D)$, $j = 1, 2$, and $s \in K$,

$$\begin{aligned} |F(g_{11}(s), \dots, g_{1v}(s)) - F(g_{21}(s), \dots, g_{2v}(s))| &= \left| \sum_{j=1}^v c_j \frac{n_j!}{2\pi i} \int_L \frac{g_{1j}(z) - g_{2j}(z)}{(z-s)^{n_j+1}} dz \right| \\ &\leq \sum_{j=1}^v |c_j| C_j \sup_{s \in L} |g_{1j}(s) - g_{2j}(s)| \leq c \sup_{1 \leq j \leq v} \sup_{s \in K_1} |g_{1j}(s) - g_{2j}(s)| \end{aligned}$$

with some constants $C_j > 0$, $j = 1, \dots, v$, and $c > 0$. Thus we have that $F \in Lip(1, \dots, 1)$, and in this case, $K_1 = \dots = K_v = K_1$.

Now we give some other classes of operators F . Let

$$S = \{g \in H(D) : g(s) \neq 0 \text{ or } g(s) \equiv 0\}.$$

Moreover, $v_1 = \sum_{j=1}^r l_j$.

Theorem 3. *Suppose that the characters χ_1, \dots, χ_d , the numbers $\alpha_1, \dots, \alpha_r$ and the sequences \mathbf{a}_{jl} , $j = 1, \dots, r$, $l = 1, \dots, l_j$, satisfy the hypotheses of Theorem 2, and that $F : H^v(D) \rightarrow H(D)$ be a continuous operator such that, for every open set $G \subset H(D)$, the set $(F^{-1}G) \cap (S^d \times H^{v_1}(D))$ is not empty. Let $K \in \mathcal{K}$ and $f(s) \in H(K)$. Then the assertion of Theorem 2 is true.*

We note that the hypothesis $(F^{-1}G) \cap (S^d \times H^{v_1}(D)) \neq \emptyset$ for every open set $G \subset H(D)$ is general but sufficiently complicated. Obviously, it is satisfied if every $g \in H(D)$ has a preimage in the set $S^d \times H^{v_1}(D)$. On the other hand, Theorem 3 implies the following modification of Theorem 2.

Theorem 4. *Suppose that the characters χ_1, \dots, χ_d , the numbers $\alpha_1, \dots, \alpha_r$ and the sequences \mathbf{a}_{jl} , $j = 1, \dots, r$, $l = 1, \dots, l_j$, satisfy the hypotheses of Theorem 2, and that $F : H^v(D) \rightarrow H(D)$ is a continuous operator such that, for every polynomial $p = p(s)$, the set $(F^{-1}\{p\}) \cap (S^d \times H^{v_1}(D))$ is not empty. Let $K \in \mathcal{K}$ and $f(s) \in H(K)$. Then the assertion of Theorem 2 is true.*

Clearly, hypothesis 2° of the class $Lip(\beta_1, \dots, \beta_v)$ implies the continuity of F . However, hypothesis 1° is weaker than the requirement $(F^{-1}\{p\}) \cap (S^d \times H^{v_1}(D)) \neq \emptyset$.

Non-vanishing of the polynomial $p(s)$ in a bounded region can be controlled by its constant term. Therefore, sometimes it is more convenient to consider operators F on the space $H^v(D_V, D) = H^d(D_V) \times H^{v_1}(D)$, where, for $V > 0$, $D_V = \{s \in \mathbb{C} : \frac{1}{2} < \sigma < 1, |t| < V\}$. Analogically, let

$$S_V = \{g \in H(D_V) : g(s) \neq 0 \text{ or } g(s) \equiv 0\}.$$

Then we have the following result.

Theorem 5. *Suppose that the characters χ_1, \dots, χ_d , the numbers $\alpha_1, \dots, \alpha_r$ and the sequences $\mathbf{a}_{jl}, j = 1, \dots, r, l = 1, \dots, l_j$, satisfy the hypotheses of Theorem 2, $K \in \mathcal{K}$, $f(s) \in H(K)$ and $V > 0$ is such that $K \subset D_V$. Let $F : H^v(D_V, D) \rightarrow H(D_V)$ be a continuous operator such that, for every polynomial $p = p(s)$, the set $(F^{-1}\{p\}) \cap (S_V^d \times H^{v_1}(D))$ is not empty. Then the assertion of Theorem 2 is true.*

For example, Theorem 5 can be applied for the operator

$$F(g_1, \dots, g_v) = c_1 g_1^{(n_1)} + \dots + c_d g_d^{(n_d)}, \quad n_1, \dots, n_d \in \mathbb{N}.$$

Now we consider approximation of analytic functions from the image of the set $S^d \times H^{v_1}(S)$ of the operator $F : H^v(D) \rightarrow H(D)$.

Theorem 6. *Suppose that the characters χ_1, \dots, χ_d , the numbers $\alpha_1, \dots, \alpha_r$ and the sequences $\mathbf{a}_{jl}, j = 1, \dots, r, l = 1, \dots, l_j$, satisfy the hypotheses of Theorem 2, and that $F : H^v(D) \rightarrow H(D)$ is a continuous operator. Let $K \subset D$ be a compact subset, and $f(s) \in F(S^d \times H^{v_1}(D))$. Then the assertion of Theorem 2 is true.*

It is not easy to describe the set $F(S^d \times H^{v_1}(D))$. The next theorem is an example with sufficiently simple set contained in $F(S^d \times H^{v_1}(D))$.

Suppose that $a_1, \dots, a_k \in \mathbb{C}$ are pairwise different numbers, and

$$H_k(D) = \{g \in H(D) : (g(s) - a_j)^{-1} \in H(D), j = 1, \dots, k\}.$$

Theorem 7. *Suppose that the characters χ_1, \dots, χ_d , the numbers $\alpha_1, \dots, \alpha_r$ and the sequences $\mathbf{a}_{jl}, j = 1, \dots, r, l = 1, \dots, l_j$, satisfy the hypotheses of Theorem 2, and that $F : H^v(D) \rightarrow H(D)$ is a continuous operator such that $F(S^d \times H^{v_1}(D)) \supset H_k(D)$. For $k = 1$, let $K \in \mathcal{K}$, $f(s) \in H(K)$ and $f(s) \neq a_1$ on K . For $k \geq 2$, let $K \subset D$ be an arbitrary compact subset, and $f(s) \in H_k(D)$. Then the assertion of Theorem 2 is true.*

For example, let $k = 2$ and $a_1 = 1, a_2 = -1$. Then Theorem 7 implies the universality of the function

$$\begin{aligned} \sin(L(s, \chi_1) + \dots + L(s, \chi_d) + \zeta(s, \alpha_1; \mathbf{a}_{11}) + \dots + \zeta(s, \alpha_1; \mathbf{a}_{1l_1}) + \dots \\ + \zeta(s, \alpha_r; \mathbf{a}_{r1}) + \dots + \zeta(s, \alpha_r; \mathbf{a}_{rl_r})). \end{aligned}$$

For this, it suffices to consider the equation

$$\frac{e^{i\Sigma(s)} - e^{-i\Sigma(s)}}{2i} = f, \quad f \in H(D), \quad a_1 = 1, \quad a_2 = -1,$$

where $\Sigma(s)$ is the sum under the sign of sin.

2 Proof of Theorem 2

Theorem 2 is a result of Theorem 1, properties of the class $Lip(\beta_1, \dots, \beta_v)$ and of the Mergelyan theorem on the approximation of analytic functions by polynomials. We state this theorem in the form of the next lemma.

Lemma 1. *Suppose that $K \subset \mathbb{C}$ is a compact subset with connected complement, and $f(s)$ is a continuous function on K which is analytic in the interior of K . Then, for every $\varepsilon > 0$, there exists a polynomial $p(s)$ such that*

$$\sup_{s \in K} |f(s) - p(s)| < \varepsilon.$$

Proof of the lemma can be found in [13] and [17].

Proof of Theorem 2. Lemma 1 implies the existence of the polynomial $p = p(s)$ such that

$$\sup_{s \in K} |f(s) - p(s)| < \frac{\varepsilon}{2}. \quad (2.1)$$

Using hypothesis 1° of the class $Lip(\beta_1, \dots, \beta_v)$, we have that, for all sets $K_1, \dots, K_d \in \mathcal{K}$, there exists an element $(g_1, \dots, g_d, g_{11}, \dots, g_{1l_1}, \dots, g_{r1}, \dots, g_{rl_r}) \in F^{-1}\{p\}$ such that $g_j(s) \neq 0$ on K_j , $j = 1, \dots, d$. Suppose that $\tau \in \mathbb{R}$ satisfies the inequalities

$$\sup_{1 \leq j \leq d} \sup_{s \in K_j} |L(s + i\tau, \chi_j) - f_j(s)| < c^{-\frac{1}{\beta}} \left(\frac{\varepsilon}{4}\right)^{\frac{1}{\beta}}, \quad (2.2)$$

$$\sup_{1 \leq j \leq r} \sup_{1 \leq l \leq l_j} \sup_{s \in K_{jl}} |\zeta(s + i\tau, \alpha_j; \mathbf{a}_{jl}) - f_{jl}(s)| < c^{-\frac{1}{\beta}} \left(\frac{\varepsilon}{4}\right)^{\frac{1}{\beta}}, \quad (2.3)$$

where the sets $K_1, \dots, K_d, K_{11}, \dots, K_{1l_1}, \dots, K_{r1}, \dots, K_{rl_r} \in \mathcal{K}$ correspond the set K in hypothesis 2° of the class $Lip(\beta_1, \dots, \beta_v)$, and $\beta = \min_{1 \leq j \leq v} \beta_j$, with notation $K_{1l} = K_{d+l}$, $j = 1, \dots, l_1, \dots, K_{rl} = K_{d+l_1+\dots+l_{r-1}+l}$, $l = 1, \dots, l_r$. Then, in view of Theorem 1, the set of τ satisfying inequalities (2.2) and (2.3) has a positive lower density. Moreover, hypothesis 2° of the class $Lip(\beta_1, \dots, \beta_v)$ shows that, for such τ ,

$$\begin{aligned} \sup_{s \in K} |F(\underline{L}(s + i\tau, \underline{\chi}, \underline{\alpha}, \underline{\mathbf{a}})) - p(s)| &\leq \sup_{1 \leq j \leq d} \sup_{s \in K_j} |L(s + i\tau, \chi_j) - f_j(s)|^{\beta_j} \\ &+ c \sup_{1 \leq j \leq r} \sup_{1 \leq l \leq l_j} \sup_{s \in K_{jl}} |\zeta(s + i\tau, \alpha_j; \mathbf{a}_{jl}) - f_{jl}(s)|^{\beta_{jl}} \leq 2cc^{-\frac{\beta}{\beta}} \left(\frac{\varepsilon}{4}\right)^{\frac{\beta}{\beta}} = \frac{\varepsilon}{2}. \end{aligned} \quad (2.4)$$

Here $\underline{\chi} = (\chi_1, \dots, \chi_d)$, $\underline{\alpha} = (\alpha_1, \dots, \alpha_r)$, $\underline{\mathbf{a}} = (\mathbf{a}_{11}, \dots, \mathbf{a}_{1l_1}, \dots, \mathbf{a}_{r1}, \dots, \mathbf{a}_{rl_r})$ and

$$\underline{L}(s + i\tau, \underline{\chi}, \underline{\alpha}, \underline{\mathbf{a}}) = (L(s, \chi_1), \dots, L(s, \chi_d), \zeta(s, \alpha_1; \mathbf{a}_{11}), \dots, \zeta(s, \alpha_1; \mathbf{a}_{1l_1}), \dots, \zeta(s, \alpha_r; \mathbf{a}_{r1}), \dots, \zeta(s, \alpha_r; \mathbf{a}_{rl_r})),$$

and $\beta_{1l} = \beta_{d+l}$, $l = 1, \dots, l_1, \dots$, $\beta_{rl} = \beta_{d+l_1+\dots+l_{r-1}+l}$, $l = 1, \dots, l_r$. Thus, by the above remark,

$$\liminf_{T \rightarrow \infty} \frac{1}{T} \text{meas} \left\{ \tau \in [0, T] : \sup_{s \in K} |F(\underline{L}(s + i\tau, \underline{\chi}, \underline{\alpha}, \underline{\mathbf{a}})) - p(s)| < \frac{\varepsilon}{2} \right\} > 0.$$

Combining this with inequality (2.1) proves the theorem.

3 Elements of probability theory

For the proof of Theorems 3 – 7, we apply a probabilistic approach based on limit theorems for weakly convergent probability measures in the space of analytic functions. We start with a limit theorem for $\underline{L}(s + i\tau, \underline{\chi}, \underline{\alpha}, \underline{\mathbf{a}})$ obtained in [5], Theorem 2.

Denote by $\mathcal{B}(X)$ the Borel σ -field of the space X . Let $\gamma = \{s \in \mathbb{C} : |s| = 1\}$ be the unit circle on the complex plane, and

$$\Omega = \prod_{p \in \mathcal{P}} \gamma_p, \quad \hat{\Omega} = \prod_{m \in \mathbb{N}_0} \gamma_m,$$

where \mathcal{P} is the set of all prime numbers, and $\gamma_p = \gamma$ for all $p \in \mathcal{P}$ and $\gamma_m = \gamma$ for all $m \in \mathbb{N}_0$. In view of the classical Tikhonov theorem, the tori Ω and $\hat{\Omega}$ with the product topology and pointwise multiplication are compact topological Abelian groups. Moreover, let

$$\underline{\Omega} = \Omega \times \hat{\Omega}_1 \times \dots \times \hat{\Omega}_r,$$

where $\hat{\Omega}_j = \hat{\Omega}$ for all $j = 1, \dots, r$. Then again $\underline{\Omega}$ is a compact topological Abelian group. This leads to the probability space $(\underline{\Omega}, \mathcal{B}(\underline{\Omega}), \underline{m}_H)$, where \underline{m}_H is the probability Haar measure on $(\underline{\Omega}, \mathcal{B}(\underline{\Omega}))$. Denote by $\omega(p)$ the projection of the element $\omega \in \Omega$ to the coordinate space γ_p , $p \in \mathcal{P}$, and by $\hat{\omega}_j(m)$ the projection of an element $\hat{\omega}_j \in \hat{\Omega}_j$ to the coordinate space γ_m , $m \in \mathbb{N}_0$, $j = 1, \dots, r$. Let $p^k \parallel m$ mean that $p^k \mid m$ but $p^{k+1} \nmid m$. Extend the function $\omega(p)$ to the set \mathbb{N} by taking

$$\omega(m) = \prod_{p^k \parallel m} \omega^k(p), \quad m \in \mathbb{N}.$$

Denote by $\underline{\omega} = (\omega, \hat{\omega}_1, \dots, \hat{\omega}_r)$ the elements of $\underline{\Omega}$, and, on the probability space $(\underline{\Omega}, \mathcal{B}(\underline{\Omega}), \underline{m}_H)$, define the $H^v(D)$ -valued random element $\underline{L}(s, \underline{\chi}, \underline{\alpha}, \underline{\omega}, \underline{\mathbf{a}})$ by the formula

$$\underline{L}(s, \underline{\chi}, \underline{\alpha}, \underline{\omega}, \underline{\mathbf{a}}) = (L(s, \omega, \chi_1), \dots, L(s, \omega, \chi_d), \zeta(s, \alpha_1, \omega_1; \mathbf{a}_{11}), \dots, \zeta(s, \alpha_1, \omega_1; \mathbf{a}_{1l_1}), \dots, \zeta(s, \alpha_r, \omega_r; \mathbf{a}_{r1}), \dots, \zeta(s, \alpha_r, \omega_r; \mathbf{a}_{rl_r})),$$

where

$$L(s, \omega, \chi_j) = \sum_{m=1}^{\infty} \frac{\chi_j(m)\omega(m)}{m^s}, \quad j = 1, \dots, d,$$

and

$$\zeta(s, \alpha_j, \omega_j; \mathbf{a}_{jl}) = \sum_{m=1}^{\infty} \frac{a_{mjl}\omega_j(m)}{(m + \alpha_j)^s}, \quad j = 1, \dots, r, \quad l = 1, \dots, l_j.$$

We note that the latter series are uniformly convergent on compact subsets of D for almost all $\underline{\omega} \in \underline{\Omega}$. Moreover, for almost $\underline{\omega} \in \underline{\Omega}$, $L(s, \omega, \chi_j)$ can be written in the form

$$L(s, \omega, \chi_j) = \prod_p \left(1 - \frac{\chi_j(p)\omega(p)}{p^s} \right)^{-1}.$$

Denote by $P_{\underline{L}}$ the distribution of the random element $\underline{L}(s, \underline{\chi}, \underline{\alpha}, \underline{\omega}, \underline{\mathbf{a}})$, i.e., the probability measure

$$P_{\underline{L}}(A) = \underline{m}_H(\underline{\omega} \in \underline{\Omega} : \underline{L}(s, \underline{\chi}, \underline{\alpha}, \underline{\omega}, \underline{\mathbf{a}}) \in A), \quad A \in \mathcal{B}(H^v(D)).$$

Then we have the following limit theorem [5].

Lemma 2. *Suppose that the numbers $\alpha_1, \dots, \alpha_r$ are algebraically independent over \mathbb{Q} . Then*

$$P_T(A) \stackrel{\text{def}}{=} \frac{1}{T} \text{meas} \{ \tau \in [0, T] : \underline{L}(s + i\tau, \underline{\chi}, \underline{\alpha}, \underline{\mathbf{a}}) \in A \}, \quad A \in \mathcal{B}(H^v(D))$$

converges weakly to $P_{\underline{L}}$ as $T \rightarrow \infty$.

For the proof a limit theorem for composite function $F(\underline{L}(s, \underline{\chi}, \underline{\alpha}, \underline{\mathbf{a}}))$, we will apply an assertion on the preservation of the weak convergence under mappings. Let X_1 and X_2 be two metric spaces, and let $u : X_1 \rightarrow X_2$ be a $(\mathcal{B}(X_1), \mathcal{B}(X_2))$ -measurable mapping, i.e.,

$$u^{-1}\mathcal{B}(X_2) \subset \mathcal{B}(X_1).$$

Then every probability measure P on $(X_1, \mathcal{B}(X_1))$ induces the unique probability measure Pu^{-1} on $(X_2, \mathcal{B}(X_2))$ defined by

$$Pu^{-1}(A) = P(u^{-1}A), \quad A \in \mathcal{B}(X_2).$$

It is well known that the continuity of u implies its $(\mathcal{B}(X_1), \mathcal{B}(X_2))$ -measurability.

Lemma 3. *Suppose that P_n converges weakly to P as $n \rightarrow \infty$, and that the mapping $u : X_1 \rightarrow X_2$ is continuous. Then $P_n u^{-1}$ converges weakly to Pu^{-1} as $n \rightarrow \infty$.*

Proof of the lemma is given in [1].

Lemma 4. *Suppose that the numbers $\alpha_1, \dots, \alpha_r$ are algebraically independent over \mathbb{Q} and that the operator $F : H^v(D) \rightarrow H(D)$ is continuous. Then*

$$P_{T,F}(A) \stackrel{\text{def}}{=} \frac{1}{T} \text{meas} \{ \tau \in [0, T] : F(\underline{L}(s + i\tau, \underline{\chi}, \underline{\alpha}, \underline{\mathbf{a}})) \in A \}, \quad A \in \mathcal{B}(H^v(D)),$$

converges weakly to $P_{\underline{L}}F^{-1}$ as $T \rightarrow \infty$.

Proof. The lemma is an immediate consequence of Lemmas 2 and 3. \square

For the proof of universality theorems for $F(\underline{L}(s + i\tau, \underline{\chi}, \underline{\alpha}, \underline{\mathbf{a}}))$, we also need the explicit form of the support of the measure $P_{\underline{L}}F^{-1}$. We apply a result of [5] on the support of the measure $P_{\underline{L}}$.

Lemma 5. *Suppose that χ_1, \dots, χ_d are pairwise non-equivalent Dirichlet characters, the numbers $\alpha_1, \dots, \alpha_r$ are algebraically independent over \mathbb{Q} , and that $\text{rank}(A_j) = l_j, j = 1, \dots, r$. Then the support of $P_{\underline{L}}$ is the set $S^d \times H^{v_1}(D)$.*

Lemma 6. *Suppose that the hypotheses of Lemma 5 are satisfied, and that the operator $F : H^v(D) \rightarrow H(D)$ is continuous. Then the support of $P_{\underline{L}}F^{-1}$ is the closure of the set $F(S^d \times H^{v_1}(D))$.*

Proof. Let g be an arbitrary element of the set $F(S^d \times H^{v_1}(D))$, and G be any open neighbourhood of g . Then $F^{-1}G$ is an open neighbourhood of a certain element of the set $S^d \times H^{v_1}(D)$. Therefore, Lemma 5 and properties of a support imply that $P_{\underline{L}}(F^{-1}G) > 0$, hence, $P_{\underline{L}}F^{-1}(G) > 0$. Moreover, in virtue of Lemma 5 again,

$$P_{\underline{L}}F^{-1}(F(S^d \times H^{v_1}(D))) = P_{\underline{L}}(S^d \times H^{v_1}(D)) = 1.$$

Since the support of $P_{\underline{L}}F^{-1}$ is a closed set, from this the lemma follows. \square

We also need one equivalent of the weak convergence of probability measures.

Lemma 7. *Let $P_n, n \in \mathbb{N}$, and P be probability measures on $(X, \mathcal{B}(X))$. Then, P_n , as $n \rightarrow \infty$, converges weakly to P if and only if, for every open set $G \subset X$,*

$$\liminf_{n \rightarrow \infty} P_n(G) \geq P(G).$$

The lemma is a part of Theorem 2.1 from [1].

4 Proofs of other universality theorems

Proof of Theorem 3. It is not difficult to see that, under hypotheses of Theorem 3, the support of the measure $P_{\underline{L}}F^{-1}$ is the whole of $H(D)$. Indeed, if $(F^{-1}G) \cap (S^d \times H^{v_1}(D)) \neq \emptyset$ for every open set $G \subset H(D)$, then we have that, for every element $g \in H(D)$ and its open neighbourhood G , there exists an element of the set $F(S^d \times H^{v_1}(D))$ which belongs to the set G . This shows

that the set $F(S^d \times H^{v_1}(D))$ is dense in $H(D)$. Since, by Lemma 6, the support of $P_{\underline{L}}F^{-1}$ is the closure of $F(S^d \times H^{v_1}(D))$, from this we obtain that the support of $P_{\underline{L}}F^{-1}$ is the whole of $H(D)$.

In view of Lemma 1, there exists a polynomial $p(s)$ satisfying inequality (2.1). Define the set

$$G = \left\{ g \in H(D) : \sup_{s \in K} |g(s) - p(s)| < \frac{\varepsilon}{2} \right\}.$$

Obviously, G is an open neighbourhood of $p(s)$ which is an element of the support of $P_{\underline{L}}F^{-1}$. Therefore, $P_{\underline{L}}F^{-1}(G) > 0$, and Lemmas 4 and 7 imply the inequality

$$\liminf_{T \rightarrow \infty} \frac{1}{T} \text{meas} \left\{ \tau \in [0, T] : F(\underline{L}(s + i\tau, \underline{\chi}, \underline{\alpha}, \underline{\mathbf{a}})) \in G \right\} > 0,$$

or, by definition of G ,

$$\liminf_{T \rightarrow \infty} \frac{1}{T} \text{meas} \left\{ \tau \in [0, T] : \sup_{s \in K} |F(\underline{L}(s + i\tau, \underline{\chi}, \underline{\alpha}, \underline{\mathbf{a}})) - p(s)| < \frac{\varepsilon}{2} \right\} > 0.$$

Combining this with (2.1) gives the assertion of the theorem.

Proof of Theorem 4. Let $\{K_l : l \in \mathbb{N}\} \subset D$ be the sequence of compact subsets such that $K_l \subset K_{l+1}$ for all $l \in \mathbb{N}$,

$$D = \bigcup_{l=1}^{\infty} K_l,$$

and, for every compact subset $K \subset D$, there exists K_l such that $K \subset K_l$. For $g_1, g_2 \in H(D)$, define

$$\rho(g_1, g_2) = \sum_{l=1}^{\infty} 2^{-l} \frac{\sup_{s \in K_l} |g_1(s) - g_2(s)|}{1 + \sup_{s \in K_l} |g_1(s) - g_2(s)|}.$$

Then ρ is a metric on $H(D)$ which induces its topology of uniform convergence on compacta. Moreover, from the definition of ρ we have that the function g_2 approximates g_1 with suitable accuracy if g_2 approximate g_1 uniformly on K_l for sufficiently large l . Obviously, we may choose the sets K_l with connected complements, for example, we can take the closed rectangles. Thus, in $H(D)$, we can limit ourselves by approximation of functions on compact subsets with connected complements.

We will show that the hypotheses of the theorem imply those of Theorem 3. Let $G \subset H(D)$ be an arbitrary non-empty open set. Then, in view of Lemma 1 and the above remark on approximation in $H(D)$, there exists a polynomial $p(s) \in G$. Therefore, the hypothesis $(F^{-1}\{p\}) \cap (S^d \times H^{v_1}(D)) \neq \emptyset$ implies that of Theorem 3 that the set $(F^{-1}G) \cap (S^d \times H^{v_1}(D))$ is non-empty. Thus, Theorem 4 is a corollary of Theorem 3.

Proof of Theorem 5. We apply the arguments used in the proof of Theorem 3 with obvious modifications.

Proof of Theorem 6. Since $f(s) \in F(S^d \times H^{v_1}(D))$, it follows from Lemma 6 that $f(s)$ is an element of the support of the measure $P_{\underline{L}}F^{-1}$. Hence, $P_{\underline{L}}F^{-1}(G) > 0$ for

$$G = \left\{ g \in H(D) : \sup_{s \in K} |g(s) - p(s)| < \varepsilon \right\}.$$

Therefore, the theorem is a consequence of Lemmas 4 and 7.

Proof of Theorem 7. First suppose that $k = 1$. By Lemma 1, there exists a polynomial $p(s)$ such that

$$\sup_{s \in K} |f(s) - p(s)| < \frac{\varepsilon}{4}. \tag{4.1}$$

Since $f(s) \neq a_1$ on K , then also $p(s) \neq a_1$ on K if $\varepsilon > 0$ is rather small. Therefore, on K we can define a continuous branch of the logarithm $\log(p(s) - a_1)$ which will be an analytic function in the interior of K . Again by Lemma 1, we can find a polynomial $q(s)$ such that

$$\sup_{s \in K} \left| p(s) - a_1 - e^{q(s)} \right| < \frac{\varepsilon}{4}. \tag{4.2}$$

Let, for brevity, $g_{a_1}(s) = a_1 + e^{q(s)}$. Then, clearly, $g_{a_1}(s) \in H(D)$, and $g_{a_1}(s) \neq a_1$ on D . Thus, $g_{a_1}(s) \in H_1(D)$. In view of Lemma 6, the support of the measure $P_{\underline{L}}F^{-1}$ contains the closure of the set $H_1(D)$. Therefore, the function $g_{a_1}(s)$ is an element of the support of the measure $P_{\underline{L}}F^{-1}$. Hence, $P_{\underline{L}}F^{-1}(G) > 0$, where

$$G = \left\{ g \in H(D) : \sup_{s \in K} |g(s) - g_{a_1}(s)| < \frac{\varepsilon}{2} \right\}.$$

This together with Lemmas 4 and 7 shows that

$$\liminf_{T \rightarrow \infty} \frac{1}{T} \text{meas} \left\{ \tau \in [0, T] : \sup_{s \in K} |F(\underline{L}(s+i\tau, \underline{\chi}, \underline{\alpha}, \underline{a})) - g_{a_1}(s)| < \frac{\varepsilon}{2} \right\} > 0. \tag{4.3}$$

Inequalities (4.1) and (4.2) imply that

$$\sup_{s \in K} |f(s) - g_{a_1}(s)| < \frac{\varepsilon}{2}.$$

This and (4.3) yield the assertion of the theorem in the case $k = 1$.

The case $k \geq 2$ is contained in Theorem 6.

References

- [1] P. Billingsley. *Convergence of Probability Measures*. Wiley, New York, 1968.
- [2] J. Genys, R. Macaitienė, S. Račkauskienė and D. Šiaučiušas. A mixed joint universality theorem for zeta-functions. *Math. Modelling and Analysis*, **15**(4):431–446, 2010.
- [3] K. Janulis. Remarks on the joint universality of Dirichlet L -functions and Hurwitz zeta-functions. *Šiauliai Math. Semin.*, **9**(17):61–70, 2014.

- [4] K. Janulis and A. Laurinčikas. Joint universality of Dirichlet L -functions and Hurwitz zeta-functions. *Ann. Univ. Sci. Budapest, Sect. Comp.*, **39**:203–214, 2013.
- [5] K. Janulis, A. Laurinčikas, R. Macaitienė and D. Šiaučiūnas. Joint universality of Dirichlet L -functions and periodic Hurwitz zeta-functions. *Math. Modelling and Analysis*, **17**(5):673–685, 2012.
- [6] R. Kačinskaitė and A. Laurinčikas. The joint distribution of periodic zeta-functions. *Studia Scientiarum Mathematicarum Hungarica*, **48**(2):257–279, 2011. <http://dx.doi.org/10.1556/SScMath.48.2011.2.1162>.
- [7] A. Laurinčikas. Joint universality of zeta-functions with periodic coefficients. *Izvestiya: Mathematics*, **74**(3):515–539, 2010. <http://dx.doi.org/10.1070/IM2010v074n03ABEH002497>.
- [8] A. Laurinčikas. Universality of the Riemann zeta-function. *Journal of Number Theory*, **130**(10):2323–2331, 2010. <http://dx.doi.org/10.1016/j.jnt.2010.04.007>.
- [9] A. Laurinčikas. On joint universality of the Riemann zeta-function and Hurwitz zeta-functions. *Journal of Number Theory*, **132**(12):2842–2853, 2012. <http://dx.doi.org/10.1016/j.jnt.2012.05.026>.
- [10] A. Laurinčikas. Universality of composite functions. In K. Matsumoto et al.(Ed.), *Functions in Number Theory and Their Probabilistic Aspects*, volume B34, pp. 191–204, RIMS, 2012. RIMS Kōkyūroku Bessatsu.
- [11] A. Laurinčikas and D. Šiaučiūnas. A mixed joint universality theorem for zeta-functions. III. In A. Laurinčikas et al.(Ed.), *Analytic Probab. Methods Number Theory*, pp. 185–195, Vilnius, 2012. TEV.
- [12] R. Macaitienė. On joint universality for zeta-functions of newforms and periodic Hurwitz zeta-functions. In K. Matsumoto et al.(Ed.), *Functions in Number Theory and Their Probabilistic Aspects*, volume B34, pp. 217–233, RIMS, 2012. RIMS Kōkyūroku Bessatsu.
- [13] S.N. Mergelyan. Uniform approximation to functions of complex variable. *Uspehi Mat. Nauk (in Russian)*, **7**(48):31–122, 1952.
- [14] H. Mishou. The joint value-distribution of the Riemann zeta-function and Hurwitz zeta-functions. *Lithuanian Mathematical Journal*, **47**(1):32–47, 2007. <http://dx.doi.org/10.1007/s10986-007-0003-0>.
- [15] V. Pocevičienė and D. Šiaučiūnas. A mixed joint universality theorem for zeta-functions. II. *Mathematical Modelling and Analysis*, **19**(1):52–65, 2014. <http://dx.doi.org/10.3846/13926292.2014.893261>.
- [16] S.M. Voronin. Theorem on the 'universality' of the Riemann zeta-function. *Mathematics of the USSR Izvestiya*, **9**(3):443–453, 1975. <http://dx.doi.org/10.1070/IM1975v009n03ABEH001485>.
- [17] J.L. Walsh. *Interpolation and Approximation by Rational Functions in the Complex Domain*. American Mathematical Society Colloquium Publications, Vol. 20, 1968.