

Contents lists available at [ScienceDirect](https://www.sciencedirect.com)

International Journal for Parasitology: Parasites and Wildlife

journal homepage: www.elsevier.com/locate/ijppaw

Subtle transcriptomic response of Eurasian perch (*Perca fluviatilis*) associated with *Triaenophorus nodulosus* plerocercoid infection

Konrad Taube^{a,*}, Kristina Noreikiene^{a,b,1}, Siim Kahar^a, Riho Gross^a, Mikhail Ozerov^c, Anti Vasemägi^{a,d,1}

^a Chair of Aquaculture, Estonian University of Life Sciences, Kreutzwaldi 46a, 51014 Tartu, Estonia

^b Institute of Biosciences, Life Sciences Center, Vilnius University Vilnius, Lithuania

^c Biodiversity Unit, University of Turku, Vesilinnantie 5, 20500 Turku, Finland

^d Swedish University of Agricultural Sciences, Sötvattenslaboratoriet, Stångholmsvägen 2, 17893 Drottningholm, Sweden

ARTICLE INFO

Keywords:

Triaenophorus nodulosus
Parasites
Transcriptome
Helminth
RNAseq
Platyhelminthes
Liver parasites

ABSTRACT

Determining the physiological effects of parasites and characterizing genes involved in host responses to infections are essential to improving our understanding of host-parasite interactions and their ecological and evolutionary consequences. This task, however, is complicated by high diversity and complex life histories of many parasite species. The use of transcriptomics in the context of wild-caught specimens can help ameliorate this by providing both qualitative and quantitative information on gene expression patterns in response to parasites in specific host organs and tissues. Here, we evaluated the physiological impact of the widespread parasite, the pike tapeworm (*Triaenophorus nodulosus*), on its second intermediate host, the Eurasian perch (*Perca fluviatilis*). We used an RNAseq approach to analyse gene expression in the liver, the target organ of *T. nodulosus* plerocercoids, and spleen which is one of the main immune organs in teleost fishes. We compared perch collected from multiple lakes consisting of individuals with (n = 8) and without (n = 6) *T. nodulosus* plerocercoids in the liver. Results revealed a small number of differentially expressed genes (DEGs, adjusted p-value ≤ 0.05) in both spleen (n = 22) and liver (n = 10). DEGs in spleen consisted of mostly upregulated immune related genes (e.g., *JUN*, *SIK1*, *THS1*), while those in the liver were often linked to metabolic functions (e.g., *FABP1*, *CADM4*, *CDAB*). However, Gene Ontology (GO) analysis showed lack of functional enrichment among DEGs. This study demonstrates that Eurasian perch displays a subtle response at a gene expression level to *T. nodulosus* plerocercoid infection. Given that plerocercoids are low-metabolic activity transmission stages, our results suggest that moderate *T. nodulosus* plerocercoid infection most likely does not provoke an extensive host immune response and have relatively low physiological costs for the host. Our findings illustrate that not all conspicuous infections have severe effects on host gene regulation.

1. Introduction

Mounting an appropriate response to a parasite infection is essential for host's fitness and survival prospects. The magnitude of such a response, involving immediate immune response, inflammation, or wound repair, is expected to be influenced by many factors, including the parasite, environmental variables, and individual host characteristics (Cox, 2001; Gopko et al., 2017; Turner et al., 2021). The use of next-generation sequencing (NGS) technologies improves our ability to

understand the molecular basis of such complex host-parasite interactions, even in the wild (Brinker and Hamers, 2007; Greenwood et al., 2016; Sudhagar et al., 2018). For instance, by characterizing tissue-specific gene expression, we can gain novel insights into the genes that are likely involved in the immune responses or tissue repair of the host in response to parasite infection (Shanmugam et al., 2012; Hoy et al., 2014; Anderson et al., 2015; Shadab et al., 2019). Consequently, this provides targets for future investigation, and even discoveries of potential anti-parasitic drug targets for various ecologically or

* Corresponding author.

E-mail addresses: konrad.taube@emu.ee (K. Taube), kriste.noreikiene@gmail.com (K. Noreikiene), siim.kahar@emu.ee (S. Kahar), riho.gross@emu.ee (R. Gross), mikoze@utu.fi (M. Ozerov), anti.vasemagi@slu.se (A. Vasemägi).

¹ Kristina Noreikiene and Anti Vasemägi contributed equally to this work.

<https://doi.org/10.1016/j.ijppaw.2023.09.009>

Received 10 July 2023; Received in revised form 20 September 2023; Accepted 21 September 2023

Available online 29 September 2023

2213-2244/© 2023 The Authors. Published by Elsevier Ltd on behalf of Australian Society for Parasitology. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

economically important species (Chandhini and Rejish Kumar, 2019; Ahmad et al., 2021a,b).

The Eurasian perch (*Perca fluviatilis*) is a widely distributed freshwater fish species native to Eurasia throughout Europe to Eastern Siberia (Behrmann-Godel and Brinker, 2016). It is a generalist feeder and has a rich parasite fauna (Kuperman, 1973; Balling and Pfeiffer, 1997; Valtonen et al., 2003; Lahnsteiner et al., 2009) hosting nearly 150 parasite species (Behrmann-Godel and Brinker, 2016). Despite a large and increasing body of research on perch biology, ecology, and genetics (Thorpe, 1977; Acerete et al., 2004; Vasemägi et al., 2023), relatively little is known about perch–parasite interactions at a molecular level; however, this situation is slowly changing, at least in part because of growing importance of Eurasian perch in aquaculture (Fontaine and Teletchea, 2019) and an increasing availability of molecular resources (Malmström et al., 2017; Roques et al., 2020; Ozerov et al., 2022). Yet, the ecological relevance and physiological impact of the majority of parasite infections on perch are currently not known, which underlines the need for performing studies outside of a contained laboratory setting (Sasser and Weber, 2023).

One of the common parasites of Eurasian perch is the pike tapeworm (*Triaenophorus nodulosus*), a widespread parasitic helminth (Cestoda) which frequently uses *P. fluviatilis* as a second intermediate host (Kuperman, 1973) to infect its definitive host, the northern pike (*Esox lucius*) (Bregazzi, 2006). Commonly, perch are infected by *T. nodulosus* as fry and juveniles by feeding on infected zooplankton (e.g., *Cyclops*, *Daphnia*, *Diaptomus*, and others (Lahnsteiner et al., 2009)). The ingested parasite migrates through the intestine of the perch and punctures the intestinal wall to reach the targeted liver tissue, where it encysts and develops into the plerocercoid phase. The parasite's migration through tissues can result of the lysis of host membranes, causing inflammation, atrophy, necrosis, and other potentially lethal consequences (Rosen, 1918; Scheuring, 1922; Kuperman, 1973; Hoffmann et al., 2006; Behrmann-Godel and Brinker, 2016). Damage to the host may be inflicted also at a later infection stage from the increased pressure on the surrounding tissues by large, encapsulated larvae which continue growth and development within the liver (Scheuring, 1922). The severity of *T. nodulosus* infection was shown to depend on parasite load, which past research has found to be between 2 and 3 plerocercoids per infected *P. fluviatilis* liver (Morley and Lewis, 2010, 2017), where the condition of the perch deteriorates with increasing parasite numbers (Behrmann-Godel and Brinker, 2016). This density-dependent effect has been also observed in pumpkinseeds (*Lepomis gibbosus*), an invasive teleost in Europe that is often occupying similar niches as the perch (Brinker and Hamers, 2000; Masson et al., 2015). Infection with *T. nodulosus* in adult perch has been also associated with a reduction in fertility (Brinker and Hamers, 2007) and ~10% reduction in growth rate (Kuperman, 1973; Brinker and Hamers, 2000; Lahnsteiner et al., 2009; Behrmann-Godel and Brinker, 2016). However, until now, the physiological effect of *T. nodulosus* on *P. fluviatilis* has been studied only using histological techniques and based on a single gene (proliferating cell nuclear antigen) (Dezfuli et al., 2014). Thus, despite high prevalence and wide distribution of *T. nodulosus* (Brinker and Hamers, 2007; Borvinskaya et al., 2019), we currently lack detailed information on the molecular pathways and processes involved in host responses to the parasite which may be currently attained using whole-transcriptome sequencing.

The aim of this study was to evaluate the response of perch to *T. Nodulosus* infection by comparing the expression profiles of naturally and moderately infected and uninfected individuals in two tissues (spleen, liver) at the whole transcriptome level. We hypothesized that (i) once *T. nodulosus* has encysted and given that plerocercoids are a relatively low-metabolic activity stage of development of a parasite (Liu et al., 2022), host responses at the level of transcript abundances are expected to be mild to moderate. This is supported by earlier work reporting relatively weak physiological effects of plerocercoid infections (Brinker and Hamers, 2007; Behrmann-Godel and Brinker, 2016), which, in less severe cases, may enable perch to survive and withstand

infection by *T. nodulosus* plerocercoids for years in its natural environment. We also hypothesized that (ii) due to a vital role of liver in haematopoiesis, detoxification and nutritional metabolism, stronger transcriptomic response in liver compared to spleen would predominantly reflect the effect of parasite on host condition, feeding and response to physical damage. On the other hand, (iii) given the spleen's key role in infection response, a greater proportion of differentially expressed genes (DEGs) in the spleen compared to liver would indicate the importance of immune response of the host against the parasite.

2. Materials and methods

2.1. Fish collection and tissue preparation

In August 2021, fifteen adult *P. fluviatilis* individuals were caught using gillnets from four lakes in Estonia. Perch was euthanized with an overdose of benzocaine and measured for weight (± 0.1 g) and length (total length; TL, mm) (Table 1). After opening the fish using a sterile scalpel, liver and spleen were dissected, snap-frozen on dry ice while in the field and subsequently stored at -80°C . In addition, we recorded the maturation status (mature, immature) and sex of the individual. The national guidelines for the care and use of animals were followed per the Estonian Animal Protection Act, and fishing permits were obtained by the Estonian Ministry of the Environment (no. 10–1/21/18).

2.2. *T. nodulosus* molecular identification

Infection status was determined by visual identification of plerocercoids in fresh *P. fluviatilis* livers. Between 2 and 5 plerocercoids were detected on each infected fish liver (data not shown) and a single plerocercoid was dissected from each of the infected specimens for molecular conformation of the species. Plerocercoid samples were fixed in 96% ethanol and stored at -20°C . DNA was extracted from whole plerocercoids using a Dneasy Tissue Kit (Qiagen) according to the manufacturer's instructions. The concentration of isolated DNA was measured with a NanoDrop ND-2000 spectrophotometer (Thermo Scientific). Amplification of the *rmlL* ribosomal subunit gene (Brabec et al., 2015) was performed in 10 μL reactions to confirm the occurrence of *T. nodulosus*. Each 10 μL polymerase chain reaction (PCR) consisted of 5 μL of 2x Typeit Buffer (Qiagen); 0.5 μL of each primer (500 nM); 2 μL of DNA (100–150 ng total) and 2 μL of nuclease-free water. The primer sequences used for *rmlL* amplification were following: Cest16Sfgen (5'-TRCCTTTTGCATCATG-3') and Cest_16SRc (5'-AATAGATAAGAACCGACCTGGC-3') (Scholz et al., 2013). The thermocycler amplification protocol consisted of initial denaturation at 95°C for 15 min, followed by 40 cycles of 30 s at 94°C , 30 s at 54°C and 90 s at 72°C ; the final extension at 72°C was 10 min. Sequencing of PCR product using Sanger method was performed at the Institute of Genomics Core Facility, University of Tartu (Estonia) from both directions, and the 22 sequences (forward and reverse) were successfully merged manually. BLAST analysis of 11 consensus sequences indicated that the infecting worms were *T. nodulosus* (GenBank ID: KR780832.1, highest sequence similarity: 97%). Sequences produced in this study can be accessed at GenBank (OR065063-OR065071).

2.3. RNA extraction

Frozen liver tissue samples were mechanically crushed in liquid nitrogen using a steel mortar and pestle to produce a homogenized powder, while frozen spleen tissue was mechanically crushed using a Retsch Mixer Mill MM 400 (Retsch) (the tissue consistency of spleen samples was not conducive to mortar and pestle homogenization). Total RNA was extracted using a NucleoSpin RNA extraction kit (MACHERY-NAGEL, Duren, Germany). RNA sample concentrations were measured with a NanoDrop 2000 (ThermoFisher), and sample quality was evaluated using a TapeStation 2200 (Agilent Technologies).

Table 1

Summary information of analysed *P. fluviatilis* samples and tissues used for this study. Sex (F for female, M for male) and infection status, location, and total length are provided.

| Sample ID | Infection status | Lake name | Lake type | Latitude | Longitude | Tissue used | Sex | Total length (mm) |
|-----------|------------------|-----------|-----------|----------|-----------|---------------|-----|-------------------|
| 1 | Infected | Kasaritsa | Clear | 57.781 | 27.03 | Liver | M | 126 |
| 2 | Infected | Kasaritsa | Clear | 57.781 | 27.03 | Liver | F | 144 |
| 3 | Infected | Lasa | Humic | 57.919 | 25.789 | Liver | M | 141 |
| 4 | Infected | Lasa | Humic | 57.919 | 25.789 | Liver, spleen | F | 161 |
| 5 | Uninfected | Lasa | Humic | 57.919 | 25.789 | Liver, spleen | M | 167 |
| 6 | Uninfected | Lasa | Humic | 57.919 | 25.789 | Liver, spleen | F | 163 |
| 7 | Uninfected | Lasa | Humic | 57.919 | 25.789 | Spleen | M | 140 |
| 8 | Uninfected | Lasa | Humic | 57.919 | 25.789 | Spleen | M | 140 |
| 9 | Infected | Saadjärv | Clear | 58.538 | 26.656 | Liver, spleen | F | 134 |
| 10 | Infected | Saadjärv | Clear | 58.538 | 26.656 | Spleen | F | 129 |
| 11 | Uninfected | Saadjärv | Clear | 58.538 | 26.656 | Liver, spleen | F | 103 |
| 12 | Uninfected | Saadjärv | Clear | 58.538 | 26.656 | Liver, spleen | M | 103 |
| 13 | Uninfected | Udriku | Humic | 59.296 | 26.064 | Liver | M | 152 |
| 14 | Infected | Udriku | Humic | 59.296 | 26.064 | Liver, spleen | F | 143 |
| 15 | Infected | Udriku | Humic | 59.296 | 26.064 | Spleen | F | 133 |

2.3.1. Library preparation, sequencing, and quality assessment

Sequencing libraries were prepared at the Novogene Cambridge Science Park (Cambridge, United Kingdom) from 300 ng total RNA. After fragmentation and end repair, poly-A tail attachment, adapter ligation, size selection, amplification, and purification, sequencing (2 × 150 bp) was performed via NovaSeq 6000 (Illumina).

The raw sequence files in fastq format returned from Novogene were run through FastQC v0.11.9 (Andrews, 2010) both before and after trimming to check the overall quality of the reads. The trimming was performed by fastp v.0.20 (Chen et al., 2018). Only reads which were both longer than 50 bp and were of mean quality of 25 were processed further.

2.3.2. Alignment and mapping

The reads which passed the quality control were mapped on the reference genome of *Perca fluviatilis* (Roques et al., 2020) using hisat2 v.2.1.1 (Kim et al., 2019) using default parameters. The produced. sam files were subsequently processed with samtools v.1.10 (Li et al., 2009). Lastly, the mapped reads were extracted in bam format for subsequent differential expression analysis.

2.4. Differential expression

Analysis for differentially expressed genes (DEGs) was performed in R v.4.2.2 (R Core Team, 2022) using the DESeq2 v.1.36.0 (Love et al., 2014) package. When designing the DESeq dataset object, we considered infection status and population as fixed factors, and the DESeq model internally corrects for potential differences in library sizes. Genes with an adjusted *p*-value (*p*-adj) ≤ 0.05 (Benjamini and Hochberg, 1995) were considered as significantly differentially expressed (DEGs) between infected and uninfected individuals. Human orthologue genes were obtained via Ozerov et al. (2022), and DEGs without human orthologues were manually searched using BLAST.

2.4.1. Gene ontology (GO) analysis

GO enrichment was performed using Gorilla (Eden et al., 2009) separately for both liver and spleen samples. DEGs were compared with all expressed genes from each respective tissue.

3. Results

Sequencing resulted in an average of 45.1 and 47.8 million raw reads for every spleen and liver sample, respectively. After trimming and filtering, a total of 432 million clean reads were retained for spleen samples, with the average number of 39.3 million reads per sample (Supplementary file 1). A total of 465 million clean reads were retained for liver samples, with the average number of reads of 42.4 million per

sample (Supplementary file 2).

RNAseq analysis for spleen samples revealed 17, 246 genes being expressed (>1 counts per million), with only 22 genes being differentially expressed between infected and uninfected fish (*p*-adj ≤ 0.05; Table 2). A heatmap generated from DEGs revealed that samples were segregating based on infection status, except for a single infected sample which clustered among uninfected fish (Fig. 1A). All except one DEG showed upregulation in infected fish (2 × 2 contingency table, Fisher's exact test *P* = 0.0015). Identified DEGs were primarily encoding for membrane-bound proteins such as adhesion molecules and receptors (*GPIA*, *SYTL3*, *BTN2A1*), kinase binding (*MIDN*, *AP1B1*, *DUSP4*), cell death and tumorigenesis (*SIK1*, *THBS1*) as well as were uncharacterized or unknown genes (LOC120571416, LOC120570956, LOC120573523, LOC120557941, si:ch73-335121.4, LOC120555327) (Table 2). Tumour suppression and related oncogenes (*JUN*, *JUND*, *RDH12*) were also observed. The majority of DEGs in the spleen relate to host immune response, metabolism, stress response, and membrane-associated proteins (Table 2). However, formal Gene Ontology (GO) analysis showed lack of functional enrichment among DEGs, most likely because of limited number of DEGs (see Fig. 2).

RNAseq analysis for liver samples revealed 15, 377 expressed genes (>1 counts per million), with only ten genes being differentially expressed (DE) between infected and uninfected fish (*p*-adj ≤ 0.05; Table 3). Like the spleen, a heatmap generated from DE genes in liver revealed incomplete segregation of samples based on infection status (Fig. 1B). Lymphocyte and cell differentiation (*ACSF2*), galactose-specific lectin nattolectin, as well as binding proteins and cell adhesion molecules and their catalysts (*LMNA*, *FABP1*, *CADM4*, *CDAB*, *APBB1IP*, *FTCD*) were among DEGs. Two DEGs (*BTN3A2*, *NRG3B*) were related to transcription factors involved in the host adaptive immune response. In contrast to spleen, up- and downregulation of DEGs in liver occurred evenly (2 × 2 contingency table, Fisher's exact test *P* = 1). Similar to the spleen, GO analysis showed lack of functional enrichment among DEGs, most likely because of limited number of DEGs.

4. Discussion

Parasitic infections can have severe effects on host physiology, immune system function, survival, and performance (Bienvenu et al., 2010). Many populations of fish are at risk to a wide range of parasites (Dezfuli et al., 2014; Behrmann-Godel and Brinker, 2016; Gopko et al., 2017; Vollset et al., 2018) which can have broad implications on the health and well-being of the host. However, not all infections have a severe or detrimental effect on the host and its fitness. In this study, we found that *T. nodulosus* plerocercoid infection evoked very subtle host response on both spleen and liver transcriptomes of adult *P. fluviatilis*. This suggests that moderate *T. nodulosus* plerocercoid load likely has

Table 2

Summary statistics and gene information for 22 DEGs ($p\text{-adj} \leq 0.05$) in spleen between infected and uninfected *P. fluviatilis* individuals. N/A indicates unknown protein.

| Genbank ID | Base mean (Infected) | Base mean (Uninfected) | log2 Fold Change | p-value | p-adj | Gene name | Gene ID |
|-----------------|----------------------|------------------------|------------------|----------|-------------|---|---------------|
| PFLUV_G00160980 | 222.75 | 99.89 | 1.97 | 1.32E-11 | 1.23E-07 | uncharacterized LOC120571416 | N/A |
| PFLUV_G00160960 | 163.54 | 17.56 | 3.38 | 1.43E-11 | 1.23E-07 | uncharacterized LOC120570956 | N/A |
| PFLUV_G00137210 | 242.75 | 54.2 | 2 | 1.60E-10 | 9.15E-07 | midnolin-like | <i>MIDN</i> |
| PFLUV_G00146270 | 1608.68 | 460.33 | 1.61 | 3.43E-10 | 1.48E-06 | salt-inducible kinase 1 | <i>SIK1</i> |
| PFLUV_G00160970 | 89.23 | 6.63 | 4.17 | 1.66E-09 | 5.69E-06 | uncharacterized protein LOC120571417 isoform X1 | N/A |
| PFLUV_G00169820 | 178.7 | 23.66 | 3.01 | 6.68E-09 | 1.91E-05 | uncharacterized protein LOC120573523 | N/A |
| PFLUV_G00042630 | 250.63 | 79.28 | 1.9 | 3.10E-08 | 7.60E-05 | probable nuclear hormone receptor HR38 | N/A |
| PFLUV_G00114330 | 2619.87 | 884.36 | 1.68 | 4.11E-08 | 8.82E-05 | transcription factor AP-1 | <i>JUN</i> |
| PFLUV_G00234600 | 3792.66 | 1154.08 | 1.75 | 3.92E-07 | 0.000748084 | thrombospondin-1-like | <i>THBS1</i> |
| PFLUV_G00093590 | 1039.83 | 845 | 0.41 | 5.16E-07 | 0.000885785 | glucose-6-phosphate isomerase a | <i>GPIA</i> |
| PFLUV_G00165910 | 197.7 | 59.92 | 1.77 | 7.91E-07 | 0.001235261 | E3 ubiquitin-protein ligase-like | N/A |
| PFLUV_G00223220 | 208.77 | 86.83 | 1.05 | 3.05E-06 | 0.004362188 | arginine vasopressin-induced protein 1-like | <i>AVPO1</i> |
| PFLUV_G00214770 | 289.01 | 151.81 | 1.01 | 4.10E-06 | 0.005413646 | synaptotagmin-like protein 3 | <i>SYTL3</i> |
| PFLUV_G00257880 | 609.91 | 415.74 | 0.48 | 9.00E-06 | 0.01017912 | retinol dehydrogenase 12: like | <i>RDH12L</i> |
| PFLUV_G00277620 | 458.37 | 200.03 | 1.88 | 9.19E-06 | 0.01017912 | uncharacterized protein LOC120555327 | N/A |
| PFLUV_G00132110 | 484.34 | 248.1 | 0.88 | 9.48E-06 | 0.01017912 | transcription factor AP-1-like | <i>AP-1</i> |
| PFLUV_G00054610 | 435.84 | 132.77 | 1.71 | 1.53E-05 | 0.015501048 | nuclear receptor subfamily 4, group A, member 1 | <i>NR4A1</i> |
| PFLUV_G00004570 | 2519.13 | 1093.24 | 1.03 | 2.37E-05 | 0.022595861 | Unknown | N/A |
| PFLUV_G00139600 | 1160.74 | 524.36 | 1.17 | 3.04E-05 | 0.027506318 | transcription factor jun-D-like | <i>JUND</i> |
| PFLUV_G00051240 | 10,006.48 | 19,495.98 | -0.95 | 3.82E-05 | 0.032055553 | butyrophilin subfamily 2 member A1-like | <i>BTN2A1</i> |
| PFLUV_G00199970 | 1436.69 | 747.55 | 0.87 | 3.95E-05 | 0.032055553 | LOW QUALITY PROTEIN: dual specificity protein phosphatase 4 | <i>DUSP4</i> |
| PFLUV_G00008200 | 176.2 | 76.43 | 1.08 | 4.10E-05 | 0.032055553 | low density lipoprotein receptor b | <i>LDLRB</i> |

only a weak effect on host metabolism and condition.

Given that the liver is a major metabolic hub and the target tissue for *T. nodulosus* plerocercoids enduring physical abrasion during encysting, we anticipated that a pronounced transcriptome-wide response to liver infection would predominantly reflect the effect of parasite on host condition, feeding and response to physical damage. Instead, *T. nodulosus* infection was associated only with a subtle transcriptomic response in perch (number of DEGs = 10), which most likely reflects a weak effect of the parasite infection on host metabolism and physiological condition. Yet, several identified DEGs in liver (e.g., *FABP1*, *BT3A2*, *LMNA*, *CADM4*, *APBB1IP*) have been shown to associate with various parasite infections and liver ailments in other host-parasite systems; for example, Alvarez Rojas et al. (2015) found an increase in immune- and fibrosis-related genes as a result of trematode *Fasciola hepatica* infection in sheep liver. Similarly, one of the upregulated genes in perch liver, *LMNA* (LOC120572191), provides increased structural integrity for the cell as component of the fibrous nuclear lamina, and has been shown to be involved with liver fibrosis (Wang et al., 2016). We also found down-regulation of inhibitors associated with wound healing

and cell death (*APBB1IP*, *CADM4*) and upregulation of neutrophil chemotaxis genes and transcription factors (*CDAB*, *Galactose-specific lectin nattectin*) associated with host immune response.

Due to the spleen's key role in immune defence, we anticipated that substantially higher proportion of DEGs in the spleen compared to liver would reflect stronger immune response of the host against the parasite. Yet, we found similar, very small number of DEGs in the spleen ($n = 22$) among which there were several genes without known human orthologs ($n = 8$). The lack of strong transcriptome-wide response may stem from a couple of processes. One such possibility is that *T. nodulosus* infection has an overall low physiological effect which does not provoke a strong host immune response and fish are able to cope with low to moderate levels of parasite burden. In contrast, many parasite infections often cause major changes in specific host tissues involving hundreds or thousands of genes reflecting a major systemic response; for example, over 200 DEGs and a strong immune functioning GO process enrichment was observed in perch eye tissues in association with *Diplostomidae* infections (Noreikiene et al., 2020). It is also possible that *T. nodulosus* is able to evade triggering the host immune system, such as the spleen or




Fig. 1. Differentially expressed genes between infected and uninfected *P. fluviatilis* in a) spleen and b) liver tissues. Filled-in and empty boxes on the top of each plot represent infected and uninfected individuals, respectively. N/A indicates unknown protein.




Fig. 2. MA and volcano plots comparing infected and uninfected spleen samples (a & b) and liver samples (c & d). In the MA plots (a & c), the log counts and the log fold change are represented on the x- and y-axis, respectively. For each volcano plot (b & d), log fold change is represented on the x-axis and the $-\log_{10}$ p-value on the y-axis, respectively. Positive fold change corresponds to upregulated genes in infected individuals.

even other immune-privileged sites (Chulanetra and Chaicumpa, 2021). For example, trematodes which can coat their surfaces in a complex which degrades host antibodies and thus allows the parasite to avoid the host’s immune response (Cortés et al., 2017; Chulanetra and Chaicumpa, 2021). *Fasciola hepatica* has also been shown to evade host immunity, despite drugs and other procedures such as vaccination and chemotherapy (Flynn and Musah-Eroje, 2020). Finally, the subtle gene response in both liver and spleen is consistent with low-metabolic

activity plerocercoid infections after host has been able to encapsulate and isolate the parasite. Therefore, it is likely that the major interactions between the host and *T. nodulosus* takes place at the time of early invasion, rather than during the established resident plerocercoid stage. Future studies should therefore focus on early infection stages of *T. nodulosus*, when the effects of parasite migration and establishment is likely more severe.

Despite this low number of DEGs, they indicate some degree of

Table 3Summary statistics and gene information for ten DEGs (p -adj ≤ 0.05) in liver between infected and uninfected *P. fluviatilis* individuals. N/A indicates unknown protein.

| Genbank ID | Base mean (Infected) | Base mean (uninfected) | log2 Fold Change | p-value | p-adj | Gene name | Gene ID |
|-----------------|----------------------|------------------------|------------------|----------|----------|---|---------|
| PFLUV_G00169300 | 111.07 | 636.76 | -2.46 | 1.32E-10 | 1.99E-06 | lamin-A-like | LMNA |
| PFLUV_G00169310 | 20.67 | 393.12 | -3.04 | 2.75E-07 | 0.00207 | butyrophilin subfamily 3 member A2-like isoform X2 | BT3A2 |
| PFLUV_G00177580 | 37.58 | 2.26 | 6.02 | 4.19E-06 | 0.018592 | medium-chain acyl-CoA ligase ACSF2: mitochondrial-like isoform X1 | ACSF2 |
| PFLUV_G00150020 | 99.55 | 9776.31 | -6.83 | 4.93E-06 | 0.018592 | galactose-specific lectin natectin-like isoform X1 | N/A |
| PFLUV_G00072130 | 982.28 | 273.76 | 2.22 | 6.51E-06 | 0.019055 | fatty acid-binding protein, liver-type-like | FABP1 |
| PFLUV_G00087640 | 63.21 | 258.32 | -1.67 | 7.58E-06 | 0.019055 | cell adhesion molecule 4 | CADM4 |
| PFLUV_G00243290 | 95.47 | 15 | 2.88 | 9.77E-06 | 0.021038 | neuregulin 3 b | NRG3B |
| PFLUV_G00041980 | 116.83 | 390.21 | -1.68 | 1.34E-05 | 0.025163 | cytidine deaminase b | CDAB |
| PFLUV_G00257980 | 870.56 | 350.03 | 1.3 | 1.69E-05 | 0.028373 | amyloid beta (A4) precursor protein-binding, family B, member 1 interacting protein | APBB1IP |
| PFLUV_G00277100 | 5813.36 | 11,217.62 | -0.83 | 1.89E-05 | 0.028451 | formimidoyltransferasecyclodeaminase | FTCD |

impact on the host. For example, one of the uncovered DEGs (*JUN*) is an oncogenic transcription factor which has been associated with anti-apoptotic properties, immune response, and cell death (Wisdom, 1999; Lukey et al., 2016). Genes belonging to this family of transcription factors have been upregulated in mice, when infected by a helminth parasite (*Taenia crassiceps* or *Taenia solium*) (Morales-Montor et al., 2004). This present study has found both an increased expression of *AP-1*, *NR4A1*, and *THBS1* orthologs in infected perch. Infection in bovines by the tick-borne *Thallia annulata* and *Thallia parva* parasites has also been shown to result in an increased expression of the *AP-1* transcription factor and an associated upregulation of *JUN* protein. Recently it has been reported that mice infected with the liver parasite *Entamoeba histolytica* showed an upregulation of *NR4A1*, a transcription factor responsible for regulation macrophages in the host (Hoenow et al., 2022). Additionally, it has been discovered that when three-spined stickleback (*Gasterosteus aculeatus*) are infected by the helminth parasites *Diplostomum pseudopathaceum* and *Schistocephalus solidus*, one of the few genes that was upregulated because of the infection was the *THBS1* (Haase et al., 2016), which is associated with both cellular matrix and cell-cell interactions (Isenberg and Roberts, 2020). Finally, upregulation of *SIK1* in infected perch coincides with an analysis by Kang et al. (2020), who found a significant correlation between *SIK1* and a negative prognosis in patients suffering from diffuse large B-cell lymphoma. While speculative, this may be relevant for other parasites, such as various liver and blood flukes (*Opisthorchis viverrini*, *Opisthorchis felinus*, and *Clonorchis sinensis*) that have been categorized as Group 1 carcinogens by the World Health Organization (Feng and Cheng, 2017). Taken together, these studies demonstrate that several identified DEGs in spleen are likely involved in host-pathogen responses in diverse systems.

Prior research has shown that perch can survive and withstand infection by *T. nodulosus* plerocercoids for years in its natural environment (Brinker and Hamers, 2007). Our findings support this by identification of only a small total number of DEGs across two tissues and corroborates earlier work by Masson et al. (2015) which demonstrated a relatively weak physiological effect (hepatosomatic index, body condition index, and others) of *T. nodulosus* plerocercoid infection on adult *P. fluviatilis*. However, our results are to some extent at odds with Brinker and Hamers (2000), who found a stronger physiological effect (on growth) which was linked to the severity of *T. nodulosus* infection. A potential reason for this difference may be related to the varying infection tolerance across perch ontogeny coupled with the selective

disappearance of the weakest individuals. Here, we sampled mature Eurasian perch individuals that have already survived their most vulnerable and challenging stages of development. Thus, it is entirely possible we would see a more pronounced effect of *T. nodulosus* infection in juveniles that initially suffer plerocercoid formation and are expected to have competing resource demands between immune function (current survival) and growth (future survival). Considering the described effects during reproduction (Brinker and Hamers, 2007), it is also possible that a stronger response to infection would be observed during the breeding season, which is especially demanding for egg producing females (Šimková et al., 2005). Due to the random nature of catching fish in the wild, we were not able to control for the degree of infection status, and it may be the case that more heavily infected perch would exhibit more substantial response against *T. nodulosus*. However, prior research on *T. nodulosus* in the liver of *P. fluviatilis* has shown a mean intensity of 2.9 parasites (Morley and Lewis, 2017), which falls comfortably into the observed range of parasite load within the present study (between 2 and 5 plerocercoids per infected liver). The findings of Morley and Lewis (2017) are in agreement with other research which has found mean infection intensities of 3.87 plerocercoids per live (SD = 1.79) (Morozínska-Gogol, 2013), as well as having a ‘normal’ infection range being defined as 1–3 plerocercoids per liver, with severe infections with >3 plerocercoids (Brinker and Hamers, 2007).

Evidence across different host-parasite systems indicates that infection can increase the likelihood of predation (Furey et al., 2021; Turner et al., 2021), but this phenomenon is far from universal (Masson et al., 2015). To maximize the likelihood for reaching its final host (northern pike), the pike flatworm may benefit from weakening perch making it more susceptible to predation and thus completing its life cycle, but this has yet to be shown conclusively (Dezfuli et al., 2014). However, if the Eurasian perch is unable to sustain and survive given parasitic burden and dies before being eaten by the pike, it does not represent an optimal reproductive outcome for *T. nodulosus*. Rather, it is in the parasite’s interest that the intermediate host can cope with the infection and live until being predated by pike. Therefore, it makes sense *prima facie* that the more optimal strategy for the parasitic flatworm would be to induce only mild response in the intermediate host and our transcriptomic analysis corroborates this. Recent research into the parasitic nematode *Contracaecum osculatatum* revealed that fish with high infection densities of *C. osculatatum* had decreased nutritional condition as evidenced by reduced standard metabolic rate, reduction in the digestive organ masses, and changes in the plasma, body and liver composition, and fish

energy source (Ryberg et al., 2020). This effect on metabolism, immune function, and growth are in broad agreement with our own study and may suggest that liver-parasites induce the same general host response, despite differences in parasite species. Therefore, in a similar vein to *T. nodulosus*, it may be advantageous for *C. osculatum* to target growth and immune functions in Baltic cod (*Gadus morhua*) as a means to reach the final stage of its life cycle within its definitive host, the grey seal (*Halichoerus grypus*).

To our knowledge, this study provides first transcriptomic insights into the Eurasian perch response to pike tapeworm plerocercoid infection. Future investigation into this host-parasite system may benefit from an increase in sample size, and to also include analysis of severely infected individuals of different ages. RNAseq studies until now typically have employed a strategy of sequencing 3–5 individuals per treatment, while using an alternative library preparation methods (such as 3'-end sequencing, also known as QuantSeq) (e.g., Moll et al., 2014; Ahmad et al., 2021) are able to screen hundreds of samples; therefore, given the ongoing rapid development of next-generation sequencing technologies and dropping sequencing costs, future studies are likely able to carry out more comprehensive screening across different stages of infection. In this study our recording system was binary, in that a given liver tissue, when examined by eye, is recorded as either being infected or uninfected and the exact degree of infection severity was not taken into account. Therefore, subsequent research should include an explicit analysis of infection intensity as well as physiological condition, as for example in Baltic cod infected by the parasitic nematode *Contracaecum osculatum* (Marnis et al., 2019; Ryberg et al., 2020; Behrens et al., 2023). Such studies may provide additional insights into how parasite load is related to *P. fluviatilis* gene expression and immune response.

While the definition of 'parasite' commonly implies that one organism benefits at the expense of another organism (Olano et al., 2006), the actual cost of the parasitism may vary from lethal to essentially negligible. Hosts may harbour dozens to many hundreds of parasites, including many different parasite species. Therefore, it has been suggested that it is more fitting to describe the impact of being parasitized as range of effects which exist on a spectrum, whose ranges can extend from mild and virtually asymptomatic, to severe or even lethal (Ebert, 1998; Abernathy et al., 2011; Bracamonte et al., 2019; Barrett and Bartholomew, 2021; Hargitai et al., 2021; Hierweger et al., 2017; Sasser and Weber, 2023). However, it is likely that there exists a bias among current transcriptomic studies focusing on highly virulent or harmful parasites, as improving our understanding of most dangerous pathogens is often prioritized in medical, veterinary, or ecological research. Yet, given the incredible diversity of parasitic life-forms, it is likely that many parasites have only a small effect on host fitness at least during a certain period. Our study therefore represents one of such examples where a parasite, despite its conspicuous nature, evokes only a subtle physiological response from the host.

Note

Supplementary data associated with this article.

Funding

Support for the study was provided by Estonian Research Council grant (PRG852 to RG), Swedish Research Council grant 2020-03916 (to AV), and the Kristjan Jaak Foundation scholarship (to KT).

Availability of data and materials

Sequence data will be made available upon publication.

Authors' contributions

KT was responsible for laboratory work, data analysis, manuscript writing, figure and table construction, and data interpretation. KN contributed to the original study design, sample collection, manuscript writing, and data interpretation. AV contributed to the original study design, sample collection, manuscript writing, and data interpretation. SK contributed to sample collection. MO contributed to data analysis and data interpretation. RG contributed to manuscript editing. All authors read and approved the final manuscript.

Ethics approval and consent to participate

None.

Consent for publication

All authors give consent for publication.

Declaration of competing interest

The authors declare no conflict of interest.

Acknowledgements

We would like to thank the CSC-IT Center for Science at the Finnish Functional Genomics Center for access to their high-performance computing resource, Puhti. We would also like to thank Magnus Lauringsson and Alfonso Diaz-Suarez for their help in the field. Lastly, we would like to thank two anonymous reviewers for providing thoughtful feedback and suggestions during the peer-review process.

Appendix A. Supplementary data

Supplementary data to this article can be found online at <https://doi.org/10.1016/j.ijppaw.2023.09.009>.

References

- Abernathy, J., Xu, D.-H., Peatman, E., Kucuktas, H., Klesius, P., Liu, Z., 2011. Gene expression profiling of a fish parasite *Ichthyophthirius multifiliis*: insights into development and senescence-associated avirulence. *Comp. Biochem. Physiol. Genom. Proteomics* 6, 382–392.
- Acerete, L., Balasch, J.C., Espinosa, E., Josa, A., Tort, L., 2004. Physiological responses in Eurasian perch (*Perca fluviatilis*, L.) subjected to stress by transport and handling. *Aquaculture* 237, 167–178.
- Ahmad, F., Debes, P.V., Nousiainen, I., Kahar, S., Pukk, L., Gross, R., Ozerov, M., Vasemägi, A., 2021a. The strength and form of natural selection on transcript abundance in the wild. *Mol. Ecol.* 30, 2724–2737.
- Ahmad, F., Debes, P.V., Pukk, L., Kahar, S., Hartikainen, H., Gross, R., et al., 2021b. Know your enemy – transcriptome of myxozoan *Tetracapsuloides bryosalmonae* reveals potential drug targets against proliferative kidney disease in salmonids. *Parasitology* 148, 726–739.
- Alvarez Rojas, C.A., Ansell, B.R., Hall, R.S., Gasser, R.B., Young, N.D., Jex, A.R., et al., 2015. Transcriptional analysis identifies key genes involved in metabolism, fibrosis/tissue repair and the immune response against *Fasciola hepatica* in sheep liver. *Parasites Vectors* 8, 124.
- Anderson, R.M., Turner, H.C., Farrell, S.H., Yang, J., Truscott, J.E., 2015. What is required in terms of mass drug administration to interrupt the transmission of schistosome parasites in regions of endemic infection? *Parasites Vectors* 8, 553.
- Andrews, S., 2010. FastQC: a Quality Control Tool for High Throughput Sequence Data. <http://www.bioinformatics.babraham.ac.uk/projects/fastqc>.
- Balling, T.E., Pfeiffer, W., 1997. Frequency distributions of fish parasites in the perch *Perca fluviatilis* L. from Lake Constance. *Parasitol. Res.* 83, 370–373.
- Barrett, D.E., Bartholomew, J.L., 2021. A tale of two fish: comparative transcriptomics of resistant and susceptible steelhead following exposure to *Ceratonova shasta* highlights differences in parasite recognition. *PLoS One* 16, e0234837.
- Behrens, J.W., Ryberg, M.P., Chondromatidou, V., Iburg, T.M., 2023. Comparative histopathology of livers from Baltic cod infected with the parasitic nematode *Contracaecum osculatum*. *J. Fish. Dis.* 46, 653–662.

- Behrmann-Godel, J., Brinker, A., 2016. Biology and Ecology of Perch Parasites, vols. 193–229.
- Benjamini, Y., Hochberg, Y., 1995. Controlling the false discovery rate: a practical and powerful approach to multiple testing. *J. Roy. Stat. Soc. B* 57, 289–300.
- Bienvenu, A.L., Gonzalez-Rey, E., Picot, S., 2010. Apoptosis induced by parasitic diseases. *Parasites Vectors* 3, 106.
- Brabec, J., Kostadinova, A., Scholz, T., Littlewood, D.T.J., 2015. Complete mitochondrial genomes and nuclear ribosomal RNA operons of two species of Diplostomum (Platyhelminthes: trematoda): a molecular resource for taxonomy and molecular epidemiology of important fish pathogens. *Parasites Vectors* 8, 336.
- Bracamonte, S.E., Johnston, P.R., Monaghan, M.T., Knopf, K., 2019. Gene expression response to a nematode parasite in novel and native eel hosts. *Ecol. Evol.* 9, 13069–13084.
- Bregazzi, P., 2006. The biology of pike, *Esox lucius L.*, in a southern eutrophic lake. *J. Fish. Biol.* 17, 91–112.
- Brinker, A., Hamers, R., 2000. *Trianaeophorus procerus* n. sp.: a new tapeworm species or a misinterpretation of some unusual *Trianaeophorus nodulosus* Pallas, 1781 phenotypic characteristics. *Bull. Eur. Assoc. Fish Pathol.* 20, 256–260.
- Brinker, A., Hamers, R., 2007. Evidence for negative impact of plerocercoid infection of *Trianaeophorus nodulosus* on *Perca fluviatilis* L. stock in Upper Lake Constance, a water body undergoing rapid reoligotrophication. *J. Fish. Biol.* 71, 129–147.
- Chandhini, S., Rejish Kumar, V.J., 2019. Transcriptomics in aquaculture: current status and applications. *Rev. Aquacult.* 11, 1379–1397.
- Chen, S., Zhou, Y., Chen, Y., Gu, J., 2018. fastp: an ultra-fast all-in-one FASTQ preprocessor. *Bioinformatics* 34, i884–i890.
- Chulanetra, M., Chaicumpa, W., 2021. Revisiting the mechanisms of immune evasion employed by human parasites. *Front. Cell. Infect. Microbiol.* 11, 702125.
- Cortés, A., Sotillo, J., Muñoz-Antolí, C., Molina-Durán, J., Esteban, J.G., Toledo, R., 2017. Antibody trapping: a novel mechanism of parasite immune evasion by the trematode *Echinostoma caproni*. *PLoS Neglected Trop. Dis.* 11, e0005773.
- Cox, F.E., 2001. Concomitant infections, parasites and immune responses. *Parasitology* 122 (Suppl. 1), S23–38.
- Dezfuli, B.S., Giari, L., Lorenzoni, M., Manera, M., Noga, E.J., 2014. Perch liver reaction to *Trianaeophorus nodulosus* plerocercoids with an emphasis on piscidins 3, 4 and proliferative cell nuclear antigen (PCNA) expression. *Vet. Parasitol.* 200, 104–110.
- Ebert, D., 1998. Experimental evolution of parasites. *Science* 282 (5393), 1432–1436.
- Eden, E., Navon, R., Steinfeld, I., Lipson, D., Yakhini, Z., 2009. Gorilla: a tool for discovery and visualization of enriched GO terms in ranked gene lists. *BMC Bioinf.* 10, 48.
- Feng, M., Cheng, X., 2017. Parasite-associated cancers (blood flukes/liver flukes). *Adv. Exp. Med. Biol.* 1018, 193–205.
- Flynn, R.J., Musah-Eroje, M., 2020. Evasion of host immunity during *Fasciola hepatica* infection. *Methods Mol. Biol.* 2137, 107–115.
- Fontaine, P., Teletchea, F., 2019. Domestication of the eurasian perch (*Perca fluviatilis*). *Animal domestication. Intech.* <https://doi.org/10.5772/intechopen.85132>.
- Furey, N.B., Bass, A.L., Miller, K.M., Li, S., Lotto, A.G., Healy, S.J., et al., 2021. Infected juvenile salmon can experience increased predation during freshwater migration. *R. Soc. Open Sci.* 8 rsos.201522, 201522.
- Gopko, M., Mikheev, V., Taskinen, J., 2017. Positive density-dependent growth supports costs sharing hypothesis and population density sensing in a manipulative parasite. *Parasitology* 144, 1511–1518.
- Greenwood, J.M., Ezquerro, A.L., Behrens, S., Branca, A., Mallet, L., 2016. Current analysis of host–parasite interactions with a focus on next generation sequencing data. *Zoology* 119, 298–306.
- Haase, D., Rieger, J.K., Witten, A., Stoll, M., Bornberg-Bauer, E., Kalbe, M., et al., 2016. Comparative transcriptomics of stickleback immune gene responses upon infection by two helminth parasites, *Diplostomum pseudospathaceum* and *Schistocephalus solidus*. *Zoology* 119, 307–313.
- Hargitai, R., Pankovics, P., Boros, Á., Mátics, R., Altan, E., Delwart, E., Reuter, G., 2021. Novel picornavirus (family *Picornaviridae*) from freshwater fishes (*Perca fluviatilis*, *Sander lucioperca*, and *Ameiurus melas*) in Hungary. *Arch. Virol.* 166, 2627–2632.
- Hierweiger, M.M., Koch, M.C., Rupp, M., Maes, P., Di Paula, N., Bruggmann, R., Kuhn, J. H., et al., 2017. Novel filoviruses, hantavirus, and rhabdovirus in freshwater fish, Switzerland. *Emerg. Infect. Dis.* 27, 3082–3091, 2021.
- Hoenow, S., Yan, K., Noll, J., Groneberg, M., Casar, C., Lory, N.C., et al., 2022. The properties of proinflammatory Ly6Chi monocytes are differentially shaped by parasitic and bacterial liver infections. *Cells* 11, 2539.
- Hoffmann, R., Meder, J., Klein, M., Osterkornj, K., Negele, R., 2006. Studies on lesions caused by plerocercoids of *Trianaeophorus nodulosus* in some fish of an alpine lake, the Königsee. *J. Fish. Biol.* 28, 701–712.
- Hoy, A.M., Lundie, R.J., Ivens, A., Quintana, J.F., Nausch, N., Forster, T., et al., 2014. Parasite-derived microRNAs in host serum as novel biomarkers of helminth infection. *PLoS Neglected Trop. Dis.* 8, e2701.
- Isenberg, J.S., Roberts, D.D., 2020. THBS1 (Thrombospondin-1). *Atlas of Genetics and Cytogenetics in Oncology and Haematology*. Available from: <http://hdl.handle.net/2042/70774>. (Accessed 10 May 2023).
- Kang, J., Yao, P., Tang, Q., Wang, Y., Zhou, Y., Huang, J., 2020. Systematic analysis of competing endogenous RNA networks in diffuse large B-cell lymphoma and Hodgkin's lymphoma. *Front. Genet.* 11, 586688.
- Kim, D., Paggi, J.M., Park, C., Bennett, C., Salzberg, S.L., 2019. Graph-based genome alignment and genotyping with HISAT2 and HISAT-genotype. *Nat. Biotechnol.* 37, 907–915.
- Kuperman, B.I., 1973. Infection of young perch by the tapeworm *Trianaeophorus nodulosus*. *SIL Proceedings* 18, 1697–1704.
- Lahnsteiner, F., Kletzl, M., Weismann, T., 2009. The risk of parasite transfer to juvenile fishes by live copepod food with the example *Trianaeophorus crassus* and *Trianaeophorus nodulosus*. *Aquaculture* 295, 120–125.
- Li, H., Handsaker, B., Wysoker, A., Fennell, T., Ruan, J., Homer, N., Marth, G., Abecasis, G., Durbin, R., 2009. 1000 Genome project data processing subgroup. The Sequence Alignment/Map format and SAMtools. *Bioinformatics* 15, 2078–2079.
- Liu, S.N., Su, X.Y., Chen, W.Q., Yu, J.W., Li, J.R., Jiang, P., Cui, J., Wang, Z.Q., Zhang, X., 2022. Transcriptome profile of plerocercoid and adult developmental stages of the neglected medical tapeworm *Spioermetra erinacei* eupoai. *Acta Trop.* 232, 106483.
- Love, M.I., Huber, W., Anders, S., 2014. Moderated estimation of fold change and dispersion for RNA-seq data with DESeq2. *Genome Biol.* 15, 550.
- Lukey, M.J., Greene, K.S., Erickson, J.W., Wilson, K.F., Cerione, R.A., 2016. The oncogenic transcription factor c-Jun regulates glutamine expression and sensitizes cells to glutamine-targeted therapy. *Nat. Commun.* 7, 11321.
- Malmström, M., Matschiner, M., Tørrisen, O.K., Jakobsen, K.S., Jentoft, S., 2017. Whole genome sequencing data and de novo draft assemblies for 66 teleost species. *Sci. Data* 4, 160132.
- Marnis, H., Kania, P.W., Syahputra, K., Zuo, S., Dirks, R.P., Buchmann, K. Transcriptomic analysis of Baltic cod (*Gadus morhua*) liver infected with *Contracaecum osculatam* third stage larvae indicates parasitic effects on growth and immune response. *Fish Shellfish Immunol.* 2019;93:965–976.
- Masson, G., Vanacker, M., Fox, M.G., Beisel, J.-N., 2015. Impact of the cestode *Trianaeophorus nodulosus* on the exotic *Lepomis gibbosus* and the autochthonous *Perca fluviatilis*. *Parasitology* 142, 745–755.
- Moll, P., Ante, M., Seitz, A., Reda, T., 2014. QuantSeq 3' mRNA sequencing for RNA quantification. *Nat. Methods* 11, i–iii.
- Morales-Montor, J., Escobedo, G., Rodriguez-Dorantes, M., Téllez-Ascencio, N., Cebón, M.A., Larraalde, C., 2004. Differential expression of AP-1 transcription factor genes c-fos and c-jun in the helminth parasites *Taenia crassiceps* and *Taenia solium*. *Parasitology* 129, 233–243.
- Morley, N.J., Lewis, J.W., 2010. Consequences of an outbreak of columnaris disease (*Flavobacterium columnare*) to the helminth fauna of perch (*Perca fluviatilis*) in the Queen Mary reservoir, south-east England. *J. Helminthol.* 84, 186–192.
- Morley, N.J., Lewis, J.W., 2017. Influence of *Trianaeophorus nodulosus* plerocercoids (Cestoda: pseudophyllidae) on the occurrence of intestinal helminths in the perch (*Perca fluviatilis*). *J. Helminthol.* 91, 711–717.
- Morozińska-Gogol, J., 2013. Parasite communities of European perch, *Perca fluviatilis* L. (Actinopterygii: perciformes: percidae) from lake Iębsko (central coast, Poland). *Annals of Parasitology* 59, 89–98.
- Noreikiene, K., Ozerov, M., Ahmad, F., Kóiv, T., Kahar, S., Gross, R., et al., 2020. Humic-acid-driven escape from eye parasites revealed by RNA-seq and target-specific metabarcoding. *Parasites Vectors* 13, 433.
- Olano, J.P., Weller, P.F., Guerrant, R.L., Walker, D.H., 2006. Principles of Parasitism: Host–Parasite Interactions, vols. 1–12. Tropical Infectious Diseases. Elsevier.
- Ozerov, M., Noreikiene, K., Kahar, S., Huss, M., Huusko, A., Kóiv, T., Sepp, M., López, M. E., Gårdmark, A., Gross, R., Vasemägi, A., 2022. Whole-genome sequencing illuminates multifaceted targets of selection to humic substances in Eurasian perch. *Mol. Ecol.* 31, 2367–2383.
- R Core Team, 2022. R: A Language and Environment for Statistical Computing. <https://www.R-project.org/>.
- Roques, C., Zahm, M., Cabau, C., Klopp, C., Bouchez, O., Donnadieu, C., et al., 2020. A Chromosome-Scale Genome Assembly of the European Perch, *Perca fluviatilis*. https://www.ncbi.nlm.nih.gov/assembly/GCF_010015445.1.
- Rosen, F., 1918. Recherches sur le développement des Cestodes; 1. Le cycle évolutif des Bothriocéphales. *Bull. Soc. Neuchâtel. Sci. Nat.* 43, 1–47.
- Ryberg, M.P., Skov, P.V., Vendramin, N., Buchmann, K., Nielsen, A., Behrens, J.W., 2020. Physiological condition of Eastern Baltic cod, *Gadus morhua*, infected with the parasitic nematode *Contracaecum osculatam*. *Conservation Physiology* 8, coaa093.
- Sasser, K.T., Weber, J.N., 2023. A call for more ecologically and evolutionarily relevant studies of immune costs. *Evol. Ecol.* 37, 203–214.
- Scheuring, L., 1922. Studien an Fischparasiten; *Trianaeophorus nodulosus* (Pallas) Rud. und die durch ihn im Fischkörper hervorgerufenen pathologischen Veränderungen. *Zeitschr. Fisch. Hilfsw.* 21, 93–204.
- Scholz, T., De Chambrier, A., Kuchta, R., Littlewood, D.T., Waeschenbach, A., 2013. *Macrobothriotaenia ficta* (Cestoda: proteocephalidea), a parasite of sunbeam snake (*Xenopeltis unicolor*): example of convergent evolution. *Zootaxa* 3640, 485–499.
- Shadab, M., Das, S., Banerjee, A., Sinha, R., Asad, M., Kamran, M., et al., 2019. RNA-Seq revealed expression of many novel genes associated with *Leishmania donovani* persistence and clearance in the host macrophage. *Front. Cell. Infect. Microbiol.* 9, 17.
- Shanmugam, D., Ralph, S.A., Carmona, S.J., Crowther, G.J., Roos, D.S., Agüero, F., 2012. Integrating and Mining Helminth Genomes to Discover and Prioritize Novel Therapeutic Targets, pp. 43–59.
- Šimková, A., Jarkovský, J., Koubková, B., Baruš, V., Prokeš, M., 2005. Associations between fish reproductive cycle and the dynamics of metazoan parasite infection. *Parasitol. Res.* 95, 65–72.
- Sudhagar, A., Kumar, G., El-Matbouli, M., 2018. Transcriptome analysis based on RNA-Seq in understanding pathogenic mechanisms of diseases and the immune system of fish: a comprehensive rReview. *Int. J. Mol. Sci.* 19, 245.

- Thorpe, J.E., 1977. Daily ration of adult perch, *Perca fluviatilis* L. during summer in Loch Leven, Scotland. *J. Fish. Biol.* 11, 55–68.
- Turner, W.C., Kamath, P.L., van Heerden, H., Huang, Y.H., Barandongo, Z.R., Bruce, S.A., et al., 2021. The roles of environmental variation and parasite survival in virulence – transmission relationships. *R. Soc. Open Sci.* 8, 210088.
- Valtonen, E., Holmes, J., Aronen, J., Rautalahti, I., 2003. Parasite communities as indicators of recovery from pollution: parasites of roach (*Rutilus rutilus*) and perch (*Perca fluviatilis*) in Central Finland. *Parasitology* 126, S43–S52.
- Vasemägi, A., Ozerov, M., Noreikiene, K., López, M.E., Gårdmark, A., 2023. Unlocking the genome of perch – from genes to ecology and back again. *Ecol. Freshw. Fish.* <https://doi.org/10.1111/eff.12713>.
- Vollset, K.W., Qviller, L., Skår, B., Barlaup, B.T., Dohoo, I., 2018. Parasitic sea louse infestations on wild sea trout: separating the roles of fish farms and temperature. *Parasites Vectors* 11, 609.
- Wang, L., Liu, P., Wan, Z.Y., Huang, S.Q., Wen, Y.F., Lin, G., et al., 2016. RNA-Seq revealed the impairment of immune defence of tilapia against the infection of

- Streptococcus agalactiae* with simulated climate warming. *Fish Shellfish Immunol.* 55, 679–689.
- Wisdom, R., 1999. c-Jun regulates cell cycle progression and apoptosis by distinct mechanisms. *EMBO J.* 18, 188–197.

Glossary

- DEG*: Differentially expressed gene
PCR: Polymerase chain reaction
RNAseq: RNA Sequencing
TL: Total length
GO: Gene ontology