

Article

Generalized Limit Theorem for Mellin Transform of the Riemann Zeta-Function

Antanas Laurinčikas^{1,*} and Darius Šiaučiušas^{2,†}

¹ Institute of Mathematics, Faculty of Mathematics and Informatics, Vilnius University, Naugarduko Str. 24, LT-03225 Vilnius, Lithuania

² Institute of Regional Development, Šiauliai Academy, Vilnius University, Vytauto Str. 84, LT-76352 Šiauliai, Lithuania; darius.siauciunas@sa.vu.lt

* Correspondence: antanas.laurincikas@mif.vu.lt

† These authors contributed equally to this work.

Abstract: In the paper, we prove a limit theorem in the sense of the weak convergence of probability measures for the modified Mellin transform $\mathcal{Z}(s)$, $s = \sigma + it$, with fixed $1/2 < \sigma < 1$, of the square $|\zeta(1/2 + it)|^2$ of the Riemann zeta-function. We consider probability measures defined by means of $\mathcal{Z}(\sigma + i\varphi(t))$, where $\varphi(t)$, $t \geq t_0 > 0$, is an increasing to $+\infty$ differentiable function with monotonically decreasing derivative $\varphi'(t)$ satisfying a certain normalizing estimate related to the mean square of the function $\mathcal{Z}(\sigma + i\varphi(t))$. This allows us to extend the distribution laws for $\mathcal{Z}(s)$.

Keywords: modified Mellin transform; Riemann zeta-function; weak convergence of probability measures

MSC: 11M06



Citation: Laurinčikas, A.; Šiaučiušas, D. Generalized Limit Theorem for Mellin Transform of the Riemann Zeta-Function. *Axioms* **2024**, *13*, 251. <https://doi.org/10.3390/axioms13040251>

Academic Editors: Dongkyu Lim, Adem Kilicman and Arjun Kumar Rathie

Received: 13 March 2024

Revised: 6 April 2024

Accepted: 8 April 2024

Published: 10 April 2024



Copyright: © 2024 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

1. Introduction

Let $s = \sigma + it$ be a complex variable. One of the most important objects of the classical analytic number theory is the Riemann zeta-function $\zeta(s)$, which is defined, for $\sigma > 1$, by the Dirichlet series

$$\zeta(s) = \sum_{m=1}^{\infty} \frac{1}{m^s}.$$

Moreover, the function $\zeta(s)$ has analytic continuation to the region $\mathbb{C} \setminus \{1\}$, and the point $s = 1$ is its simple pole with residue 1. The first value distribution results for $\zeta(s)$ with real s were obtained by Euler. Riemann was the first mathematician who began to study [1] $\zeta(s)$ with complex variables, proved the functional equation for $\zeta(s)$, obtained its analytic continuation, proposed a means of using $\zeta(s)$ for the investigation of the asymptotic prime number distribution law

$$\pi(x) = \sum_{p \leq x} 1, \quad x \rightarrow \infty,$$

and stated some hypotheses on $\zeta(s)$. The most important hypothesis, now called the Riemann hypothesis, states that all zeros of $\zeta(s)$ in the region $\sigma \geq 0$ are located on the line $\sigma = 1/2$. Riemann's ideas concerning $\pi(x)$ were correct, and Hadamard [2] and de la Vallée Poussin [3], using them, independently proved that

$$\lim_{x \rightarrow \infty} \pi(x) \left(\int_2^x \frac{du}{\log u} \right)^{-1} = 1.$$

However, the Riemann hypothesis remains open at present; it is among the seven Millennium Problems of mathematics [4]. In the theory of $\zeta(s)$, there are other important problems. One of them is connected to the asymptotics of moments

$$M_k(\sigma, T) \stackrel{\text{def}}{=} \int_0^T |\zeta(\sigma + it)|^{2k} dt, \quad k > 0, \sigma \geq \frac{1}{2},$$

as $T \rightarrow \infty$. For example, at the moment the asymptotics of $M_k(\sigma, T)$, $\sigma = 1/2$ is known only for $k = 1$ and $k = 2$; see [5]. For the investigation of $M_k(\sigma, T)$, Motohashi proposed (see [6,7]) to use the modified Mellin transforms

$$\mathcal{Z}_k(s) = \int_1^\infty \left| \zeta\left(\frac{1}{2} + ix\right) \right|^{2k} x^{-s} dx, \quad k \in \mathbb{N}.$$

Let $g(x)$ be a certain function, e.g., $g(x)x^{\sigma-1} \in L(0, \infty)$, and

$$G(s) = \int_0^\infty g(x)x^{s-1} dx.$$

Then, using the Mellin inverse formula leads to the following equality (see [8]):

$$\int_1^\infty g\left(\frac{x}{T}\right) \left| \zeta\left(\frac{1}{2} + ix\right) \right|^{2k} dx = \frac{1}{2\pi i} \int_{c-i\infty}^{c+i\infty} G(s) T^s \mathcal{Z}_k(s) ds$$

with a certain $c > 1$. This shows that a suitable choice of the function $g(x)$ reduces investigations of $M_k(1/2, T)$ to those of properties of $\mathcal{Z}_k(s)$. The latter assertion inspired the creation of the analytic theory of the functions $\mathcal{Z}_k(s)$.

In this paper, we limit ourselves to the probabilistic value distribution of the function $\mathcal{Z}(s) \stackrel{\text{def}}{=} \mathcal{Z}_1(s)$ only. Before this, we recall some known results of the function $\mathcal{Z}(s)$.

Let $\gamma = 0.577 \dots$ denote the Euler constant and $E(T)$ be defined by

$$\int_0^T \left| \zeta\left(\frac{1}{2} + it\right) \right|^2 dt = T \log \frac{T}{2\pi} + (2\gamma - 1)T + E(T).$$

Moreover, let

$$F(t) = \int_1^t E(t) dt - \pi T \quad \text{and} \quad F_1(T) = \int_1^T F(t) dt.$$

The analytic behavior of the function $\mathcal{Z}(s)$ was described in [9] and forms the following theorem.

Theorem 1 ([9]). *The function $\mathcal{Z}(s)$ is analytically continuable to the region $\sigma > -3/4$, except the point $s = 1$, which is a double pole, and*

$$\mathcal{Z}(s) = \frac{1}{(s-1)^2} + \frac{2\gamma - \log 2\pi}{s-1} - E(1) + \pi(s+1) + s(s+1)(s+2) \int_1^\infty F_1(x)x^{-s-3} dx.$$

Moreover, the estimates

$$\mathcal{Z}(\sigma + it) \ll_\epsilon t^{1-\sigma+\epsilon}, \quad 0 \leq \sigma \leq 1, t \geq t_0 > 0,$$

and

$$\int_1^T |\mathcal{Z}(\sigma + it)|^2 dt \ll_\epsilon \begin{cases} T^{3-4\sigma+\epsilon} & \text{if } 0 \leq \sigma \leq 1/2, \\ T^{2-2\sigma+\epsilon} & \text{if } 1/2 \leq \sigma \leq 1, \end{cases} \tag{1}$$

are valid.

Here and in what follows, ϵ is an arbitrary fixed positive number that is not always the same, and the notation $x \ll_\epsilon y$, $x \in \mathbb{C}$, $y > 0$, means that there is a constant $c = c(\epsilon) > 0$ such that $|x| \leq cy$.

In [10], Bohr proposed to characterize the asymptotic behavior of the Riemann zeta-function by using a probabilistic approach. This idea is acceptable because the value distribution of $\zeta(s)$ is quite chaotic. Denote by JA the Jordan measure of the set $A \subset \mathbb{R}$. Then, Bohr, jointly with Jessen, roughly speaking, obtained in [11,12] that, for $\sigma > 1/2$ and every rectangle $R \subset \mathbb{C}$ with edges parallel to the axes, there exists a limit

$$\lim_{T \rightarrow \infty} J\{t \in [0, T] : \zeta(\sigma + it) \in R\}.$$

In modern terminology, the Bohr–Jessen theorem is stated as a limit theorem on weakly convergent probability measures. Let $\mathcal{B}(\mathbb{X})$ stand for the Borel σ -field of the space \mathbb{X} (in general, topological), and let P_n , $n \in \mathbb{N}$, and P be probability measures defined on $(\mathbb{X}, \mathcal{B}(\mathbb{X}))$. By this definition, P_n converges weakly to P as $n \rightarrow \infty$ ($P_n \xrightarrow[n \rightarrow \infty]{w} P$) if

$$\lim_{n \rightarrow \infty} \int_{\mathbb{X}} g dP_n = \int_{\mathbb{X}} g dP$$

for every real continuous bounded function g on \mathbb{X} . Let $\mathbf{L}A$ stand for the Lebesgue measure of a measurable set $A \subset \mathbb{R}$. Then, the modern version of the Bohr–Jessen theorem is of the following form: for every fixed $\sigma > 1/2$, there exists a probability measure P_σ on $(\mathbb{C}, \mathcal{B}(\mathbb{C}))$ such that

$$\frac{1}{T} \mathbf{L}\{t \in [0, T] : \zeta(\sigma + it) \in A\}, \quad A \in \mathcal{B}(\mathbb{C}),$$

converges weakly to P_σ as $T \rightarrow \infty$.

The first probabilistic limit theorems for the function $\mathcal{Z}(s)$ were discussed in [13]. For $A \in \mathcal{B}(\mathbb{C})$, set

$$Q_{T,\sigma}(A) = \frac{1}{T} \mathbf{L}\{t \in [0, T] : \mathcal{Z}(\sigma + it) \in A\}.$$

Assuming that $\sigma > 1/2$, it was obtained that there is a probability measure Q_σ on $(\mathbb{C}, \mathcal{B}(\mathbb{C}))$ such that $Q_{T,\sigma} \xrightarrow[T \rightarrow \infty]{w} Q_\sigma$. On the other hand, for every $\kappa > 0$, we have

$$\frac{1}{T} \mathbf{L}\{t \in [0, T] : |\mathcal{Z}(\sigma + it)| \geq \kappa\} \leq \frac{1}{\kappa T} \int_0^T |\mathcal{Z}(\sigma + it)| dt \leq \frac{1}{\kappa} \left(\frac{1}{T} \int_0^T |\mathcal{Z}(\sigma + it)|^2 dt \right)^{1/2}.$$

This, together with Theorem 1, implies that, for $1/2 < \sigma < 1$,

$$\lim_{T \rightarrow \infty} \frac{1}{T} \mathbf{L}\{t \in [0, T] : |\mathcal{Z}(\sigma + it)| \geq \kappa\} = 0.$$

The latter equality remains valid also for $\sigma > 1$. Thus, the limit measure Q_σ is degenerated at the point $s = 0$. In order to avoid this situation, we propose to consider $\mathcal{Z}(\sigma + i\varphi(t))$ with a certain function $\varphi(t)$. Moreover, it is more convenient to deal with $t \in [T, 2T]$ because, in this case, additional restrictions for $\varphi(t)$ with $t = 0$ are not needed.

Denote

$$I_\sigma(T) = \int_1^T |\mathcal{Z}(\sigma + it)|^2 dt.$$

We suppose that $\varphi(t)$ is a positive increasing to $+\infty$ differentiable function with a monotonically decreasing derivative, such that

$$\frac{I_{\sigma-\varepsilon}(\varphi(T))}{\varphi'(T)} \ll T, \quad T \rightarrow \infty.$$

The class of such functions $\varphi(t)$ is denoted by W_σ . Consider the weak convergence for

$$P_{T,\sigma}(A) = \frac{1}{T} \mathbf{L}\{t \in [T, 2T] : \mathcal{Z}(\sigma + i\varphi(\tau)) \in A\}, \quad A \in \mathcal{B}(\mathbb{C}).$$

In this case, we have, by $\varepsilon \rightarrow 0$, that

$$\frac{I_\sigma(\varphi(T))}{\varphi'(T)} \ll T,$$

and

$$\frac{1}{T} \int_T^{2T} |\mathcal{Z}(\sigma + i\varphi(t))|^2 dt = \frac{1}{T} \int_{\varphi(T)}^{\varphi(2T)} \frac{1}{\varphi'(t)} |\mathcal{Z}(\sigma + iu)|^2 du \leq \frac{1}{T\varphi'(2T)} I_\sigma(\varphi(2T)) \ll 1 \quad (2)$$

for $\varphi(t) \in W_\sigma$. Thus, we cannot claim that the limit measure for $P_{T,\sigma}$ is degenerated at zero.

Now, we state a limit theorem for $P_{T,\sigma}$.

Theorem 2. *Assume that $\sigma \in (1/2, 1)$ is a given fixed number, and $\varphi(t) \in W_\sigma$. Then, on $(\mathbb{C}, \mathcal{B}(\mathbb{C}))$, there exists a probability measure P_σ such that $P_{T,\sigma} \xrightarrow[T \rightarrow \infty]{w} P_\sigma$.*

In virtue of Theorem 1, we see that

$$I_{\sigma-\varepsilon}(T) \ll T^{\alpha_\sigma}$$

with certain $0 < \alpha_\sigma < 1$. Take $\varphi(t) = (\log t)^{\beta_\sigma}$, $t \geq 2$, $\beta_\sigma > 0$. Then, $\varphi'(t)$ is decreasing, and

$$\frac{I_{\sigma-\varepsilon}(\varphi(T))}{\varphi'(T)} \ll T(\log T)^{\alpha_\sigma\beta_\sigma+1-\beta_\sigma} \ll T$$

if we choose

$$\beta_\sigma = (1 - \alpha_\sigma)^{-1}.$$

This shows that $(\log T)^{\beta_\sigma}$ is an element of the class W_σ .

Theorem 2 shows that the asymptotic behavior of the function $\mathcal{Z}(s)$ on vertical lines is governed by a certain probabilistic law, and this confirms the chaos in its value distribution. Moreover, Theorem 2 is an example of the application of probability methods in analysis. Thus, it continues a tradition initiated in works [11,12] and developed by Selberg [14], Joyner [15], Bagchi [16], Korolev [17,18], Kowalski [19], Lamzouri, Lester and Radziwill [20,21], Steuding [22], and others; see also a survey paper [23]. We note that a generalization of Theorem 2 for the functional spaces can be applied for approximation problems of some classes of functions.

We divide the proof of Theorem 2 into several parts. First, we discuss weak convergence on a certain group. The second part is devoted to the case related to an integral. Further, we consider a measure defined by an absolutely convergent improper integral. In the last part, Theorem 2 is proven. For proofs of all assertions on weak convergence,

the notions of relative compactness as well as of tightness and convergence in distribution are employed.

2. Fourier Transform Method

Let $b > 1$ be a fixed finite number, and

$$\mathbb{I}_b = \prod_{x \in [1,b]} \{s \in \mathbb{C} : |s| = 1\}.$$

The Cartesian product \mathbb{I}_b consists of all functions $i : [1, b] \rightarrow \{s \in \mathbb{C} : |s| = 1\}$. On \mathbb{I}_b , the product topology and operation of pointwise multiplication can be defined. This reduces \mathbb{I}_b to a compact topological group. We will give a limit lemma for probability measures on $(\mathbb{I}_b, \mathcal{B}(\mathbb{I}_b))$.

For $A \in \mathcal{B}(\mathbb{I}_b)$, put

$$V_{T,b}(A) = \frac{1}{T} \mathbf{L} \left\{ t \in [T, 2T] : \left(x^{-i\varphi(t)} : x \in [1, b] \right) \in A \right\}.$$

Lemma 1. *Suppose that the function $\varphi(t)$ has a monotonically decreasing derivative $\varphi'(t)$ such that*

$$(\varphi'(T))^{-1} = o(T), \quad T \rightarrow \infty. \tag{3}$$

Then $V_{T,b}$ converges weakly to a certain probability measure V_b as $T \rightarrow \infty$.

Proof. We use the Fourier transform approach. Denote the elements of \mathbb{I}_b by $i = \{i_x : x \in [1, b]\}$. Then, the Fourier transform $f_{T,b}(\underline{k}), \underline{k} = (k_x : k_x \in \mathbb{Z}, x \in [1, b])$ of the measure $V_{T,b}$ is the integral

$$f_{T,b}(\underline{k}) = \int_{\mathbb{I}_b} \left(\prod_{x \in [1,b]} i_x^{k_x} \right) dV_{T,b},$$

where only a finite number of integers k_x are not zeros. Therefore, the definition of $V_{T,b}$ yields

$$f_{T,b}(\underline{k}) = \frac{1}{T} \int_T^{2T} \left(\prod_{x \in [1,b]} x^{-ik_x \varphi(t)} \right) dt = \frac{1}{T} \int_T^{2T} \exp \left\{ -i\varphi(t) \sum_{x \in [1,b]} k_x \log x \right\} dt. \tag{4}$$

For brevity, let $A_b(\underline{k}) = \sum_{k \in [1,b]} k_x \log x$. Then, the second mean value theorem, (4), and (3) give

$$\begin{aligned} \operatorname{Re} f_{T,b}(\underline{k}) &= \frac{1}{T} \int_T^{2T} \cos(\varphi(t) A_b(\underline{k})) dt = \frac{1}{A_b(\underline{k}) T} \int_T^{2T} \frac{1}{\varphi'(t)} d \sin(\varphi(t) A_b(\underline{k})) \\ &\ll \frac{1}{|A_b(\underline{k})|} \frac{1}{\varphi'(2T) T} = o(1), \quad T \rightarrow \infty, \end{aligned}$$

provided that $A_b(\underline{k}) \neq 0$. Clearly, the same estimate holds for $\operatorname{Im} f_{T,b}(\underline{k})$. Hence, for $A_b(\underline{k}) \neq 0$,

$$\lim_{T \rightarrow \infty} f_{T,b}(\underline{k}) = 0. \tag{5}$$

Obviously,

$$f_{T,b}(\underline{k}) = 1$$

if $A_b(\underline{k}) = 0$. This and (5) show that

$$V_{T,b} \xrightarrow[T \rightarrow \infty]{w} V_b,$$

where V_b is a probability measure on $(\mathbb{I}_b, \mathcal{B}(\mathbb{I}_b))$ defined by the Fourier transform

$$f_b(k) = \begin{cases} 1 & \text{if } A_b(k) = 0, \\ 0 & \text{if } A_b(k) \neq 0. \end{cases}$$

□

Now, we will apply Lemma 1 for the measure defined by means of a certain finite sum. Let $\theta > 1/2$ be a fixed number, and, for $x, y \in [1, \infty)$,

$$u(x, y) = \exp\left\{-\left(\frac{x}{y}\right)^\theta\right\}.$$

Moreover, we use the notation $\widehat{\zeta}(t) = |\zeta(1/2 + it)|^2$. Consider the n th integral sum

$$U_{n,b,y}(\sigma + i\varphi(t)) = \frac{b-1}{n} \sum_{l=1}^n \widehat{\zeta}(a_l) u(a_l, y) a_l^{-\sigma - i\varphi(t)}, \quad n \in \mathbb{N},$$

where $a_l \in [x_{l-1}, x_l]$ and $x_l = 1 + ((b-1)/n)l$.

For $A \in \mathcal{B}(\mathbb{C})$, set

$$P_{T,n,b,y}(A) = \frac{1}{T} \mathbf{L}\left\{t \in [T, 2T] : U_{n,b,y}(\sigma + i\varphi(t)) \in A\right\}.$$

For simplicity, here and in the following, we omit the dependence on σ of some objects. Before the statement of the limit lemma for $P_{T,n,b,y}$, we will present some lower estimates for the mean square $I_\sigma(T)$. For this, we will apply the following general lemma from [8]. Let $\mathcal{F}(s)$ be the modified Mellin transform of $f(x)$, i.e.,

$$\mathcal{F}(s) = \int_1^\infty f(x)x^{-s} dx.$$

Lemma 2 ([8], Lemma 5). *Let $f(x) \in C^\infty[2, \infty]$ be a real-valued function such that*

1°

$$\int_1^X |f^{(k)}(x)| dx \ll_{\varepsilon,k} X^{1+\varepsilon}, \quad k \in \mathbb{N}_0;$$

2° $\mathcal{F}(s)$ has analytic continuation to the half-plane $\sigma > 1/2$, except for a pole of order l at the point $s = 1$;

3° For $\sigma > 1/2$, $\mathcal{F}(s)$ is of polynomial growth in $|t|$.

Then, for $1/2 < \sigma < 1$ and any fixed $\varepsilon > 0$,

$$\int_T^{2T} f^2(x) dx \ll_\varepsilon \log^{l-1} T \int_{T/2}^{5T/2} |f(x)| dx + T^{2\sigma-1} \int_0^{T^{1+\varepsilon}} |\mathcal{F}(\sigma + it)|^2 dt.$$

Lemma 3. *For $1/2 < \sigma < 1$, and any $\varepsilon > 0$, the estimate*

$$I_\sigma(T) \gg_\varepsilon T^{2-2\sigma-\varepsilon}$$

holds.

Proof. As usual, denote by $Z(t)$, $t \in \mathbb{R}$, the Hardy function, i.e.,

$$Z(t) = \zeta\left(\frac{1}{2} + it\right) \chi^{-1/2}\left(\frac{1}{2} + it\right),$$

where

$$\chi(s) = \frac{\zeta(s)}{\zeta(1-s)}.$$

It is well known that $Z(t)$ is a real-valued function satisfying $|Z(t)| = |\zeta(1/2 + it)|$. Moreover, the estimate [8]

$$Z^{(k)}(t) \ll_k t^{-1/4}(\log T)^{k+1} + \sum_{m \leq \sqrt{t/(2\pi)}} m^{-1/2} \left(\log \frac{\sqrt{t/(2\pi)}}{m} \right)^k \tag{6}$$

holds. Take $f(x) = Z^2(x)$. Then, we have

$$\mathcal{F}(s) = \int_1^\infty Z^2(x)x^{-s} dx = \int_1^\infty \left| \zeta\left(\frac{1}{2} + ix\right) \right|^2 x^{-s} dx = \mathcal{Z}(s).$$

In view of Theorem 1 and (6), the function satisfies the hypotheses of Lemma 1 with $l = 2$. Thus, for $1/2 < \sigma < 1$,

$$\int_T^{2T} f^2(t) dt = \int_T^{2T} \left| \zeta\left(\frac{1}{2} + it\right) \right|^4 dt \ll_\epsilon \log T \int_{T/2}^{5T/2} \left| \zeta\left(\frac{1}{2} + it\right) \right|^2 dt + T^{2\sigma-1} \int_0^{T^{1+\epsilon}} |\mathcal{Z}(\sigma + it)|^2 dt.$$

Since [5]

$$\int_0^T \left| \zeta\left(\frac{1}{2} + it\right) \right|^4 dt = \frac{1}{2\pi^2} T \log^4 T + O(T \log^3 T)$$

and

$$\int_0^T \left| \zeta\left(\frac{1}{2} + it\right) \right|^2 dt \ll T \log T,$$

this implies

$$T \log^4 T \ll_\epsilon T^{2\sigma-1} \int_0^{T^{1+\epsilon}} |\mathcal{Z}(\sigma + it)|^2 dt.$$

Consequently,

$$I_\sigma(T) \gg_\epsilon T^{(2-2\sigma)/(1+\epsilon)} \gg_\epsilon T^{2-2\sigma-\epsilon}.$$

□

Lemma 4. Assume that $\sigma \in (1/2, 1)$ is a given fixed number, and $\varphi(t) \in W_\sigma$. Then, on $(\mathbb{C}, \mathcal{B}(\mathbb{C}))$, there exists a probability measure $P_{n,b,y}$ such that $P_{T,n,b,y} \xrightarrow[T \rightarrow \infty]{w} P_{n,b,y}$.

Proof. Lemma 3 implies that, for $\sigma \in (1/2, 1)$, $I_\sigma(T) \rightarrow \infty$ as $T \rightarrow \infty$. Therefore, if $\varphi(t) \in W_\sigma$, then

$$\frac{1}{\varphi'(T)} \ll TI_\sigma^{-1}(\varphi(T)) = o(T)$$

as $T \rightarrow \infty$. Thus, the application of Lemma 1 is possible.

Consider the mapping $v_{n,b} : \mathbb{I}_b \rightarrow \mathbb{C}$ defined by

$$v_{n,b}(i) = \frac{b-1}{n} \sum_{l=1}^n \widehat{\zeta}(a_l) u(a_l, y) a_l^{-\sigma} i_{a_l}. \tag{7}$$

Since the latter sum is finite, and \mathbb{I}_b is equipped with the product topology, the mapping $v_{n,b}$ is continuous. Moreover, in view of (7),

$$v_{n,b}(x^{-i\varphi(t)} : x \in [1, b]) = \frac{b-1}{n} \sum_{l=1}^n \widehat{\zeta}(a_l) u(a_l, y) a_l^{-\sigma-i\varphi(t)} = U_{n,b,y}(\sigma + i\varphi(t)).$$

Hence, for $A \in \mathcal{B}(\mathbb{C})$,

$$\begin{aligned} P_{T,n,b,y}(A) &= \frac{1}{T} \mathbf{L} \left\{ t \in [T, 2T] : v_{n,b}(x^{-i\varphi(t)} : x \in [1, b]) \in A \right\} \\ &= \frac{1}{T} \mathbf{L} \left\{ t \in [T, 2T] : (x^{-i\varphi(t)} : x \in [1, b]) \in v_{n,b}^{-1}A \right\} = V_{T,b}(v_{n,b}^{-1}A), \end{aligned} \tag{8}$$

where $V_{T,b}$ is from Lemma 1. The continuity of the mapping $v_{n,b}$ implies its $(\mathcal{B}(\mathbb{I}_b), \mathcal{B}(\mathbb{C}))$ -measurability. Therefore, the mapping $v_{n,b}$ and any probability measure P on $(\mathbb{I}_b, \mathcal{B}(\mathbb{I}_b))$ define the unique probability measure $Pv_{n,b}^{-1}$ on $(\mathbb{C}, \mathcal{B}(\mathbb{C}))$ given by

$$Pv_{n,b}^{-1}(A) = P(v_{n,b}^{-1}A), \quad A \in \mathcal{B}(\mathbb{C}).$$

See Section 2 of [24]. Thus, by (8), we have $P_{T,n,b,y} = V_{T,b}v_{n,b}^{-1}$. Therefore, Lemma 1, the continuity of $v_{n,b}$, and the principle of the preservation of weak convergence under continuity mappings (Theorem 5.1 of [24]) show that

$$P_{T,n,b,y} \xrightarrow[T \rightarrow \infty]{w} P_{n,b,y},$$

where $P_{n,b,y} = V_b v_{n,b}^{-1}$, and V_b is the limit measure in Lemma 1. \square

3. Limit Lemma for Integral

Denote

$$\mathcal{Z}_{b,y}(\sigma + i\varphi(t)) = \int_1^b \widehat{\zeta}(x) u(x, y) x^{-\sigma-i\varphi(t)} dx,$$

and, for $A \in \mathcal{B}(\mathbb{C})$, set

$$P_{T,b,y}(A) = \frac{1}{T} \mathbf{L} \left\{ t \in [T, 2T] : \mathcal{Z}_{b,y}(\sigma + i\varphi(t)) \in A \right\}.$$

In this section, we will prove the weak convergence for $P_{T,b,y}$ as $T \rightarrow \infty$. Before this, we recall some known probabilistic results. Let $\{Q\}$ be a certain family of probability measures on $(\mathbb{X}, \mathcal{B}(\mathbb{X}))$. The family $\{Q\}$ is called tight if, for every $\delta > 0$, there is a compact set $K \subset \mathbb{X}$ such that

$$Q(K) > 1 - \delta$$

for all $Q \in \{Q\}$. The family $\{Q\}$ is said to be relatively compact if every sequence contains a subsequence weakly convergent to a certain probability measure on $(\mathbb{X}, \mathcal{B}(\mathbb{X}))$. The Prokhorov theorem connects two above notions, and, for convenience, we state it as the following lemma.

Lemma 5. *If a family of probability measures is tight, then it is relatively compact.*

The proof of the lemma is given in [24], Theorem 5.1.

Moreover, we recall one useful property on convergence in distribution. Let ζ_n and ζ be \mathbb{X} -valued random elements defined on the probability space $(\Omega, \mathcal{F}, \mu)$ with distributions P_n and P , respectively. Then, ζ_n converges in distribution to ζ as $n \rightarrow \infty$ $\left(\frac{\mathcal{D}}{n \rightarrow \infty}\right)$ if

$$P_n \xrightarrow[n \rightarrow \infty]{w} P.$$

Now, we state a lemma on convergence in distribution.

Lemma 6. Assume that the metric space (\mathbb{X}, d) is separable, and ζ_{nk}, ζ_n are \mathbb{X} -valued random elements defined on the same probability space $(\Omega, \mathcal{F}, \mu)$. Let

$$\zeta_{nk} \xrightarrow[n \rightarrow \infty]{\mathcal{D}} \zeta_k$$

and

$$\zeta_k \xrightarrow[k \rightarrow \infty]{\mathcal{D}} \zeta.$$

If, for every $\delta > 0$,

$$\lim_{k \rightarrow \infty} \limsup_{n \rightarrow \infty} \mu\{d(\zeta_{nk}, \eta_k) \geq \delta\} = 0,$$

then

$$\eta_n \xrightarrow[n \rightarrow \infty]{\mathcal{D}} \zeta.$$

The lemma is proven in [24], Theorem 3.2.

Lemma 7. Assume that $\sigma \in (1/2, 1)$ is a given fixed number, and $\varphi(t) \in W_\sigma$. Then, on $(\mathbb{C}, \mathcal{B}(\mathbb{C}))$, there exists a probability measure $P_{b,y}$ such that $P_{T,b,y} \xrightarrow[T \rightarrow \infty]{w} P_{b,y}$.

Proof. First, we will show that $\mathcal{Z}_{b,y}(\sigma + i\varphi(t))$ is close in a certain sense to $U_{n,b,y}(\sigma + i\varphi(t))$. Let

$$J_{T,n} \stackrel{\text{def}}{=} \frac{1}{T} \int_T^{2T} \left| \mathcal{Z}_{b,y}(\sigma + i\varphi(t)) - U_{n,b,y}(\sigma + \varphi(t)) \right| dt.$$

Clearly,

$$J_{T,n}^2 \leq \frac{1}{T} \int_T^{2T} \left| \mathcal{Z}_{b,y}(\sigma + i\varphi(t)) - U_{n,b,y}(\sigma + \varphi(t)) \right|^2 dt. \tag{9}$$

We have

$$\begin{aligned} & \int_T^{2T} \left| \mathcal{Z}_{b,y}(\sigma + i\varphi(t)) \right|^2 dt \\ &= \int_T^{2T} \left(\int_1^b \widehat{\zeta}(x) u(x,y) x^{-\sigma - i\varphi(t)} dx \right) \left(\int_1^b \widehat{\zeta}(x) u(x,y) x^{-\sigma + i\varphi(t)} dx \right) dt \\ &= T \int_{\substack{1 \\ x_1=x_2}}^b \int_{\substack{1 \\ x_2}}^b \widehat{\zeta}(x_1) \widehat{\zeta}(x_2) u(x_1,y) u(x_2,y) x_1^{-\sigma} x_2^{-\sigma} dx_1 dx_2 \\ & \quad + \int_{\substack{1 \\ x_1 \neq x_2}}^b \int_{\substack{1 \\ x_2}}^b \widehat{\zeta}(x_1) \widehat{\zeta}(x_2) u(x_1,y) u(x_2,y) x_1^{-\sigma} x_2^{-\sigma} \left(\int_T^{2T} \left(\frac{x_1}{x_2} \right)^{i\varphi(t)} dt \right) dx_1 dx_2. \end{aligned} \tag{10}$$

Since

$$\operatorname{Re} \int_T^{2T} \left(\frac{x_1}{x_2}\right)^{i\varphi(t)} dt = \left(\log\left(\frac{x_1}{x_2}\right)\right)^{-1} \int_T^{2T} \frac{1}{\varphi'(t)} d \sin\left(\varphi(t) \log\left(\frac{x_1}{x_2}\right)\right) \ll \left|\log \frac{x_1}{x_2}\right|^{-1} \frac{1}{\varphi'(2T)},$$

and the same bound is true for the imaginary part of the latter integral, we obtain by (10) that

$$\int_T^{2T} \left| \mathcal{Z}_{b,y}(\sigma + i\varphi(t)) \right|^2 dt = o(T), \quad T \rightarrow \infty. \tag{11}$$

Reasoning similarly, we find

$$\begin{aligned} \int_T^{2T} \left| U_{n,b,y}(\sigma + i\varphi(t)) \right|^2 dt &= T \left(\frac{b-1}{n}\right)^2 \sum_{l=1}^n \widehat{\zeta}^2(a_l) u^2(a_l, y) a_l^{-2\sigma} \\ &+ O\left(\left(\frac{b-1}{n}\right)^2 \sum_{l_1=1}^n \sum_{\substack{l_2=1 \\ l_1 \neq l_2}}^n \widehat{\zeta}(a_{l_1}) \widehat{\zeta}(a_{l_2}) u(a_{l_1}, y) u(a_{l_2}, y) a_{l_1}^{-\sigma} a_{l_2}^{-\sigma} \left|\log \frac{a_{l_1}}{a_{l_2}}\right|^{-1}\right). \end{aligned} \tag{12}$$

Thus,

$$\lim_{n \rightarrow \infty} \limsup_{T \rightarrow \infty} \frac{1}{T} \int_T^{2T} \left| U_{n,b,y}(\sigma + i\varphi(t)) \right|^2 dt = 0. \tag{13}$$

By (9),

$$\begin{aligned} J_{T,n}^2 &\ll \frac{1}{T} \left(\int_T^{2T} \left| \mathcal{Z}_{b,y}(\sigma + i\varphi(t)) \right|^2 dt + \left(\int_T^{2T} \left| \mathcal{Z}_{b,y}(\sigma + i\varphi(t)) \right|^2 dt \int_T^{2T} \left| U_{n,b,y}(\sigma + i\varphi(t)) \right|^2 dt \right)^{1/2} \right. \\ &\left. + \int_T^{2T} \left| U_{n,b,y}(\sigma + i\varphi(t)) \right|^2 dt \right). \end{aligned}$$

Therefore, (11) and (13) yield

$$\lim_{n \rightarrow \infty} \limsup_{T \rightarrow \infty} J_{T,n} = 0. \tag{14}$$

Now, we will deal with the sequence $\{P_{n,b,y} : n \in \mathbb{N}\}$. By (12), we have

$$\begin{aligned} &\sup_{s \in \mathbb{N}} \limsup_{T \rightarrow \infty} \frac{1}{T} \int_T^{2T} \left| U_{n,b,y}(\sigma + i\varphi(t)) \right| dt \\ &\ll \sup_{s \in \mathbb{N}} \limsup_{T \rightarrow \infty} \left(\frac{1}{T} \int_T^{2T} \left| U_{n,b,y}(\sigma + i\varphi(t)) \right|^2 dt \right)^{1/2} \\ &\ll \sup_{n \in \mathbb{N}} \frac{b-1}{n} \left(\sum_{l=1}^n \widehat{\zeta}^2(a_l) u^2(a_l, y) a_l^{-2\sigma} \right)^{1/2} \leq C_{b,y,\sigma} < \infty \end{aligned} \tag{15}$$

because

$$\lim_{n \rightarrow \infty} \frac{b-1}{n} \sum_{l=1}^n \widehat{\zeta}^2(a_l) u^2(a_l, y) a_l^{-2\sigma} = \int_1^b \widehat{\zeta}^2(x) u^2(x, y) x^{-2\sigma} dx.$$

Take a random variable θ_T given on the probability space $(\Omega, \mathcal{F}, \mu)$ that is uniformly distributed on $[T, 2T]$. Consider the complex-valued random variables

$$x_{T,n,b,y} = x_{T,n,b,y}(\sigma) = U_{n,b,y}(\sigma + i\varphi(\theta_T)),$$

and $x_{n,b,y}(\sigma)$ with the distribution $P_{n,b,y,\sigma}$. Then, rewrite the assertion of Lemma 4 in the form

$$x_{T,n,b,y} \xrightarrow[T \rightarrow \infty]{\mathcal{D}} x_{n,b,y}. \tag{16}$$

Fix $\delta > 0$. Then, in view of (15) and (16),

$$\begin{aligned} \mu \left\{ \left| x_{n,b,y}(\sigma) \right| > \delta^{-1} C_{b,y,\sigma} \right\} &\leq \sup_{n \in \mathbb{N}} \limsup_{T \rightarrow \infty} \mu \left\{ \left| x_{T,n,b,y}(\sigma) \right| > \delta^{-1} C_{b,y,\sigma} \right\} \\ &\leq \sup_{n \in \mathbb{N}} \limsup_{T \rightarrow \infty} \frac{\delta}{C_{b,y,\sigma}} \int_T^{2T} \left| U_{n,b,y}(\sigma + i\varphi(t)) \right| dt \leq \delta. \end{aligned} \tag{17}$$

The set $K = \{s \in \mathbb{C} : |s| \leq \delta^{-1} C_{b,y,\sigma}\}$ is compact in \mathbb{C} . Moreover, by (17),

$$\mu \left\{ x_{n,b,y} \in K \right\} = 1 - \mu \left\{ x_{n,b,y} \notin K \right\} > 1 - \delta$$

for all $n \in \mathbb{N}$. This and the definition of $x_{n,b,y}$ show that, for all $n \in \mathbb{N}$,

$$P_{n,b,y,\sigma}(K) > 1 - \delta.$$

This means that the sequence $\{P_{n,b,y,\sigma} : n \in \mathbb{N}\}$ is tight. Therefore, by Lemma 5, it is relatively compact. Hence, there exists a subsequence $\{P_{n_l,b,y,\sigma}\} \subset \{P_{n,b,y,\sigma}\}$ and a probability measure $P_{b,y,\sigma}$ on $(\mathbb{C}, \mathcal{B}(\mathbb{C}))$ such that $P_{n_l,b,y,\sigma} \xrightarrow[l \rightarrow \infty]{w} P_{b,y,\sigma}$. In other words,

$$x_{n_l,b,y} \xrightarrow[l \rightarrow \infty]{\mathcal{D}} P_{b,y,\sigma}.$$

This, (16), and (14) show that all hypotheses of Lemma 6 for $x_{T,n,b,y}$, $x_{n_l,b,y}$ and

$$y_{T,b,y} = y_{T,b,y}(\sigma) = \mathcal{Z}_{b,y}(\sigma + i\varphi(\theta_T))$$

are satisfied. Thus, we have

$$y_{T,b,y} \xrightarrow[T \rightarrow \infty]{\mathcal{D}} P_{b,y,\sigma},$$

which proves the lemma. \square

4. Case of Improper Integral

This section is devoted to a limit lemma for the integral

$$\mathcal{Z}_y(\sigma + i\varphi(t)) = \int_1^\infty \widehat{\zeta}(x) u(x, y) x^{-\sigma - i\varphi(t)} dx.$$

It is well known that $\zeta(1/2 + ix) \ll (1 + |x|)^{1/6}$. Therefore, the integral for $\mathcal{Z}(\sigma + i\varphi(t))$ converges absolutely for $\sigma > \widehat{\sigma}$ with every finite $\widehat{\sigma}$.

For $A \in \mathcal{B}(\mathbb{C})$, let

$$P_{T,y,\sigma}(A) = \frac{1}{T} \mathbf{L} \{t \in [T, 2T] : \mathcal{Z}_y(\sigma + i\varphi(t)) \in A\}.$$

Lemma 8. Assume that $\sigma \in (1/2, 1)$ is a given fixed number, and $\varphi(t) \in W_\sigma$. Then, there is a probability measure $P_{y,\sigma}$ on $(\mathbb{C}, \mathcal{B}(\mathbb{C}))$ such that $P_{T,y,\sigma} \xrightarrow[T \rightarrow \infty]{w} P_{y,\sigma}$.

Proof. We use a similar method as in the proof of Lemma 7. We begin with a mean value

$$J_{T,y} \stackrel{\text{def}}{=} \frac{1}{T} \int_0^T \left| \mathcal{Z}_y(\sigma + i\varphi(t)) - \mathcal{Z}_{b,y}(\sigma + i\varphi(t)) \right| dt.$$

Clearly, the absolute convergence of the integral for $\mathcal{Z}_y(\sigma + i\varphi(t))$ shows that, for every fixed $y > 0$,

$$\begin{aligned} \mathcal{Z}_y(\sigma + i\varphi(t)) - \mathcal{Z}_{b,y}(\sigma + i\varphi(t)) &= \int_b^\infty \widehat{\zeta}(x) u(x, y) x^{-\sigma - i\varphi(t)} dx \\ &\ll \int_b^\infty \widehat{\zeta}(x) u(x, y) x^{-\sigma} dx = o_y(1) \end{aligned}$$

as $b \rightarrow \infty$. Hence, we obtain

$$\lim_{b \rightarrow \infty} \limsup_{T \rightarrow \infty} J_{T,y} = 0. \tag{18}$$

Let $y_{b,y}(\sigma)$ be the complex-valued random variable with distribution $P_{b,y,\sigma}$, and, in the notation of Lemma 7,

$$y_{T,b,y} = y_{T,b,y}(\sigma) = \mathcal{Z}_{b,y}(\sigma + i\varphi(\theta_T)).$$

Then, by Lemma 7,

$$y_{T,b,y} \xrightarrow[T \rightarrow \infty]{D} y_{b,y}. \tag{19}$$

Moreover, in virtue of (11),

$$\sup_{b \geq 1} \limsup_{T \rightarrow \infty} \frac{1}{T} \int_0^{2T} \left| \mathcal{Z}_{b,y}(\sigma + i\varphi(t)) \right| dt \leq C_{y,\sigma} < \infty.$$

This together with (19) gives, for $\delta > 0$,

$$\begin{aligned} \mu \left\{ \left| y_{b,y} \right| > \delta^{-1} C_{y,\sigma} \right\} &\leq \sup_{b \geq 1} \limsup_{T \rightarrow \infty} \mu \left\{ \left| y_{b,y} \right| > \delta^{-1} C_{y,\sigma} \right\} \\ &\leq \sup_{b \geq 1} \limsup_{T \rightarrow \infty} \frac{\delta}{C_{y,\sigma}} \int_0^{2T} \left| \mathcal{Z}_{b,y}(\sigma + i\varphi(t)) \right| dt \leq \delta. \end{aligned}$$

Taking a set $K = \{s \in \mathbb{C} : |s| \leq \delta^{-1} C_{y,\sigma}\}$, from this, we deduce that

$$\mu \left\{ y_{b,y} \in K \right\} > 1 - \delta.$$

This implies that the family $\{P_{b,y,\sigma} : b \geq 1\}$ is tight. Therefore, in view of Lemma 5, it is relatively compact. Thus, there is a sequence $\{P_{b_l,y,\sigma}\}$ and a probability measure $P_{y,\sigma}$ on $(\mathbb{C}, \mathcal{B}(\mathbb{C}))$ such that

$$y_{b_l,y,\sigma} \xrightarrow[l \rightarrow \infty]{D} P_{y,\sigma}.$$

This, (19), (18), and the application of Lemma 6 complete the proof of the lemma. \square

5. Proof of Theorem 2

We recall that

$$u(x, y) = \exp\left\{-\left(\frac{x}{y}\right)^\theta\right\}, \quad x, y \in [1, \infty),$$

with a fixed $\theta > 1/2$. For brevity, set

$$f(s, y) = \frac{1}{\theta} \Gamma\left(\frac{s}{\theta}\right) y^s,$$

where $\Gamma(s)$ is the Euler gamma-function. For the approximation of $\mathcal{Z}(\sigma + i\varphi(t))$ by $\mathcal{Z}_y(\sigma + \varphi(t))$, we use the representation

$$\mathcal{Z}_y(s) = \frac{1}{2\pi i} \int_{\theta - i\infty}^{\theta + i\infty} \mathcal{Z}(s + z) f(z, y) dz, \quad \frac{1}{2} < \sigma < 1, \tag{20}$$

obtained in [25], Lemma 9.

Lemma 9. *Under the hypotheses of Theorem 2,*

$$\lim_{y \rightarrow \infty} \limsup_{T \rightarrow \infty} \frac{1}{T} \int_T^{2T} |\mathcal{Z}(\sigma + i\varphi(t)) - \mathcal{Z}_y(\sigma + i\varphi(t))| dt = 0.$$

Proof. Let $\theta_1 = -\varepsilon$ and $\theta = 1/2 + \varepsilon$. The integrand in (20) has a double pole $z = 1 - s$ and a simple pole $z = 0$ lying in $\theta_1 < \text{Re} z < \theta$. Therefore, by the residue theorem and (20), we have

$$\mathcal{Z}_y(s) - \mathcal{Z}(s) = \frac{1}{2\pi i} \int_{\theta_1 - i\infty}^{\theta_1 + i\infty} \mathcal{Z}(s + z) f(z, y) dz + r_y(s),$$

where

$$r_y(s) = \text{Res}_{z=1-s} \mathcal{Z}(s) f(s, y). \tag{21}$$

From this, we obtain

$$\begin{aligned} & \mathcal{Z}_y(\sigma + i\varphi(t)) - \mathcal{Z}(\sigma + i\varphi(t)) \\ &= \frac{1}{2\pi} \int_{-\infty}^{\infty} \mathcal{Z}(\sigma - \varepsilon + i\varphi(t) + i\tau) f(-\varepsilon + i\tau, y) d\tau + r_y(\sigma + i\varphi(t)) \\ &\ll \int_{-\infty}^{\infty} |\mathcal{Z}(\sigma - \varepsilon + i\varphi(t) + i\tau)| |f(-\varepsilon + i\tau, y)| d\tau + |r_y(\sigma + i\varphi(t))|. \end{aligned}$$

Thus,

$$\frac{1}{T} \int_T^{2T} |\mathcal{Z}(\sigma + i\varphi(t)) - \mathcal{Z}_y(\sigma + i\varphi(t))| dt \ll I_{T,y},$$

where

$$\begin{aligned}
 I_{T,y} &\stackrel{\text{def}}{=} \int_{-\infty}^{\infty} \left(\frac{1}{T} \int_T^{2T} |\mathcal{Z}(\sigma - \varepsilon + i\varphi(t) + i\tau)| dt \right) |f(-\varepsilon + i\tau, y)| d\tau \\
 &\quad + \frac{1}{T} \int_T^{2T} |r_y(\sigma + i\varphi(t))| dt = I_{T,y}^{(1)} + I_{T,y}^{(2)}.
 \end{aligned}
 \tag{22}$$

To estimate $I_{T,y}^{(1)}$, we observe that

$$\begin{aligned}
 \frac{1}{T} \int_T^{2T} |\mathcal{Z}(\sigma - \varepsilon + i\varphi(t) + i\tau)| dt &\leq \left(\frac{1}{T} \int_T^{2T} |\mathcal{Z}(\sigma - \varepsilon + i\varphi(t) + i\tau)|^2 dt \right)^{1/2} \\
 &= \left(\frac{1}{T} \int_T^{2T} |\mathcal{Z}(\sigma - \varepsilon + i\varphi(t) + i\tau)|^2 \frac{\varphi'(t) dt}{\varphi'(t)} \right)^{1/2} \\
 &\ll \left(\frac{1}{T\varphi'(2T + |\tau|)} \int_0^{\varphi(2T + |\tau|)} |\mathcal{Z}(\sigma - \varepsilon + iu)|^2 du \right) \\
 &\ll \left(\frac{I_{\sigma - \varepsilon} \varphi(2T + |\tau|)}{T\varphi'(2T + |\tau|)} \right)^{1/2} \\
 &\ll \left(\frac{2T + |\tau|}{T} \right)^{1/2} \ll (1 + |\tau|)^{1/2}.
 \end{aligned}
 \tag{23}$$

For the gamma-function, the estimate

$$\Gamma(\sigma + it) \ll \exp\{-c|t|\}, \quad c > 0,
 \tag{24}$$

is valid. Therefore,

$$f(-\varepsilon + i\tau, y) \ll y^{-\varepsilon} \exp\{-c_1|\tau|\}, \quad c_1 > 0.$$

This together with (23) leads to the bound

$$I_{T,y}^{(1)} \ll y^{-\varepsilon} \int_{-\infty}^{\infty} (1 + |\tau|)^{1/2} \exp\{-c_1|\tau|\} d\tau \ll y^{-\varepsilon}.
 \tag{25}$$

Let $a = 2\gamma - \log 2\pi$. In view of the formula for $\mathcal{Z}(s)$ in Theorem 1,

$$\begin{aligned}
 r_y(s) &= f'(1 - s, y) + af(1 - s, y) \\
 &= \frac{1}{\theta^2} \Gamma' \left(\frac{1 - s}{\theta} \right) y^{1-s} + \frac{1}{\theta} \Gamma \left(\frac{1 - s}{\theta} \right) y^{1-s} \log y + \frac{a}{\theta} \Gamma \left(\frac{1 - s}{\theta} \right) y^{1-s} \\
 &= \frac{y^{1-s}}{\theta} \Gamma \left(\frac{1 - s}{\theta} \right) \left(\frac{1}{\theta} \frac{\Gamma'}{\Gamma} \left(\frac{1 - s}{\theta} \right) + \log y + a \right).
 \end{aligned}$$

Hence, the estimates (24) and

$$\frac{\Gamma'}{\Gamma}(\sigma + it) \ll \log(|t| + 2)$$

yield

$$I_{T,y}^{(2)} \ll_{\theta} y^{1-\sigma} \log y \frac{1}{T} \int_T^{2T} \exp\left\{-\frac{c}{\theta} \varphi(t)\right\} \log \varphi(t) dt$$

$$\ll_{\theta} y^{1-\sigma} \log y \exp\left\{-\frac{c}{2\theta} \varphi(T)\right\}.$$

This, (25), and (22) show that

$$I_{T,y} \ll_{\delta} y^{-\varepsilon} + y^{1-\sigma} \log y \exp\left\{-\frac{c}{2\theta} \varphi(T)\right\}.$$

Therefore,

$$\lim_{y \rightarrow \infty} \limsup_{T \rightarrow \infty} \frac{1}{T} \int_T^{2T} |\mathcal{Z}(\sigma + i\varphi(t)) - \mathcal{Z}_y(\sigma + i\varphi(t))| dt = 0 \tag{26}$$

because $\varphi(T) \rightarrow \infty$ as $T \rightarrow \infty$. \square

Now, we return to the limit measure $P_{y,\sigma}$ of Lemma 8.

Lemma 10. *Under the hypotheses of Theorem 2, the family $\{P_{y,\sigma} : y \in [1, \infty)\}$ is tight.*

Proof. We have

$$\frac{1}{T} \int_T^{2T} |\mathcal{Z}_y(\sigma + i\varphi(t))| dt \leq \frac{1}{T} \int_T^{2T} |\mathcal{Z}(\sigma + i\varphi(t)) - \mathcal{Z}_y(\sigma + i\varphi(t))| dt + \frac{1}{T} \int_T^{2T} |\mathcal{Z}(\sigma + i\varphi(t))| dt.$$

Therefore, by (2) and (26),

$$\sup_{y \geq 1} \limsup_{T \rightarrow \infty} \frac{1}{T} \int_T^{2T} |\mathcal{Z}_y(\sigma + i\varphi(t))| dt \leq C < \infty. \tag{27}$$

Let

$$z_{T,y} = z_{T,y}(\sigma) = \mathcal{Z}_y(\sigma + i\varphi(\theta_T)),$$

and $z_y = z_y(\sigma)$ be the complex-valued random variable with the distribution $P_{y,\sigma}$. Then, the statement of Lemma 8 can be written as

$$z_{T,y} \xrightarrow[T \rightarrow \infty]{\mathcal{D}} z_y. \tag{28}$$

From this and (27), we obtain that, for every $\delta > 0$,

$$\mu\{|z_y| > \delta^{-1}C\} \leq \sup_{y \geq 1} \limsup_{T \rightarrow \infty} \mu\{|z_{T,y}| > \delta^{-1}C\} \leq \frac{\delta}{TC} \int_T^{2T} |\mathcal{Z}_y(\sigma + i\varphi(t))| dt \leq \delta.$$

This shows that, for $K = \{s \in \mathbb{C} : |s| \leq \delta^{-1}C\}$,

$$P_{y,\sigma}(K) \geq 1 - \delta,$$

and the lemma is proven. \square

Proof of Theorem 2. Lemma 10 together with Lemma 5 implies that the family $\{P_{y,\sigma}\}$ is relatively compact. Therefore, there is a sequence $\{P_{y_k,\sigma}\} \subset \{P_{y,\sigma}\}$ weakly convergent to a certain probability measure P_σ on $(\mathbb{C}, \mathcal{B}(\mathbb{C}))$ as $k \rightarrow \infty$. This also can be written as

$$z_{y_k,\sigma} \xrightarrow[k \rightarrow \infty]{\mathcal{D}} P_\sigma. \quad (29)$$

Define one more random variable,

$$z_T = z_T(\sigma) = \mathcal{Z}(\sigma + i\varphi(\theta_T)).$$

Then, Lemma 9 implies, for every $\delta > 0$,

$$\begin{aligned} & \lim_{k \rightarrow \infty} \limsup_{T \rightarrow \infty} \mu\{|z_T - z_{T,y_k}| > \delta\} \\ & \leq \lim_{k \rightarrow \infty} \limsup_{T \rightarrow \infty} \frac{1}{\delta T} \int_T^{2T} |\mathcal{Z}(\sigma + i\varphi(t)) - \mathcal{Z}_{y_k}(\sigma + i\varphi(t))| dt = 0. \end{aligned}$$

This, (28), and (29) together with Lemma 6 prove that

$$z_T \xrightarrow[T \rightarrow \infty]{\mathcal{D}} P_\sigma.$$

The theorem is proven. \square

6. Conclusions

In the paper, we considered the asymptotic behavior of the modified Mellin transform of the square of the Riemann zeta-function by using a probabilistic approach. We proved a limit theorem on the weak convergence of probability measures defined by shifts $\mathcal{Z}(\sigma + i\varphi(t))$, $1/2 < \sigma < 1$, where $\varphi(t)$ is a differentiable increasing to infinity function with a monotonically decreasing derivative $\varphi'(t)$ satisfying a certain estimate connected to the mean square of the function $\mathcal{Z}(s)$. We expect that such normalization of the function $\mathcal{Z}(s)$ extends the class of limit distributions for $\mathcal{Z}(s)$. Our future plans are related to a similar theorem in the space of analytic functions.

Author Contributions: Conceptualization, A.L. and D.Š.; methodology, A.L. and D.Š.; investigation, A.L. and D.Š.; writing—original draft preparation, A.L. and D.Š. All authors have read and agreed to the published version of the manuscript.

Funding: This research received no external funding.

Data Availability Statement: Data are contained within the article.

Conflicts of Interest: The authors declare no conflicts of interest.

References

1. Riemann, B. Über die Anzahl der Primzahlen unterhalb einer gegebenen Grösse. *Monatsber. Preuss. Akad. Wiss. Berlin* **1859**, 671–680.
2. Hadamard, J. Sur les zéros de la fonction $\zeta(s)$ de Riemann. *Comptes Rendus Acad. Sci. Paris* **1896**, 122, 1470–1473.
3. de la Vallée-Poussin, C.J. Recherches analytiques sur la théorie des nombres premiers, I–III. *Ann. Soc. Sci. Brux.* **1896**, 20, 183–256, 281–362, 363–397.
4. The Millennium Prize Problems. Available online: <https://www.claymath.org/millennium-problems/> (accessed on 1 March 2024).
5. Ivič, A. *The Riemann Zeta-Function*; John Wiley & Sons: New York, NY, USA, 1985.
6. Motohashi, Y. A relation between the Riemann zeta-function and the hyperbolic Laplacian. *Ann. Sc. Norm. Super. Pisa Cl. Sci. IV. Ser.* **1995**, 22, 299–313.
7. Motohashi, Y. *Spectral Theory of the Riemann Zeta-Function*; Cambridge University Press: Cambridge, UK, 1997.
8. Ivič, A. On some conjectures and results for the Riemann zeta-function and Hecke series. *Acta Arith.* **2001**, 99, 115–145. [[CrossRef](#)]
9. Ivič, A.; Jutila, M.; Motohashi, Y. The Mellin transform of powers of the zeta-function. *Acta Arith.* **2000**, 95, 305–342. [[CrossRef](#)]

10. Bohr, H. Über das Verhalten von $\zeta(s)$ in der Halbebene $\sigma > 1$. *Nachr. Akad. Wiss. Göttingen II Math. Phys. Kl.* **1911**, 409–428.
11. Bohr, H.; Jessen, B. Über die Wertverteilung der Riemanschen Zetafunktion, erste Mitteilung. *Acta Math.* **1930**, *54*, 1–35. [[CrossRef](#)]
12. Bohr, H.; Jessen, B. Über die Wertverteilung der Riemanschen Zetafunktion, zweite Mitteilung. *Acta Math.* **1932**, *58*, 1–55. [[CrossRef](#)]
13. Laurinćikas, A. Limit theorems for the Mellin transform of the square of the Riemann zeta-function. I. *Acta Arith.* **2006**, *122*, 173–184. [[CrossRef](#)]
14. Selberg, A. Old and New Conjectures and Results about a Class of Dirichlet Series. In Proceedings of the Amalfi Conference on Analytic Number Theory, Maiori, Italy, 25–29 September 1989; University of Salerno: Salerno, Italy, 1992; pp. 367–385
15. Joyner, D. *Distribution Theorems of L-Functions*; Ditman Research Notes in Math; Longman Scientific: Harlow, UK, 1986.
16. Bagchi, B. The Statistical Behaviour and Universality Properties of the Riemann Zeta-Function and Other Allied Dirichlet Series. Ph.D. Thesis, Indian Statistical Institute, Calcutta, India, 1981.
17. Korolev, M.A. Gram’s law in the theory of the Riemann zeta-function. Part 1. *Proc. Steklov Inst. Math.* **2016**, *292*, 1–146. [[CrossRef](#)]
18. Korolev, M.A. Gram’s law in the theory of the Riemann zeta-function. Part 2. *Proc. Steklov Inst. Math.* **2016**, *294*, 1–78. [[CrossRef](#)]
19. Kowalski, E. *An Introduction to Probabilistic Number Theory*; Cambridge Studies in Advanced Mathematics (192); Cambridge University Press: Cambridge, UK, 2021.
20. Lamzouri, Y.; Lester, S.; Radziwill, M. An effective universality theorem for the Riemann zeta function. *Comment. Math. Helv.* **2018**, *93*, 709–736. [[CrossRef](#)]
21. Lamzouri, Y.; Lester, S.; Radziwill, M. Discrepancy bounds for the distribution of the Riemann zeta-function and applications. *J. Anal. Math.* **2019**, *139*, 453–494. [[CrossRef](#)]
22. Steuding, J. *Value-Distribution of L-Functions*; Lecture Notes Math; Springer: Berlin/Heidelberg, Germany; New York, NY, USA, 2007; Volume 1877.
23. Matsumoto, K. Probabilistic value-distribution theory of zeta-functions. *Sugaku Expo.* **2004**, *17*, 51–71.
24. Billingsley, P. *Convergence of Probability Measures*, 2nd ed.; John Wiley & Sons: New York, NY, USA, 1999.
25. Korolev, M.; Laurinćikas, A. On the approximation by Mellin transform of the Riemann zeta-function. *Axioms* **2023**, *12*, 520. [[CrossRef](#)]

Disclaimer/Publisher’s Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.