



# DAMSS

DATA ANALYSIS  
METHODS FOR SOFTWARE  
SYSTEMS



**15th Conference on**

# **DATA ANALYSIS METHODS for Software Systems**

**November 28–30, 2024**

Druskininkai, Lithuania, Hotel “Europa Royale”

<https://www.mii.lt/DAMSS>

**Co-Chairmen:**

Prof. **Gintautas Dzemyda** (Vilnius University, Lithuanian Academy of Sciences)

Dr. **Saulius Maskeliūnas** (Lithuanian Computer Society)

**Programme Committee:**

Dr. **Jolita Bernatavičienė** (Lithuania)

Prof. **Juris Borzovs** (Latvia)

Prof. **Janis Grundspenkis** (Latvia)

Prof. **Janusz Kacprzyk** (Poland)

Prof. **Ignacy Kaliszewski** (Poland)

Prof. **Bożena Kostek** (Poland)

Prof. **Tomas Krilavičius** (Lithuania)

Prof. **Olga Kurasova** (Lithuania)

Assoc. Prof. **Tatiana Tchemisova** (Portugal)

Assoc. Prof. **Gintautas Tamulevičius** (Lithuania)

Prof. **Julius Žilinskas** (Lithuania)

**Organizing Committee:**

Dr. **Jolita Bernatavičienė**

Prof. **Olga Kurasova**

Assoc. Prof. **Viktor Medvedev**

**Laima Paliulionienė**

Assoc. Prof. **Martynas Sabaliauskas**

Prof. **Povilas Treigys**

**Contacts:**

Dr. Jolita Bernatavičienė

*jolita.bernatavicienne@mif.vu.lt*

Tel. (+370 5) 2109 315

Prof. Olga Kurasova

*olga.kurasova@mif.vu.lt*

Copyright © 2024 Authors. Published by Vilnius University Press.

This is an Open Access article distributed under the terms of the Creative Commons Attribution Licence, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

<https://doi.org/10.15388/DAMSS.15.2024>

ISBN 978-609-07-1112-5 (digital PDF)

© Vilnius University, 2024

# Spectral and Molecular Dynamics Studies of Carotenoids and Stilbene in Complex Systems for Supercomputing

Mindaugas Mačernis, Goda Bankovskaitė,  
Jonas Franukevičius, Darius Abramavicius

Institute of Chemical Physics, Faculty of Physics  
Vilnius University

*mindaugas.macernis@ff.vu.lt*

Molecular dynamics (MD) can utilize thousands of cores for research studies [1]. However, the Newtonian laws lack quantum physics phenomena, which are essential for understanding molecular properties in solvents, crystals, and amorphous materials [1-4]. Quantum MD methods, such as Car-Parrinello and Atom-Centered Density Matrix Propagation (ADMP) models or AIMD, can be updated to include excited-state dynamics, but understanding remains limited regarding physical properties and computational benchmarks [2-4]. In this study, we present results and discussions on the challenges of Quantum MD for selected carotenoids (Cars), stilbene, and other specific molecules in complex systems.

According to MD and DFT Raman analysis of lycopene and  $\beta$ -cyclodextrin interactions, we demonstrate that the Raman  $\nu_1$  mode shifts to lower or higher frequencies depending on the position of lycopene within the  $\beta$ -cyclodextrin. While MD provides insights into possible structural properties, it remains sensitive to the chosen environment, making it challenging to accurately model the complex dynamics of the lycopene and  $\beta$ -cyclodextrin system. Separate classical and quantum analyses are required.

Stilbene molecules are relatively small compared to carotenoids, but new methods are needed to interpret experimental data [4]. The combination of various Quantum MD approaches provides a deeper understanding, though it makes the required computational resources less predictable.

Moreover, phosphorus-phosphorus through-space indirect spin-spin J-couplings in different conformations, as derived from QMD, present challenges for conformer identification (Fig. 1). Using our proposed Molecular Dynamics Conformation Probability analysis, we identified at least five distinct conformers in CH(PPh)<sub>2</sub>.




Fig. 1. The P-P distance from Quantum MD is crucial for understanding NMR-based quantum computing

In conclusion, we discuss the limitations of MD and QMD methods in larger scale supercomputers.

**Acknowledgements:** Theoretical analysis and calculations were supported by the Research Council of Lithuania (Grant No. S-MIP-23-48). Computations were performed using high-performance supercomputer “VU HPC” Saulėtekis of Vilnius University at Faculty of Physics.

## References

- [1] M. Macernis, J. Franukevicius et al. arXiv preprint arXiv:2210.00934 (2022)
- [2] M. Macernis, A. Bockuviene et al. , J. Mol. Stuct. 1226, 129362 (2021)
- [3] M. Macernis, S. Streckaite et al. J. Phys. Chem. A126, 6 , 813-824 (2022)
- [4] Halimski, I, M. Macernis, D. Abramavicius et al. PCCP 26 (36), 23692-23702 (2024)