






MAIN

Longitudinal interplay between subjective stress, anxiety, depression, and well-being in internet-based stress recovery intervention for nurses

Inga Truskauskaitė¹ , Austėja Dumarkaitė¹ , Auguste Nomeikaite¹ , Gerhard Andersson^{2,3}  and Evaldas Kazlauskas¹ 

¹Institute of Psychology, Vilnius University, Vilnius, Lithuania, ²Department of Behavioural Sciences and Learning, Linköping University, Linköping, Sweden and ³Department of Clinical Neuroscience, Karolinska Institute, Stockholm, Sweden

Corresponding author: Inga Truskauskaitė; Email: inga.truskauskaitė@fsf.vu.lt

(Received 11 January 2024; revised 20 September 2024; accepted 9 October 2024)

Abstract

Background: Cognitive behavioural therapy (CBT) interventions are effective in reducing subjective stress. Nevertheless, the longitudinal links between mental health indicators are rarely studied in intervention research. Therefore, it is unknown how the intervention effects are sustained.

Aim: The current study investigated mechanisms explaining sustained intervention effects in a sample of medical nurses who receive a CBT-based internet-delivered stress recovery program.

Method: A single-group longitudinal study design with three measurement points, pre-test, post-test, and 3-month follow-up, was used in the current study. The sample consisted of nurses and assistant nurses from Lithuania ($n = 111$, age: $M (SD) = 41.69$ years (10.85)) who had participated in a 6-week CBT internet intervention targeting stress recovery. Data were collected as the randomised control trial, the treatment samples were combined, and the data were analysed using cross-lagged panel analysis with four variables representing the psychological well-being and symptoms of stress, anxiety, and depression.

Results: The results revealed that decreased anxiety and increased psychological well-being at post-test predicted reduced stress levels at the 3-month follow-up. In addition, decreased anxiety at post-test predicted decreased depression at follow-up.

Conclusions: Decreased anxiety and increased well-being could explain the sustainability of reduced stress following a CBT-based internet intervention for nurses. The implications of this for research and practice are discussed.

Keywords: anxiety; cross-lagged effect modelling; depression; internet-based CBT; stress

Introduction

Prolonged elevated stress often has devastating negative long-term effects on physical and mental health (Cohen, 2000; Haight *et al.*, 2023; O'Connor *et al.*, 2021). Moreover, stress imposes a substantial financial burden on society, mainly due to loss of productivity and increased demand for healthcare services (Hassard *et al.*, 2018). For some occupations, such as healthcare professionals, acute stress seems to be an inherent part of daily work (Mingote Adán *et al.*, 2004). Substantial evidence shows that stress-management interventions can effectively reduce subjective stress, with multi-modal, relaxation-based, and cognitive interventions showing the most beneficial effects in reducing stress symptoms (Estevez Cores *et al.*, 2021).

The increasing demand for psychological services (e.g. American Psychological Association, 2021) and the rapid development of secure information technologies have encouraged the growth of internet-delivered psychological interventions (Andersson, 2018). In the face of the burden brought on the world by the COVID-19 pandemic, the world has faced the reality that, at times, internet interventions might be the only available option when addressing mental health issues when face-to-face contact is not recommended (Wind *et al.*, 2020). In general, internet-delivered interventions are seen as a more cost-effective alternative to face-to-face therapy (Andersson *et al.*, 2014). Being also more accessible, internet interventions are seen as particularly valuable when addressing the lack of trained therapists, especially in rural or remote locations (Lattie *et al.*, 2022).

Numerous randomised controlled trials (RCTs) have demonstrated the efficacy of internet-delivered interventions on various mental health outcomes, such as anxiety and mood disorders (Andersson *et al.*, 2019). A meta-analysis of 13 RCTs published between 2010 and 2021 confirmed the efficacy of internet interventions targeting elevated stress as a primary outcome and also documented secondary effects on measures of anxiety and depression (Svärdman *et al.*, 2022). Moreover, in nine out of 10 studies with follow-up measures, the effects of internet interventions on stress sustained ($n = 4$) or even slightly increased ($n = 5$) over time (Svärdman *et al.*, 2022). Nevertheless, most existing studies have focused on the effectiveness of interventions without examining how the effects are sustained or how different mental health symptoms interact. Moreover, it has been suggested that alongside a reduction of symptoms, an increase in well-being should be confirmed to establish intervention effects (Cloninger, 2006). However, the interplay between the increase in psychological well-being and the reduction of symptoms is rarely addressed in intervention research, including research on the effects of CBT-based internet interventions.

To address these gaps in previous research, the current study aimed to investigate longitudinal links between symptoms of stress, anxiety, and depression, as well as psychological well-being, in a help-seeking sample of medical nurses who participated in the internet-delivered CBT-based intervention FOREST, targeting stress reduction (Jovarauskaite *et al.*, 2021; the description of the intervention FOREST is provided in the Method section). In the current study, we defined stress as the subjective feeling of lack of control and confidence to handle difficulties in important areas of life (Cohen and Williamson, 1988). Anxiety was defined as a feeling of nervousness and inability to control worrying; and depression as a lack of interest or pleasure when doing things and feeling down or hopeless (Kroenke *et al.*, 2009). Psychological well-being was defined as the subjective experience of positive emotional states, vitality, and sense of purpose (Bech, 2004).

As was found in the previously conducted RCT, FOREST had immediate and follow-up effects, decreasing stress and depression and increasing well-being (Dumarkaite *et al.*, 2023). Symptoms of depression and anxiety tend to be interlinked over time (Jacobson and Newman, 2017). Therefore, we expected that anxiety and depression would predict each other over time. There is abundant literature linking exposure to stressful life events and mental health. Some longitudinal research shows that stressful events precede mental health deterioration (e.g. Schneider *et al.*, 2021), and some report bi-directionality of associations between exposure to stress and mental health (e.g. Schönfeld *et al.*, 2019). However, there has been less research on longitudinal links between subjective stress and mental health. Some evidence shows that subjective stress is an even stronger predictor of overall health than exposure to stressful events (Shields *et al.*, 2023). Also, the evidence from prospective longitudinal research showed that higher levels of subjective stress were a risk factor for future mental health problems in healthcare workers (Van Steenkiste, 2022). Keeping this in mind and also considering that the current study was conducted in the context of stress recovery intervention, we expected that lower levels of subjective stress would predict a decrease in depression and anxiety symptoms and an increase in well-being over time. Nevertheless, most studies that inform intervention research are conducted in non-intervention settings. Addressing longitudinal links between stress, anxiety, depression, and well-being in the help-seeking sample that reported intervention-induced changes in mental health indicators

would allow a better understanding of what further outcomes we might expect after fostering mental health.

Method

Participants

The participants of the internet-delivered stress recovery program FOREST (Dumarkaite *et al.*, 2023; Jovarauskaite *et al.*, 2021) were included in the current study. Participants were nurses and assistant nurses from both study arms of the RCT: an intervention group (IG) and a delayed treatment group (DTG), which received the intervention after a 3-month follow-up assessment. The inclusion criteria were as follows: (1) being a licensed nurse actively working within the national healthcare system; (2) providing written informed consent electronically; (3) completing a baseline assessment prior to randomization; (4) sufficient proficiency in Lithuanian to understand the intervention content and instructions; (5) possessing a computer, tablet, smartphone, or similar device with internet access. Exclusion criteria were as follows: (1) experiencing an acute psychiatric crisis; (2) a high risk of suicide; (3) alcohol or drug addiction; (4) current experience of interpersonal violence. Overall, 208 applicants registered for participation, and 184 met the inclusion and did not meet the exclusion criteria (93 were assigned to the IG and 91 to the DTG). Of these, 73 participants were excluded from the analysis because (1) they were from the DTG group and had not completed pre-test measurement questionnaires ($n = 29$); (2) they had never logged in to the program ($n = 23$); (3) they had not completed the post-test and/or follow-up assessment ($n = 21$). The final sample of the current study consisted of 111 participants (65 from the IG and 46 from the DTG). The study flowchart is presented in Fig. S1 in the Supplementary material. Demographic characteristics of the current study sample ($n = 111$) collected at enrolment are presented in Table 1.

We found no differences between participants of the current study ($n = 111$) and excluded participants ($n = 73$) at the enrolment in terms of demographic characteristics, including gender ($\chi^2(1) = 0.89, p = .35$), age ($t_{182} = 1.21, p = .23$), job position ($\chi^2(1) = 0.00, p = .98$), education ($\chi^2(2) = 0.98, p = .62$), working status ($\chi^2(2) = 1.99, p = .37$), work experience ($\chi^2(3) = 3.86, p = .28$), subjective income ($\chi^2(2) = 0.13, p = .94$), partnership status ($\chi^2(1) = 0.22, p = .64$), seeing a psychologist ($\chi^2(1) = 2.84, p = .09$), and usage of psychopharmacological medicine due to mental health difficulties ($\chi^2(1) = 1.08, p = .30$). Also, there were no differences between included and excluded participants regarding any outcome measures at the enrolment: perceived stress ($M (SD) = 7.86 (2.33)$ vs $M (SD) = 7.89 (3.20)$; $t_{121.58} = 0.06, p = .95$), anxiety ($M (SD) = 2.88 (1.56)$ vs $M (SD) = 2.63 (1.95)$; $t_{130.36} = -0.93, p = .35$), depression ($M (SD) = 2.56 (1.49)$ vs $M (SD) = 2.56 (1.75)$; $t_{182} = 0.13, p = .99$), and psychological well-being ($M (SD) = 9.41 (4.65)$ vs $M (SD) = 9.84 (5.22)$; $t_{182} = 0.59, p = .56$).

Procedures

The participants were all from Lithuania and were enrolled via disseminating information through press releases to media, healthcare institutions, and specialised social networks. Detailed enrolment procedures and participant eligibility criteria have been reported previously (Dumarkaite *et al.*, 2023; Jovarauskaite *et al.*, 2021). Informed consent was obtained from all participants before data collection. In the current study, the IG data collected at T1, T2 and T3 and the DTG data collected at T3, T4 and T5 were merged into a single data pool and represented pre-test, post-test, and follow-up, respectively. Thus, the current study presents the secondary analysis of the IG data collected at T1 and T2, as well as IG and DTG data collected at T3, while the data collected at T4 and T5 has not been analysed previously.

Table 1. Demographic characteristics of the participants at the enrolment ($n = 111$)

Variable	n (%)
Gender	
Female	109 (98.2)
Male	2 (1.8)
Age	
M (SD)	41.69 (10.85)
Range	23–59
Position	
Assistant nurse	6 (5.4)
Nurse	105 (94.6)
Education	
Secondary or lower	1 (0.9)
Higher or non-university higher	65 (58.6)
Higher university	45 (40.5)
Working status	
Part-time	6 (5.4)
Full-time	45 (40.5)
More than full-time	60 (54.1)
Working experience	
<2 years	14 (12.6)
2–5 years	15 (13.5)
6–10 years	13 (11.7)
>10 years	69 (62.2)
Income	
Lower than average	13 (11.7)
Average	81 (73.0)
Higher than average	17 (15.3)
Partnership (long-term)	
No	27 (24.3)
Yes	84 (75.7)
Visiting a psychologist	
No	101 (91.0)
Yes	10 (9.0)
Taking psychopharmacological medicine	
No	106 (95.5)
Yes	5 (4.5)

Program FOREST

The online cognitive behavior therapy-based stress recovery program used in the study FOREST aimed to enhance the healthcare staff's stress recovery skills. FOREST also included mindfulness principles that have been shown to assist in the modulation of brain-behaviour interaction involved in recovery from stress (Johnson *et al.*, 2014). FOREST consisted of six sessions and lasted six weeks. Each session was unlocked on a weekly basis and covered the following topics: (1) Introduction; (2) Relaxation; (3) Psychological detachment; (4) Mastery; (5) Control; (6) Keeping the change alive. The topics of sessions 2 to 5 represented the four elements of the stress recovery experiences (Sonnetag and Fritz, 2007). FOREST consisted of three main elements that can be found throughout the whole program: psychoeducation (text and video recordings), exercises (written and audio recordings), and communication with a psychologist. The psychologist's role included providing feedback to the participants once they completed exercises and replying to personal messages from program participants. FOREST was developed specifically for healthcare staff experiencing stress by clinical psychologists and researchers at Vilnius University. More details on the program and its entire content can be found in the study protocol (Jovarauskaite *et al.*, 2021) and the FOREST RCT study (Dumarkaitė *et al.*, 2023).

Measures

Stress

The Perceived Stress Scale (PSS-4) (Cohen and Williamson, 1988) was used to evaluate the changes in self-reported perceived stress. The PSS-4 consists of four questions (e.g. 'In the last month, how often have you felt that you were unable to control the important things in your life?'); each question is ranked on a 5-point Likert scale ranging from 0 ('never') to 4 ('very often'). A higher score indicates more pronounced perceived stress. In the current study, Cronbach's alpha for the PSS-4 at pre-test was $\alpha = .65$.

Symptoms of anxiety and depression

The Patient Health Questionnaire (PHQ-4) (Kroenke *et al.*, 2009) was used to evaluate the changes in self-reported anxiety and depression symptoms. The PHQ-4 consists of four items and two subscales: the anxiety subscale (e.g. 'Feeling nervous, anxious or on edge') and depression subscale (e.g. 'Little interest or pleasure in doing things'); each item is ranked on a 4-point Likert scale ranging from 0 ('not at all') to 3 ('nearly every day'). A higher score indicates more pronounced anxiety and depression symptoms. In the current study, Cronbach's alpha for the PHQ-4 anxiety subscale at pre-test was $\alpha = .83$, and for the PHQ-4 depression subscale at pre-test was $\alpha = .75$.

Psychological well-being

The World Health Organization Well-being Index (WHO-5) (Bech, 2004) was used to evaluate the changes in self-reported psychological well-being. The WHO-5 consists of five items (e.g. 'I have felt cheerful and in good spirits'); each item is ranked on a 6-point Likert scale ranging from 0 ('at no time') to 5 ('all of the time'). A higher score indicates more pronounced psychological well-being. In the current study, Cronbach's alpha for the WHO-5 at pre-test was $\alpha = .87$.

Data analyses

The current study aimed to investigate the longitudinal interplay between mental health indicators when participating in the internet-based stress-reduction intervention. As a preliminary analysis, we assessed the change on four indicators: symptoms of stress, anxiety, and depression, as well as the level of psychological well-being using the latent change modelling approach (Duncan *et al.*, 2013). We ran a series of latent change models, indicating the estimated baseline levels and the estimated change in variables from pre- to post-test and from pre-test to follow-up.

To analyse the interplay between mental health indicators over time, we conducted a cross-lagged panel analysis (Selig and Little, 2012), with four variables representing symptoms of stress, anxiety, and depression, as well as the level of psychological well-being measured at three time-points, i.e. pre-test, post-test, and 3-month follow-up. When running the autoregressive cross-lagged model, the cross-lagged effects were estimated by controlling for the stability paths between the pre-test and post-test and between post-test and follow-up, as well as within-time correlations among all variables at all measurement points. All analyses were conducted using the sum scores of the variables with the robust maximum likelihood estimator in Mplus 8.2 (Muthén and Muthén, 1998–2017). The model fit of the cross-lagged model was evaluated using the comparative fit index (CFI), the Tucker–Lewis index (TLI), the root mean square error of approximation (RMSEA), and χ^2 test (Kline, 2011). The CFI/TLI values higher than .90 and RMSEA values below .08 indicate an acceptable fit; CFI/TLI values higher than .95, RMSEA values below .06, and insignificant χ^2 test represent a good fit. The full information maximum likelihood (FIML) algorithm was used for handling missing data in the current study.




Figure 1. The univariate trajectories of stress, anxiety, depression, and psychological well-being over the study period ($n = 111$). The variables were measured in different scales and are incomparable with one another; stars next to the estimated means represent the difference at post-test or follow-up in comparison with pre-test: ** $p < .01$, *** $p < .001$.




Figure 2. Standardised significant estimates of the cross-lagged panel model ($n = 111$). Numbers on the lines represent standardised beta coefficients; numbers within rectangles represent standardised R^2 coefficients; * $p < .05$, ** $p < .01$, *** $p < .001$.

Results

The latent change analysis indicated statistically significant reductions in stress, anxiety, and depression symptoms and a significant increase in well-being from pre- to post-test (see Fig. 1). At follow-up, the decreases in stress and depression symptoms and the increase in well-being were sustained, while anxiety levels returned to pre-intervention levels. The results of the autoregressive cross-lagged panel analysis indicated that the model fitted data well ($\chi^2(16) = 17.43, p = .358, CFI/TLI = .997/.988, RMSEA [90\% CI] = .028 [.000, .094]$). Significant stability paths and cross-lagged effects are presented in Fig. 2. The within-time correlations are reported in Table 2. As shown in

Table 2. Within-time cross-lagged panel correlations between study variables

	Variable at the same time point		
	Stress	Anxiety	Depression
T1 Anxiety	.53	—	—
T1 Depression	.58	.72	—
T1 Well-being	-.50	-.56	-.56
T2 Anxiety	.33	—	—
T2 Depression	.49	.56	—
T2 Well-being	-.35	-.33	-.42
T3 Anxiety	.62	—	—
T3 Depression	.59	.61	—
T3 Well-being	-.55	-.67	-.50

All correlations are significant at $p < .001$.

Fig. 2, the levels of anxiety symptoms at the post-test positively predicted the levels of stress and depression symptoms at follow-up. Additionally, the levels of psychological well-being at the post-test negatively predicted stress symptoms at follow-up. No cross-lagged effects were observed from the pre-test to the post-test.

Discussion

The current study aimed to investigate the longitudinal links between symptoms of stress, anxiety, and depression and the psychological well-being of the participants of the CBT-based internet intervention FOREST, targeting stress recovery. Our previous RCT study (Dumarkaite *et al.*, 2023) confirmed that the intervention positively affected nurses' mental health and the effects on stress and depression symptoms, as well as well-being, sustained at a 3-month follow-up. The same within-group effects were confirmed in the current study using the combined sample of intervention participants who received intervention right after enrolment and delayed treatment group data. Overall, the results demonstrate that reduced anxiety and increased well-being predict the sustainability of the effects of reduced stress. Additionally, we found that reduced anxiety appears to contribute to sustaining intervention effects on reduced depression symptoms.

In contrast to our expectations, we did not find that reduced subjective stress was linked with decreased symptoms of poor mental health. In contrast, increased mental health contributed to sustaining lower subjective stress levels achieved through participation in the intervention. Even though there is strong evidence that exposure to stressful events precedes the symptoms of anxiety and depression (Schneider *et al.*, 2021), our results suggest that when addressing elevated subjective stress, we need to target other mental health outcomes, such as anxiety and psychological well-being if we want effects to be sustained. Also, contrary to our expectations, we did not find bi-directional longitudinal links between anxiety and depression symptoms. The results suggest that reduced anxiety plays a role in sustaining intervention effects on depression symptoms but not vice versa.

Previous research has demonstrated that one of the primary mechanisms behind improved mental health following mindfulness-based stress-reduction interventions is decreased repetitive negative thinking (Gu *et al.*, 2015). It is also known that the reduction of repetitive negative thinking is usually paired with reduced anxiety (Monteregge *et al.*, 2020). Although the design of the current study did not allow testing mediation mechanisms of intervention effects, it has been shown that reduced anxiety is one of the predictors of sustained intervention effects. Therefore, in stress reduction intervention research, more attention should be paid to targeting decreased anxiety through reduced repetitive negative thinking or other potential mechanisms.

The results also highlighted the importance of improved psychological well-being when targeting elevated stress and promoting mental health. There are many mechanisms for tackling well-being (Diener and Biswas-Diener, 2019). Some of them, such as CBT and mindfulness, were also used in the FOREST intervention. The results suggest that increased well-being contributes to sustaining reduced stress. From the perspective of the experiences of recovery from stress model (Sonnentag and Fritz, 2007), positive emotionality, which is an internal part of psychological well-being (Bech, 2004), might be seen as internal resources, providing a sense of self-esteem and self-confidence (Philippe *et al.*, 2018) and thus might assist in coping with everyday challenges and protecting from stress (Cohen and Williamson, 1988). Therefore, future intervention research should consider measuring psychological well-being as a secondary intervention outcome.

In summary, based on the findings of the current study, when applying internet-delivered interventions targeted at sustained stress reduction in clinical research and practice settings, specialists should consider incorporating more anxiety-reduction strategies. Moreover, interventions targeting stress reduction should consider focusing not only on immediate relief but also on longer-term strategies that could potentially increase overall well-being. Also, based on the findings of the current study, researchers and clinicians should consider monitoring anxiety and well-being as early indicators of whether the stress-reduction intervention has the potential to have long-lasting results. Finally, when personalising the interventions targeting stress, the baseline level of anxiety and well-being could be used as indicators when offering different levels of support or booster sessions to individuals with higher levels of anxiety or lower levels of well-being.

The current study's findings should be seen in light of its limitations. The study's main limitation is the combined study sample, resulting in different measurement times for the intervention group (IG) and delayed treatment group (DTG). Even though we did not find any differences between groups at pre-test and post-test and, in most cases, at follow-up, stress levels were higher at follow-up in the DTG, possibly indicating that long-term effects on stress were more sustainable in the IG, compared with DTG. Even though the waiting conditions do not seem to have negative effects on intervention outcomes when participating in stress reduction interventions (Elliott and Brown, 2002), in help-seeking samples (this was also the case in the current study), waiting for the intervention might introduce some dissatisfaction and, in turn, the final intervention effects might be less beneficial (Gunnarsson *et al.*, 2023). Additionally, the study was conducted with a sample of self-referred nurses, indicating the risk of volunteering bias and the results should be generalised to other samples with caution. Moreover, the very short versions of stress, anxiety, and depression measures were used in the study and could limit the comparison of the results with other studies. Finally, the follow-up effects of the study were relatively short-term. Future studies should test the predictors of the sustained intervention effects with longer follow-ups.

Conclusion

The results of the current study provide evidence of the importance of reducing anxiety and promoting psychological well-being when targeting stress in internet interventions, as the post-intervention levels of anxiety and psychological well-being are associated with the sustainability of the intervention effects on reduced stress over time.

Supplementary material. The supplementary material for this article can be found at <https://doi.org/10.1017/S1352465824000456>

Data availability statement. The data that support the findings of this study are available from the corresponding author, upon reasonable request.

Acknowledgements. None.

Author contributions. **Inga Truskauskaitė:** Conceptualization-Lead, Data curation-Equal, Formal analysis-Lead, Funding acquisition-Equal, Investigation-Equal, Methodology-Equal, Project administration-Supporting, Visualization-Lead, Writing - original draft-Lead, Writing - review & editing-Lead; **Austeja Dumarkaitė:** Conceptualization-Supporting, Data curation-Equal, Investigation-Lead, Methodology-Supporting, Project administration-Equal, Writing - review & editing-Supporting; **Auguste Nomeikaitė:** Investigation-Supporting, Project administration-Supporting, Writing - review & editing-Supporting; **Gerhard Andersson:** Funding acquisition-Supporting, Methodology-Supporting, Resources-Equal, Supervision-Supporting, Validation-Supporting, Writing - review & editing-Supporting; **Evaldas Kazlauskas:** Conceptualization-Supporting, Funding acquisition-Lead, Investigation-Equal, Methodology-Equal, Project administration-Equal, Resources-Equal, Supervision-Lead, Validation-Lead, Writing - review & editing-Equal.

Financial support. This work was supported by the European Regional Development Fund according to the supported activity 'Research Projects Implemented by World-class Researcher Groups' under grant no. 01.2.2-LMT-K-718 (project no. 01.2.2-LMT-K-718-03-0072).

Competing interests. All authors declare no competing interests.

Ethical standards. The study was approved by the Psychology Research Ethics Committee of Vilnius University (reference no. 2021-03-22/61). All participants gave informed consent to participate in the study and for the results to be published. The authors assert that all procedures contributing to this work comply with the ethical standards of the relevant national and institutional committees and with the Helsinki Declaration of 1975, as revised in 2008. The trial was registered at www.clinicaltrials.gov (NCT04817995: <https://clinicaltrials.gov/ct2/show/NCT04817995>, 30 March 2021).

References

- American Psychological Association (2021). *Worsening mental health crisis pressures the psychologist workforce*. <https://www.apa.org/pubs/reports/practitioner/covid-19-2021>
- Andersson, G. (2018). Internet interventions: past, present and future. *Internet Interventions*, 12, 181–188. <https://doi.org/10.1016/j.invent.2018.03.008>
- Andersson, G., Carlbring, P., Titov, N., & Lindfors, N. (2019). Internet interventions for adults with anxiety and mood disorders: a narrative umbrella review of recent meta-analyses. *Canadian Journal of Psychiatry*, 64, 465–470. <https://doi.org/10.1177/0706743719839381>
- Andersson, G., Cuijpers, P., Carlbring, P., Riper, H., & Hedman, E. (2014). Guided internet-based vs. face-to-face cognitive behavior therapy for psychiatric and somatic disorders: a systematic review and meta-analysis. *World Psychiatry*, 13, 288–295. <https://doi.org/10.1002/wps.20151>
- Bech, P. (2004). Measuring the dimension of psychological general well-being by the WHO-5. *Quality of Life Newsletter*, 15–16.
- Cloninger, C. R. (2006). The science of well being: an integrated approach to mental health and its disorders. *Psychiatria Danubina*, 5, 71–76.
- Cohen, J. I. (2000). Stress and mental health: a biobehavioral perspective. *Issues in Mental Health Nursing*, 21, 185–202. <https://doi.org/10.1080/016128400248185>
- Cohen, S., & Williamson, G. (1988). Perceived stress in a probability sample of the United States. In Spacapan, S. & Oskamp, S. (eds), *The Social Psychology of Health* (pp. 31–68). SAGE.
- Diener, E., & Biswas-Diener, R. (2019). Well-being interventions to improve societies. *Global Happiness Council, Global Happiness and Well-Being Policy Report*, 95–110.
- Dumarkaitė, A., Truskauskaitė, I., Andersson, G., Jovarauskaite, L., Jovaisiene, I., Nomeikaitė, A., & Kazlauskas, E. (2023). The efficacy of the internet-based stress recovery intervention FOREST for nurses amid the COVID-19 pandemic: a randomized controlled trial. *International Journal of Nursing Studies*, 138, 104408. <https://doi.org/10.1016/j.ijnurstu.2022.104408>
- Duncan, T. E., Duncan, S. C., & Strycker, L. A. (2013). *An Introduction to Latent Variable Growth Curve Modeling: Concepts, Issues, and Application*. Routledge.
- Elliott, S. A., & Brown, J. S. L. (2002). What are we doing to waiting list controls? *Behaviour Research and Therapy*, 40, 1047–1052. [https://doi.org/10.1016/S0005-7967\(01\)00082-1](https://doi.org/10.1016/S0005-7967(01)00082-1)
- Estevez Cores, S., Sayed, A. A., Tracy, D. K., & Kempton, M. J. (2021). Individual-focused occupational health interventions: a meta-analysis of randomized controlled trials. *Journal of Occupational Health Psychology*, 26, 189–203. <https://doi.org/10.1037/ocp0000249>
- Gu, J., Strauss, C., Bond, R., & Cavanagh, K. (2015). How do mindfulness-based cognitive therapy and mindfulness-based stress reduction improve mental health and wellbeing? A systematic review and meta-analysis of mediation studies. *Clinical Psychology Review*, 37, 1–12. <https://doi.org/10.1016/j.cpr.2015.01.006>
- Gunnarsson, K. U., McCambridge, J., & Bendtsen, M. (2023). Reactions to being allocated to a waiting list control group in a digital alcohol intervention trial. *Patient Education and Counseling*, 107, 107572. <https://doi.org/10.1016/j.pcc.2022.11.014>

- Haight, B. L., Peddie, L., Crosswell, A. D., Hives, B. A., Almeida, D. M., & Puterman, E. (2023). Combined effects of cumulative stress and daily stressors on daily health. *Health Psychology, 42*, 325–334. <https://doi.org/10.1037/hea0001281>
- Hassard, J., Teoh, K. R. H., Visockaite, G., Dewe, P., & Cox, T. (2018). The cost of work-related stress to society: a systematic review. *Journal of Occupational Health Psychology, 23*, 1–17. Educational Publishing Foundation. <https://doi.org/10.1037/ocp0000069>
- Jacobson, N. C., & Newman, M. G. (2017). Anxiety and depression as bidirectional risk factors for one another: a meta-analysis of longitudinal studies. *Psychological Bulletin, 143*, 1155–1200. <https://doi.org/10.1037/bul0000111>
- Johnson, D. C., Thom, N. J., Stanley, E. A., Haase, L., Simmons, A. N., Shih, P. B., Thompson, W. K., Potterat, E. G., Minor, T. R., & Paulus, M. P. (2014). Modifying resilience mechanisms in At-Risk individuals: a controlled study of mindfulness training in marines preparing for deployment. *American Journal of Psychiatry, 171*, 844–853. <https://doi.org/10.1176/appi.ajp.2014.13040502>
- Jovarauskaite, L., Dumarkaite, A., Truskauskaitė-Kuneviciene, I., Jovaisiene, I., Andersson, G., & Kazlauskas, E. (2021). Internet-based stress recovery intervention FOREST for healthcare staff amid COVID-19 pandemic: study protocol for a randomized controlled trial. *Trials, 22*, 559. <https://doi.org/10.1186/s13063-021-05512-1>
- Kline, R. B. (2011). *Principles and Practice of Structural Equation Modeling* (3rd edn). Guilford Press.
- Kroenke, K., Spitzer, R. L., Williams, J. B. W., & Löwe, B. (2009). An ultra-brief screening scale for anxiety and depression: the PHQ-4. *Psychosomatics, 50*, 613–621.
- Lattie, E. G., Stiles-Shields, C., & Graham, A. K. (2022). An overview of and recommendations for more accessible digital mental health services. *Nature Reviews Psychology, 1*, 87–100. <https://doi.org/10.1038/s44159-021-00003-1>
- Mingote Adán, J. C., Moreno Jiménez, B., & Gálvez Herrer, M. (2004). Burnout and the health of the medical professionals: review and preventive options. *Medicina Clinica, 123*, 265–270. <https://doi.org/10.1157/13065203>
- Monteregge, S., Tsagkalidou, A., Cuijpers, P., & Spinhoven, P. (2020). The effects of different types of treatment for anxiety on repetitive negative thinking: a meta-analysis. *Clinical Psychology: Science and Practice, 27*, 1–20. <https://doi.org/10.1111/cpsp.12316>
- Muthén, L. K., & Muthén, B. O. (1998–2017). *Mplus User's Guide*, 8th edn. Los Angeles, CA: Muthén & Muthén.
- O'Connor, D. B., Thayer, J. F., & Vedhara, K. (2021). Stress and health: a review of psychological processes. *Annual Review of Psychology, 72*, 663–688. <https://doi.org/10.1146/annurev-psych-062520-122331>
- Philippe, F. L., Dobbin, A. E., Ross, S., & Houle, I. (2018). Resilience facilitates positive emotionality and integration of negative memories in need satisfying memory networks: an experimental study. *Journal of Positive Psychology, 13*, 586–596. <https://doi.org/10.1080/17439760.2017.1365158>
- Schneider, R. L., Long, E. E., Arch, J. J., & Hankin, B. L. (2021). The relationship between stressful events, emotion dysregulation, and anxiety symptoms among youth: longitudinal support for stress causation but not stress generation. *Anxiety, Stress and Coping, 34*, 157–172. <https://doi.org/10.1080/10615806.2020.1839730>
- Schönfeld, P., Brailovskaia, J., Zhang, X. C., & Margraf, J. (2019). Self-efficacy as a mechanism linking daily stress to mental health in students: a three-wave cross-lagged study. *Psychological Reports, 122*, 2074–2095. <https://doi.org/10.1177/0033294118787496>
- Selig, J. P., & Little, T. D. (2012). Autoregressive and cross-lagged panel analysis for longitudinal data. In *Handbook of Developmental Research Methods* (pp. 265–278). Guilford Press.
- Shields, G. S., Fasset-Carman, A., Gray, Z. J., Gonzales, J. E., Snyder, H. R., & Slavich, G. M. (2023). Why is subjective stress severity a stronger predictor of health than stressor exposure? A preregistered two-study test of two hypotheses. *Stress and Health, 39*, 87–102. <https://doi.org/10.1002/smi.3165>
- Sonnentag, S., & Fritz, C. (2007). The recovery experience questionnaire: development and validation of a measure for assessing recuperation and unwinding from work. *Journal of Occupational Health Psychology, 12*, 204–221. <https://doi.org/10.1037/1076-8998.12.3.204>
- Svärdman, F., Sjöwall, D., & Lindsäter, E. (2022). Internet-delivered cognitive behavioral interventions to reduce elevated stress: a systematic review and meta-analysis. *Internet Interventions, 29*, 100553. <https://doi.org/10.1016/j.invent.2022.100553>
- Van Steenkiste, E., Schoofs, J., Gilis, S., & Messiaen, P. (2022). Mental health impact of COVID-19 in frontline healthcare workers in a Belgian Tertiary care hospital: a prospective longitudinal study. *Acta Clinica Belgica, 77*, 533–540. <https://doi.org/10.1080/17843286.2021.1903660>
- Wind, T. R., Rijkeboer, M., Andersson, G., & Riper, H. (2020). The COVID-19 pandemic: the 'black swan' for mental health care and a turning point for e-health. *Internet Interventions, 20*, 100317. <https://doi.org/10.1016/j.invent.2020.100317>

Cite this article: Truskauskaitė I, Dumarkaite A, Nomeikaite A, Andersson G, and Kazlauskas E (2024). Longitudinal interplay between subjective stress, anxiety, depression, and well-being in internet-based stress recovery intervention for nurses. *Behavioural and Cognitive Psychotherapy*. <https://doi.org/10.1017/S1352465824000456>