

THE 67TH INTERNATIONAL



OPEN READINGS

CONFERENCE FOR STUDENTS OF PHYSICS AND NATURAL SCIENCES

**BOOK OF
ABSTRACTS** | **2024**



Vilnius
University

VILNIUS UNIVERSITY PRESS

Editors:

Martynas Keršys
Rimantas Naina
Vincentas Adomaitis
Emilijus Maskvytis

Cover and Interior Design:

Goda Grybauskaitė

Vilnius University Press
9 Saulėtekio Av., III Building, LT-10222 Vilnius
info@leidykla.vu.lt, www.leidykla.vu.lt/en/
www.knygynas.vu.lt, www.journals.vu.lt

Bibliographic information is available
on the Lithuanian Integral Library Information System (LIBIS) portal www.ibiblioteka.lt
ISBN 978-609-07-1051-7 (PDF)

© Vilnius University, 2024

THE STUDY OF THE ATMOSPHERES AND MIXING PROCESSES OF MAGNETICALLY ACTIVE RS CVN GIANTS

Barkha Bale¹, Grazina Tautvaisiene¹, Renata Minkeviciute¹, Arnas Drazdauskas¹, Edit Stonkute¹, Sarunas Mikolaitis¹

¹Institute of Theoretical Physics and Astronomy (ITPA), Vilnius University
barkha.bale@ff.stud.vu.lt

Our primary focus is the examination of RS CVn stars, aiming to ascertain the carbon isotope $^{12}\text{C}/^{13}\text{C}$ and C/N ratios within these chromospherically active celestial bodies. Analyzing the abundances of carbon and nitrogen in their atmospheres offers valuable insights into chemical composition alterations induced by stellar evolution. The determinations of carbon and nitrogen abundances, along with the C/N ratios, serve as pivotal tools in the field of stellar evolution studies. To explore the influence of magnetic activity on mixing processes within the atmospheres of magnetically active stars, we are conducting an exhaustive investigation of CNO abundances and carbon isotope ratios in a selected group of RS CVn giant stars. High-resolution spectra, acquired from the 1.65 m telescope at the Moletai Astronomical Observatory of Vilnius University, were utilized for this study. The analysis focused on the C2 Swan (1,0) band head at 5135 Å and the C2 Swan (0,1) band head at 5635.5 Å to determine carbon abundance. Additionally, the interval 7990 – 8010 Å, encompassing the $^{12}\text{C}/^{14}\text{N}$ and $^{13}\text{C}/^{14}\text{N}$ bands, was employed for nitrogen abundance and carbon isotope ratio analysis. Oxygen abundance was derived from the forbidden [O I] line at 6300.31 Å. Our findings support earlier observations, indicating that in low-mass chromospherically active RS CVn stars, extra-mixing processes may commence below a luminosity function bump in red giants.




Fig. 1. The investigated stars in the log g versus Teff diagram along with the PADOVA evolutionary tracks taken from Bressan et al. (2012). The red symbols indicate the stars which are below the red giant branch (RGB) luminosity bump, the blue colour indicates the stars which are at the RGB bump, and the black colour marks the stars that are above the RGB bump.