

THE 67<sup>TH</sup> INTERNATIONAL



# OPEN READINGS

CONFERENCE FOR STUDENTS OF PHYSICS AND NATURAL SCIENCES

**BOOK OF  
ABSTRACTS** | **2024**



Vilnius  
University

VILNIUS UNIVERSITY PRESS

Editors:

Martynas Keršys  
Rimantas Naina  
Vincentas Adomaitis  
Emilijus Maskvytis

Cover and Interior Design:

Goda Grybauskaitė

Vilnius University Press  
9 Saulėtekio Av., III Building, LT-10222 Vilnius  
info@leidykla.vu.lt, [www.leidykla.vu.lt/en/](http://www.leidykla.vu.lt/en/)  
[www.knygynas.vu.lt](http://www.knygynas.vu.lt), [www.journals.vu.lt](http://www.journals.vu.lt)

Bibliographic information is available  
on the Lithuanian Integral Library Information System (LIBIS) portal [www.ibiblioteka.lt](http://www.ibiblioteka.lt)  
ISBN 978-609-07-1051-7 (PDF)

© Vilnius University, 2024

# NIELSEN IDENTITIES IN THE STANDARD MODEL

Nojus Danyla<sup>1</sup>, Simonas Draukšas<sup>2</sup>

<sup>1</sup>Faculty of Physics, Vilnius University, Lithuania

<sup>2</sup>Institute of Theoretical Physics and Astronomy, Vilnius University, Lithuania

[nojus.danyla@gmail.com](mailto:nojus.danyla@gmail.com)

It is not always clear whether definitions of physical quantities is really physical, for example, it may be that the definition is gauge-dependent. One of investigating gauge-dependence is the use of Nielsen Identities [1] --- a generalised form of Slavnov-Taylor Identities [2], which come from BRST symmetry [3, 4]. These Identities enable us to express dependence on the gauge parameter in terms correlation functions containing BRST sources related to the gauge parameters.

This study focuses on understanding Nielsen Identities in the Standard Model, in which we investigate the gauge dependence of quark self-energies. We explore dependencies of quark self-energies on the gauge parameter in two distinct ways: 1) by calculating the Nielsen identities via a BRST Lagrangian; 2) by calculating quark self-energies and taking the derivative with respect to the gauge parameter. The goal is to check the consistency between the results obtained through both approaches.

In this presentation, we will introduce all the needed concepts as well as our progress.

- 
- [1] . Nielsen, On the gauge dependence of spontaneous symmetry breaking in gauge theories, Nuclear Physics B, Dec. 1975, 101, 173–188.
  - [2] . Gambino and P. A. Grassi, Nielsen identities of the SM and the definition of mass, Physical Review D, Aug. 2000, 62, 076002.
  - [3] . Becchi, A. Rouet, and R. Stora, Renormalization of gauge theories, Annals of Physics, June 1976, 98, 287–321.
  - [4] . V. Tyutin, Gauge Invariance in Field Theory and Statistical Physics in Operator Formalism, 2008.