THE 67TH INTERNATIONAL

OPEN READINGS



CONFERENCE FOR STUDENTS OF PHYSICS AND NATURAL SCIENCES

BOOK OF 2024 ABSTRACTS



VILNIUS UNIVERSITY PRESS

Editors:

Martynas Keršys Rimantas Naina Vincentas Adomaitis Emilijus Maskvytis

Cover and Interior Design:

Goda Grybauskaitė

Vilnius University Press 9 Saulėtekio Av., III Building, LT-10222 Vilnius info@leidykla.vu.lt, www.leidykla.vu.lt/en/ www.knygynas.vu.lt, www.journals.vu.lt

Bibliographic information is available on the Lithuanian Integral Library Information System (LIBIS) portal www.ibiblioteka.lt ISBN 978-609-07-1051-7 (PDF)

© Vilnius University, 2024

EVALUATING METHODS FOR ESTIMATING THE HURST EXPONENT IN TIME SERIES A COMPARATIVE ANALYSIS OF ACCURACY AND APPLICATION

Danielius Kundrotas¹, Rytis Kazakevičius²

¹Faculty of Physics, Vilnius University, Saulėtekio av. 9, Vilnius LT-10222, Lithuania
²Institute of Theoretical Physics and Astronomy, Vilnius University, Saulėtekio av. 3, Vilnius LT-10257, Lithuania danielius.kundrotas@ff.stud.vu.lt

The Hurst exponent (H) is a statistical measure used to understand the long-term memory of time series, characterized by persistence and anti-persistence phenomena. Accurate estimation of H is crucial in diverse fields, such as finance, geophysics, and telecommunications, due to its ability to indicate the future tendency of a system based on its historical behavior. Numerous methods have been developed for estimating H, each with varying degrees of complexity and accuracy. This research reviews these methods, focusing on popular ones like box-counting, Katz, detrended fluctuation analysis, mean squared displacement (MSD), and Higuchi. To evaluate the reliability of these methods, fractional Brownian motion was used as a benchmark due to its well-defined Hurst parameter. Our analysis revealed that the Higuchi method and MSD provided the most accurate their effectiveness. This work contributes to the field by identifying the most reliable numerical methods for estimating the Hurst exponent, providing a foundation for future studies and practical applications in analyzing time series with long-range dependencies.