

THE 67TH INTERNATIONAL



OPEN READINGS

CONFERENCE FOR STUDENTS OF PHYSICS AND NATURAL SCIENCES

**BOOK OF
ABSTRACTS** | **2024**



Vilnius
University

VILNIUS UNIVERSITY PRESS

Editors:

Martynas Keršys
Rimantas Naina
Vincentas Adomaitis
Emilijus Maskvytis

Cover and Interior Design:

Goda Grybauskaitė

Vilnius University Press
9 Saulėtekio Av., III Building, LT-10222 Vilnius
info@leidykla.vu.lt, www.leidykla.vu.lt/en/
www.knygynas.vu.lt, www.journals.vu.lt

Bibliographic information is available
on the Lithuanian Integral Library Information System (LIBIS) portal www.ibiblioteka.lt
ISBN 978-609-07-1051-7 (PDF)

© Vilnius University, 2024

CALCIUM-INDUCED HETERODIMERIZATION OF S100A8 WITH S100A1 TRIGGERS AMYLOID FIBRILLATION

Viktorija Karalkevičiūtė¹, Ieva Baronaitė¹, Darius Šulskis¹, Vytautas Smirnovas¹

¹ Sector of Amyloid Research, Institute of Biotechnology, Life Sciences Centre, Vilnius University, Lithuania
viktorija.karalkeviciute@gmc.stud.vu.lt

S100 is a family of calcium-binding proteins, consisting of isoforms with structural similarity but functional diversity [1]. S100 proteins regulate various proteins involved in cellular functions like calcium homeostasis, cell growth, differentiation, cytoskeleton dynamics, and energy metabolism [2]. Several members are known to be important in neurodegeneration by signaling neuroinflammation and forming amyloid fibrils. One of them is S100A9, which is well-studied, but the roles of S100A1 and S100A8 remain relatively unexplored.

S100A1 is predominantly expressed in the brain, skeletal and cardiac muscles [3]. S100A1 interacts with tau, RAGE, and RyR - proteins that participate Alzheimer's disease (AD) cascade [4]. Another family member S100A8 is mostly found in neutrophils and monocytes [5] and plays a role in neurological disease pathology as well. S100A8 homodimers can independently induce neuroinflammation [6] and their overexpression in AD patients leads to activation of microglia [7, 8]. However, it is known that S100A8 can form a heterodimer with S100A9 called calprotectin [9], but interaction with S100A1 is still not investigated. Both S100A1 and S100A8 are expressed in the cerebral cortex as per the Human Protein Atlas (<https://www.proteinatlas.org/>) [10] and share structural similarities [1]. Thus, our main goal was to elucidate their potential complex formation.

To explore the aggregation kinetics of the S100A1/A8 complex, we employed the Thioflavin T Fluorescence Assay, unveiling calcium concentration-dependent amyloid formation. In addition, Atomic Force Microscopy (AFM) was used to visualize the S100A1/A8 fibrils and, Differential Scanning Fluorimetry (DSF) to quantify protein stabilities. In conclusion, our research contributes new findings to the understanding of S100A1 and S100A8 aggregation dynamics, offering valuable insights into their relevance to various diseases.

-
- [1] P. Singh and S. A. Ali, "Multifunctional Role of S100 Protein Family in the Immune System: An Update," *Cells*, vol. 11, no. 15, Jan. 2022.
 - [2] R. Donato et al., "Functions of S100 Proteins," *Curr. Mol. Med.*, vol. 13, no. 1, pp. 24–57, Jan. 2013.
 - [3] N. T. Wright et al., "S100A1: Structure, Function, and Therapeutic Potential," *Curr. Chem. Biol.*, vol. 3, no. 2, pp. 138–145, May 2009.
 - [4] J. S. Cristóvão and C. M. Gomes, "S100 Proteins in Alzheimer's Disease," *Front. Neuroscience*, vol. 13, 2019.
 - [5] K. Sunahori et al., "S100A8/A9 Heterodimer and Proinflammatory Cytokine Production in Rheumatoid Arthritis," *Arthritis Res. Ther.*, vol. 8, no. 3, p. R69, 2006.
 - [6] T. Vogl et al., "Pro-Inflammatory S100A8 and S100A9 Proteins," *Int. J. Mol. Sci.*, vol. 13, no. 3, Mar. 2012.
 - [7] M. Sidoryk-Wegrzynowicz et al., "Astrocytes in Mouse Models of Tauopathies," *Acta Neuropathol. Commun.*, vol. 5, no. 1, p. 89, Nov. 2017.
 - [8] H. L. Weiner and D. Frenkel, "Immunology and Immunotherapy of Alzheimer's Disease," *Nat. Rev. Immunol.*, vol. 6, no. 5, May 2006.
 - [9] I. P. Kornödörfer et al., "Crystal Structure of the Human Calprotectin," *J. Mol. Biol.*, vol. 370, no. 5, pp. 887–898, Jul. 2007.
 - [10] M. Uhlen et al., "Knowledge-based Human Protein Atlas," *Nat. Biotechnol.*, vol. 28, no. 12, pp. 1248–1250, Dec. 2010.