

THE 67TH INTERNATIONAL



OPEN READINGS

CONFERENCE FOR STUDENTS OF PHYSICS AND NATURAL SCIENCES

**BOOK OF
ABSTRACTS**

2024



Vilnius
University

VILNIUS UNIVERSITY PRESS

Editors:

Martynas Keršys
Rimantas Naina
Vincentas Adomaitis
Emilijus Maskvytis

Cover and Interior Design:

Goda Grybauskaitė

Vilnius University Press
9 Saulėtekio Av., III Building, LT-10222 Vilnius
info@leidykla.vu.lt, www.leidykla.vu.lt/en/
www.knygynas.vu.lt, www.journals.vu.lt

Bibliographic information is available
on the Lithuanian Integral Library Information System (LIBIS) portal www.ibiblioteka.lt
ISBN 978-609-07-1051-7 (PDF)

© Vilnius University, 2024

DEVELOPMENT OF THE SCREENING SYSTEM FOR TRANSPOSABLE ELEMENTS

Bartė Žiliukaitė¹, Gytis Druteika¹, Tautvydas Karvelis¹

¹Department of Protein-DNA Interactions, Institute of Biotechnology, Life Sciences Center, Vilnius University, Lithuania
barte.ziliukaite@chgf.stud.vu.lt

Prokaryotic CRISPR-Cas systems are responsible for the adaptive immunity against foreign DNA, which is based on guide RNA-dependent DNA or RNA nuclease activity. Nowadays, these systems are adapted and widely used for genome editing; however, they are inefficient, prone to off-target activity, and require PAM sequences that may greatly limit possible targets in the genome. In addition, gene editing with Cas proteins relies on homologous recombination repair, which is tied to active cell division and, therefore, is not applicable to some cell types [1]. All these limitations of CRISPR-Cas systems raise demand for novel genome editing tools. Recent studies and bioinformatic analysis show that Cas12 nucleases may have evolved from TnpB – proteins encoded by IS200/IS605 family insertion sequences (IS) [2]. IS, as well as plasmids, bacteriophages, and transposons, are mobile genetic elements (MGE) that are abundant in both prokaryotes and eukaryotes and play an important role in the evolution and expression of the host genome [3]. The wide variety and evolutionary proximity of IS to Cas proteins make IS a potential source of novel genome editing tools. This study aimed to develop a screening system for transposable elements. Based on the previously described work [4,5], we constructed vectors, which contain an active transposase and an antibiotic resistance gene-carrying minimal transposable element (mini-Tn). We conducted experiments to select bacterial strains that are most suitable for transposition assay, optimized bacterial conjugation conditions, and determined the best conditions for transposase expression. These results contribute to the development of a robust IS screening system, which will facilitate the detection of new IS that could lead to novel genome editing tools.

-
- [1] Strecker J, Ladha A, Gardner Z, et al. RNA-guided DNA insertion with CRISPR-associated transposases. *Science*. 2019;365(6448):48-53. doi:10.1126/science.aax9181
- [2] Sasnauskas G, Tamulaitiene G, Druteika G, et al. TnpB structure reveals minimal functional core of Cas12 nuclease family. *Nature*. 2023;616(7956):384-389. doi:10.1038/s41586-023-05826-x
- [3] Siguier P, Gourbeyre E, Chandler M. Bacterial insertion sequences: their genomic impact and diversity. *FEMS Microbiol Rev*. 2014;38(5):865-891. doi:10.1111/1574-6976.12067
- [4] Kaczmarek Z, Czarnocki-Cieciura M, Górecka-Minakowska KM, et al. Structural basis of transposon end recognition explains central features of Tn7 transposition systems. *Mol Cell*. 2022;82(14):2618-2632.e7. doi:10.1016/j.molcel.2022.05.005
- [5] Meers C, Le HC, Pesari SR, et al. Transposon-encoded nucleases use guide RNAs to promote their selfish spread. *Nature*. 2023;622(7984):863-871. doi:10.1038/s41586-023-06597-1