

THE 67TH INTERNATIONAL



OPEN READINGS

CONFERENCE FOR STUDENTS OF PHYSICS AND NATURAL SCIENCES

**BOOK OF
ABSTRACTS**

2024



Vilnius
University

VILNIUS UNIVERSITY PRESS

Editors:

Martynas Keršys
Rimantas Naina
Vincentas Adomaitis
Emilijus Maskvytis

Cover and Interior Design:

Goda Grybauskaitė

Vilnius University Press
9 Saulėtekio Av., III Building, LT-10222 Vilnius
info@leidykla.vu.lt, www.leidykla.vu.lt/en/
www.knygynas.vu.lt, www.journals.vu.lt

Bibliographic information is available
on the Lithuanian Integral Library Information System (LIBIS) portal www.ibiblioteka.lt
ISBN 978-609-07-1051-7 (PDF)

© Vilnius University, 2024

INTERACTION OF ALKYLPHOSPHOLIPIDS WITH TETHERED BILAYER LIPID MEMBRANES

Rūta Bagdonaitė¹, Artūras Polita¹

¹Institute of Biochemistry, Life Sciences Center, Vilnius University, Saulėtekio av. 7, Vilnius, LT-10257, Lithuania
ruta.bagdonaitė@bchi.stud.vu.lt

Synthetic alkylphospholipids (APLs) are stable analogs of lysophosphatidylcholine, which are being investigated as potential antitumor agents. APLs exert their activities by acting on cell membranes [1], however understanding of their interaction with lipid bilayers is still lacking. In this work we investigated three APLs differing in hydrocarbon chain length and headgroup composition – miltefosine (hexadecylphosphocholine), edelfosine (1-O-octadecyl-2-O-methyl-sn-glycero-phosphocholine) and perifosine (Octadecyl-(1,1-dimethyl-4-piperidyl) phosphate). We used tethered bilayer lipid membranes (tBLMs) formed on self-assembled monolayer modified gold surface as an artificial membrane model [2] and measured its interaction with APLs. We employed electrochemical impedance spectroscopy (EIS) to assess defect formation and fluorescence lifetime imaging microscopy (FLIM) to observe membrane structural changes through microviscosity measurements. Incubation of APLs with tBLMs composed of dopc and cholesterol induced dielectric membrane damage in a time and concentration-dependent manner, with miltefosine being less damaging compared to edelfosine and perifosine at the same concentrations. APLs' induced damage was also demonstrated to be partially reversible at moderate concentrations, reaching a point in EIS spectra after which it shifts back towards lower defect densities. FLIM measurements showed that besides defect formation APLs also induce changes in membrane organization and form mobile micron-scale cholesterol-rich domains.

[1] W.J.vanBlitterswijk, M.Verheij, AnticancerMechanismsAndClinicalApplicationOfAlkylphospholipids (Biochim. Biophys. Acta - Mol. Cell Biol. Lipids, 2013)
[2] T.Penkaukas, G.Preta, BiologicalApplicationsOfTetheredBilayerLipidMembranes (Biochimie, 2019)