

THE 67TH INTERNATIONAL



OPEN READINGS

CONFERENCE FOR STUDENTS OF PHYSICS AND NATURAL SCIENCES

**BOOK OF
ABSTRACTS**

2024



Vilnius
University

VILNIUS UNIVERSITY PRESS

Editors:

Martynas Keršys
Rimantas Naina
Vincentas Adomaitis
Emilijus Maskvytis

Cover and Interior Design:

Goda Grybauskaitė

Vilnius University Press
9 Saulėtekio Av., III Building, LT-10222 Vilnius
info@leidykla.vu.lt, www.leidykla.vu.lt/en/
www.knygynas.vu.lt, www.journals.vu.lt

Bibliographic information is available
on the Lithuanian Integral Library Information System (LIBIS) portal www.ibiblioteka.lt
ISBN 978-609-07-1051-7 (PDF)

© Vilnius University, 2024

PHOTOSENSITIZER TPPS₄ AGGREGATION AND AGGREGATE TYPE SPECTROSCOPIC STUDIES IN NEUTRAL AND HIGHLY ACIDIC PH ENVIRONMENTS

Greta Tamoliūnaitė^{1,2}, Vilius Poderys¹, Ričardas Rotomskis^{1,3}

¹Biomedical Physics Laboratory of National Cancer Institute, Vilnius, Lithuania

²Faculty of Chemistry and Geosciences, Vilnius University, Vilnius, Lithuania

³Faculty of Physics, Vilnius University, Vilnius, Lithuania

greta.tamoliunaite@chgf.stud.vu.lt

Cancer development in patients is at an utmost high. A progressive cancer treatment is photodynamic therapy (PDT), that utilizes photosensitizer, light and molecular oxygen to initiate selective cancer cells death by singlet oxygen, produced by the long lived triplet state of the photosensitizer.

The absorption spectra changes of 5,10,15,20-Tetrakis(4-sulfonatophenyl)porphyrin (TPPS₄), a photosensitizer proposed for PDT, were measured with Varian Cary 50 spectrophotometer in spectral region 350 nm - 750 nm, in deionized water at different pH (achieved by adding into the aqueous solution HCl (1M) and NaOH(1M)) and different concentrations ($1 \times 10^{-2} \text{M}$ - $1 \times 10^{-6} \text{M}$). It is well known that in acidic environment TPPS₄ molecules can form J-aggregates, that exhibit characteristic absorption bands at 490 nm and 709 nm. These aggregates are formed because negatively charged SO₃⁻ group interacts with the positively charged core of protonated porphyrin ring [1]. At neutral pH, the TPPS₄ core is neutral, therefore J-aggregates are unable to form, but spectral changes were detected at high TPPS₄ concentrations. A significant Red-shift of Q₁ and Q₄ absorption bands and the intensity redistribution of central Q₂ and Q₃ bands show a new type of aggregate, possibly face-to-face, formation.

At very acidic pH=-1 all SO₃⁻ are protonated, therefore there are no electrostatic interactions of the positively charged core of porphyrin ring and neutral SO₃⁻, but, with increasing TPPS₄ concentration, absorption spectrum changes were detected, which are slightly similar to J-aggregate formation. Our studies have shown that, regardless of the pH of the solution, TPPS₄ form aggregates at high concentrations, that are characterized by a lower quantum efficiency of singlet oxygen generation, therefore it is necessary to control the concentration of TPPS-based photodrugs, injected into the patient, because a too high dose of a photosensitizer can lead to aggregation and, as a result, the decrease in singlet oxygen generation, which will determine the lower efficiency of PDT.




Fig. 1. TPPS₄ aggregation and aggregate types in different pH environments.

[1] Rotomskis R, Augulis R, Snitka V, Valiokas R, Liedberg B, Hierarchical Structure of TPPS₄ J-Aggregates on Substrate Revealed by Atomic Force Microscopy, J. Phys. Chem. B 2004 108(9), 2833-2838. DOI: 10.1021/jp036128v