

SCIENTIFIC REPORTS



OPEN

Inhibiting mevalonate pathway enzymes increases stromal cell resilience to a cholesterol-dependent cytotoxin

Sholeem Griffin¹, Giulio Preta^{1,2} & Iain Martin Sheldon¹

Animal health depends on the ability of immune cells to kill invading pathogens, and on the resilience of tissues to tolerate the presence of pathogens. *Trueperella pyogenes* causes tissue pathology in many mammals by secreting a cholesterol-dependent cytotoxin, pyolysin (PLO), which targets stromal cells. Cellular cholesterol is derived from squalene, which is synthesized via the mevalonate pathway enzymes, including HMGCR, FDPS and FDFT1. The present study tested the hypothesis that inhibiting enzymes in the mevalonate pathway to reduce cellular cholesterol increases the resilience of stromal cells to PLO. We first verified that depleting cellular cholesterol with methyl- β -cyclodextrin increased the resilience of stromal cells to PLO. We then used siRNA to deplete mevalonate pathway enzyme gene expression, and used pharmaceutical inhibitors, atorvastatin, alendronate or zaragozic acid to inhibit the activity of HMGCR, FDPS and FDFT1, respectively. These approaches successfully reduced cellular cholesterol abundance, but mevalonate pathway enzymes did not affect cellular resilience equally. Inhibiting FDFT1 was most effective, with zaragozic acid reducing the impact of PLO on cell viability. The present study provides evidence that inhibiting FDFT1 increases stromal cell resilience to a cholesterol-dependent cytotoxin.

Health and the ability to counter pathogenic microbes depends on an organism's immunity and resilience^{1,2}. Immunity, or resistance, is the ability to reduce the pathogen burden by killing infecting microbes. Resilience, or tolerance, is the ability to limit the impact of pathogens on health, by tolerating a given microbial burden²⁻⁴. The ability to tolerate pathogenic bacteria largely depends on the resilience of the host's tissue cells to damaging bacterial virulence factors. Cholesterol-dependent cytotoxins are a common virulence factor secreted by pathogenic bacteria, and they have a high affinity for cholesterol in the plasma membrane of mammalian cells, where they form 30–50 nm diameter pores^{5,6}. These pores allow leakage of molecules across the plasma membrane, resulting in cell death and tissue damage. Most cellular cholesterol is located in the plasma membrane of animal cells, where it constitutes almost half of the lipid molecules⁷. Cholesterol synthesis depends on the production of squalene by the mevalonate pathway⁸. The mevalonate pathway enzymes are common drug targets, used to limit cellular cholesterol synthesis for the control of hypercholesterolemia⁹. Here we explored whether inhibiting the mevalonate pathway to reduce cellular cholesterol in tissue cells could also increase their resilience to cholesterol-dependent cytotoxins.

Trueperella pyogenes is a Gram-positive bacterium found on the skin and mucosa of many animals, and it causes pathology in several tissues, including mucosa, liver, and skin¹⁰. Postpartum uterine disease in cattle is the most economically important disease associated with *T. pyogenes* infection, typically affecting 20 to 40% of animals after parturition¹¹⁻¹³. Uterine disease costs the USA and EU dairy industry about \$2 billion/year in lost production, infertility, and treatment costs¹¹. The presence of *T. pyogenes* correlates with the severity of endometrial pathology, the extent of the subsequent infertility, and infusion of *T. pyogenes* recapitulates the disease¹⁴⁻¹⁶. *Trueperella pyogenes* causes inflammation and damage of the stromal compartment of the endometrium, once the surface epithelium is breached during parturition. The main virulence factor secreted by *T. pyogenes* is a cholesterol-dependent cytotoxin, pyolysin (PLO)¹⁰. Bovine endometrial stromal cells are highly sensitive to

¹Swansea University Medical School, Swansea University, Swansea, SA2 8PP, United Kingdom. ²Institute of Biochemistry, Vilnius University, Sauletekio 7, Vilnius LT-10257, Lithuania. Correspondence and requests for materials should be addressed to I.M.S. (email: i.m.sheldon@swansea.ac.uk)




Figure 1. The mevalonate pathway leads to cholesterol synthesis. Acetoacetyl CoA and acetyl CoA are converted to squalene, which is subsequently converted to cholesterol. Important enzymes in the pathway include 3-hydroxy-3-methyl-glutaryl-coenzyme A reductase (HMGCR), farnesyl pyrophosphate synthase (FDPS), and farnesyl diphosphate farnesyltransferase 1 (FDFT1, also commonly called squalene synthase). Each of these enzymes can be inhibited by statins, bisphosphonates, and zaragozic acid, respectively; cholesterol can be depleted using methyl- β -cyclodextrin (MBCD).

cytolysis caused by PLO¹⁶. Whilst antimicrobial therapy limits the severity of disease, there are no treatments to increase the ability of tissues to tolerate *T. pyogenes*. The gap in knowledge is how to manipulate cellular resilience to help tissues tolerate pathogens.

There are three pools of cholesterol in the plasma membrane of cells: an essential pool, a sphingomyelin sequestered pool, and a labile pool of cholesterol¹⁷. The labile cholesterol can be depleted *in vitro* using methyl- β -cyclodextrin (MBCD), which is a cyclic oligosaccharide that binds cholesterol^{18,19}. Reducing cellular cholesterol with MBCD increases stromal cell resilience to PLO^{16,20}. Cellular cholesterol abundance is highly regulated and depends on the balance amongst cholesterol synthesis, cholesterol efflux, and cholesterol uptake from low-density lipoproteins^{9,21}. Cellular cholesterol synthesis uses a series of enzymes, with the mevalonate pathway providing the rate-limiting process⁸. The mevalonate pathway initially condenses two acetyl-CoA molecules to form acetoacetyl-CoA, which are converted to 3-hydroxy-3-methyl-glutaryl-CoA (HMG-CoA) by HMG-CoA synthase, before HMG-CoA reductase (HMGCR, EC 1.1.1.34) yields mevalonate (Fig. 1). A series of enzymes, ending with farnesyl pyrophosphate synthase (FDPS, EC 2.5.1.10), then convert mevalonate to farnesyl pyrophosphate. Farnesyl pyrophosphate is a substrate for several enzymes, but the most important for cholesterol synthesis is farnesyl diphosphate farnesyltransferase 1 (FDFT1, EC 2.5.1.21), which is also called squalene synthase^{8,22}.

We reasoned that depleting cellular cholesterol might increase the resilience of cells to PLO. The most common drugs used to reduce hypercholesterolemia regulate the mevalonate pathway include: statins, such as atorvastatin, which inhibit HMGCR; nitrogen-containing bisphosphonates, such as alendronate, which inhibit FDPS; and, zaragozic acid A, derived from fungi, which inhibit FDFT1^{9,23–25}. The present study tested the hypothesis that inhibiting enzymes in the mevalonate pathway to reduce cellular cholesterol increases the resilience of stromal cells to PLO.

Results

PLO causes cytolysis. To explore the resilience of stromal cells to cholesterol-dependent cytolysins, we chose to use primary bovine endometrial stromal cells (BESC) because they are the main target of *T. pyogenes*¹⁶. In addition, to examine whether cellular resilience mechanisms are conserved across species, we used telomerase-immortalized human endometrial stromal cells (HESC)²⁶. Using pure populations of stromal cells removes potential confounding effects of immune cells, and helps focus on mechanisms that impact tissue cell resilience, rather than immunity^{2,4}. Recombinant pyolysin protein (PLO) was generated and purity was confirmed using SDS-PAGE (Supplementary Fig. S1), as described previously^{16,27,28}. The specific activity of PLO was 628,338 HU/mg protein, as determined using a hemolysis assay, and there was very little endotoxin contamination (1.5 EU/mg protein).

The BESC and HESC were cultured in 24-well plates until 70% confluent and then challenged directly with PLO because, unlike most cholesterol-dependent cytolysins, PLO is spontaneously active without the addition of thiol reducing agents^{6,10}. To limit the potential effects of cellular uptake of cholesterol or serum binding to PLO, serum-free media were used as described previously^{20,29}. After challenging the cells for 2 h with a range of concentrations of PLO, cell viability was assessed by the mitochondria-dependent reduction of 3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide (MTT) to formazan in viable cells³⁰. In this colorimetric assay we defined 100% viability as the OD₅₇₀ measurement for cells in control medium, and 0% viability as the OD₅₇₀ for cells lysed using Triton X-100. We assumed that a reduction in OD₅₇₀ was a reflection of cytotoxicity, as described previously for experiments using cholesterol-dependent cytolysins^{16,20,31}. The correlation between MTT OD₅₇₀ measurements and the number of live cells was confirmed previously using trypan blue exclusion and counting the number of live cells using a hemocytometer¹⁶. The percentage cell viability was calculated as the OD₅₇₀ of PLO-treated cells relative to OD₅₇₀ values for cells in control medium. The viability of BESC and HESC decreased in a PLO concentration-dependent manner (Fig. 2A,B). For subsequent experiments, we selected a 2 h challenge with 100 HU/ml PLO for BESC and 200/ml HU PLO for HESC.

We confirmed the utility of challenging BESC and HESC with 100 and 200 HU/ml PLO, respectively, by measuring the leakage of lactate dehydrogenase (LDH) from the cytosol into the supernatant, which is a proxy for the formation of pores in plasma membranes by cholesterol-dependent cytolysins³². There was increased abundance of LDH in the supernatants of both BESC and HESC challenged with PLO, compared with control (Fig. 2C,D). To verify that PLO activity is cholesterol-dependent, we also depleted cellular cholesterol by treating cells with MBCD, which binds to cholesterol and prevents the formation of pores by cholesterol-dependent cytolysins^{16,18,19,33}. When BESC or HESC were treated with MBCD for 24 h, and then challenged with PLO for 2 h, there was no significant increase in supernatant LDH concentrations, compared with control (Fig. 2C,D).

As a control, to verify the cytolytic activity was caused by PLO, we used a mutant PLO protein, which has a protein-stabilizing disulphide bridge (DS-PLO), that binds to plasma membranes but does not form pores²⁸. Challenging cells with PLO caused a marked reduction in viability of both cell types, whereas an equivalent amount of DS-PLO did not cause cytolysis (Fig. 2E,F). Taken together, these data provide evidence that BESC and HESC are sensitive to the cholesterol-dependent cytolysin PLO.

RNA interference of *FDFT1* increases cellular resilience to PLO. We used RNA interference and pharmaceutical inhibitors to explore the role of the mevalonate pathway in cellular resilience to PLO. In the first approach, BESC and HESC were transfected with short interfering RNA (siRNA) sequences targeting key genes in the mevalonate pathway *HMGCR*, *FDPS* and *FDFT1* (Table 1). The transfection and RNA interference technique in serum-free media did not significantly affect cell viability or cell permeability, as determined by MTT and LDH assays, respectively (Supplementary Fig. S2). However, mRNA expression was reduced by 48%, 90% and 69% with siRNA targeting *HMGCR*, *FDPS* and *FDFT1*, respectively, in BESC (Fig. 3A), and by 91%, 89% and 91%, respectively, in HESC (Fig. 3B). Cellular concentrations of cholesterol were also measured to quantify the effectiveness of the siRNA. Each of the siRNA significantly reduced the concentrations of cholesterol in BESC and HESC, and the concentrations were similar to cells treated with MBCD to deplete cholesterol (Fig. 3C,D).

The BESC and HESC were next transfected with siRNA for 48 h and then challenged with PLO for 2 h. Cells transfected with control scramble siRNA had a reduction in cell viability when challenged with PLO, as determined by MTT assay (Fig. 3E,F), and increased accumulation of LDH in supernatants (Fig. 3G,H). Using MBCD to deplete cellular cholesterol as a positive control reduced the impact of PLO in BESC and HESC. As might be expected with primary cells, the effect of siRNA in BESC was somewhat variable. Although siRNA targeting *FDPS* did not significantly rescue BESC when challenged with PLO, depleting *HMGCR* and *FDFT1* increased cell resilience to PLO (Fig. 3E). However, depletion of *HMGCR*, *FDPS* and *FDFT1* did not significantly change the BESC leakage of LDH when challenged with PLO (Fig. 3G). In HESC, only siRNA depletion of *FDFT1* significantly increased cell resilience to PLO and reduced leakage of LDH (Fig. 3G,H). Together these data provide evidence that siRNA targeting mevalonate pathway enzymes reduces cellular cholesterol, but only depletion of specific enzymes, such as *FDFT1*, enhances stromal cell resilience to PLO.

Inhibitors of mevalonate pathway enzymes enhance cellular resilience to PLO. For our second approach to explore cellular resilience to PLO, we selected atorvastatin to inhibit *HMGCR*, the bisphosphonate alendronate to inhibit *FDPS*, and zaragozic acid to inhibit *FDFT1*. In order to determine if the inhibitors might reduce cell viability *per se*, BESC and HESC were incubated with the manufacturer's recommended concentrations of 1 μM atorvastatin, 10 μM alendronate and 10 μM zaragozic acid for 48 h in serum free media, prior to evaluating cell viability using the MTT assay. Apart from atorvastatin in HESC, the inhibitors did not significantly reduce BESC or HESC viability (Fig. 4A,B). However, each of the inhibitors reduced cellular cholesterol in BESC and HESC, compared with control (Fig. 4C,D).




Figure 2. The effect of PLO on stromal cells. (A) BESC isolated from 3 animals, and (B) HESC from 3 independent passages were challenged for 2 h with control medium (■), or media containing the indicated concentrations of PLO (■), and cell viability evaluated by MTT assay. Data are presented as mean (SEM), and analyzed by one-way ANOVA with Dunnett's multiple comparison post-hoc test; values differ from control, ** $P < 0.01$, *** $P < 0.001$. (C) BESC isolated from 5 animals, and (D) HESC from 3 independent passages were grown in control serum-free medium or medium containing methyl- β -cyclodextrin for 24 h to deplete cholesterol, and then challenged for 2 h with control medium (■), or media containing PLO (■) using 100 HU/ml for BESC and 200 HU/ml for HESC, and leakage of LDH was measured in cell supernatants. Data are presented as mean (SEM), and analyzed by one-way ANOVA with Dunnett's multiple comparison post-hoc test; values differ from control, ** $P < 0.01$. (E) BESC isolated from 3 animals, and (F) HESC from 3 independent passages, were challenged for 24 h with control serum-free medium (■), or media containing PLO (■) or DS-PLO (DS, ■), and cell viability evaluated by MTT assay. Data are presented as mean (SEM) percent viability of control, and analyzed by one-way ANOVA with Dunnett's multiple comparison post-hoc test; values differ from control, ** $P < 0.01$, *** $P < 0.001$.

One concern with the inhibitors was that observations might be misleading if the inhibitors could bind PLO. To test this, each inhibitor was incubated with PLO for 1 h prior to addition of horse red blood cells and measurement of hemolysis. Cholesterol was used as a positive control to bind PLO and reduce hemolysis. However, the mevalonate pathway inhibitors or MBCD did not significantly alter the hemolytic effect of PLO (Fig. 4E).

Having established the mevalonate pathway inhibitors could reduce cellular cholesterol, BESC and HESC were incubated for 48 h with a range of concentrations of atorvastatin, alendronate or zaragozic acid, prior to challenge with PLO for 2 h. In BESC, there was no significant effect of atorvastatin, but alendronate and particularly zaragozic acid increased the resilience of cells to PLO (Fig. 5A). In HESC, there was no significant effect of alendronate, but atorvastatin and particularly zaragozic acid increased the resilience of cells to PLO (Fig. 5B). To

Species	Gene	Direction	Sequence (5' → 3')
<i>Bos taurus</i>	HMGCR	Sense	CAGCAUGGAUUAUGAACAAUU
		Antisense	UUGUUCAAUAUCCAUGCUG
<i>Bos taurus</i>	FDPS	Sense	GCACAGACAUCCAGGACAAUU
		Antisense	UUGUCCUGGAUGUCUGUGCUU
<i>Bos taurus</i>	FDFT1	Sense	GCGAGAAGGGAGAGAGUUUUU
		Antisense	AAACUCUCUCCCUUCUCGC
<i>Homo sapiens</i>	HMGCR	Sense	GGAUAAACCAGAGAAAGAAUU
		Antisense	UUUCUUUCUCGGUUUAUCC
<i>Homo sapiens</i>	FDPS	Sense	GCAGAAGGAGGCUGAGAAUU
		Antisense	UUUCUCAGCCUCCUUCUGC
<i>Homo sapiens</i>	FDFT1	Sense	GCAAGGAGGAAGAGAGUUUUU
		Antisense	GAACUCUCUCCCUUCUGC

Table 1. siRNA sequence for target gene knockdown.

further examine the effect of zaragozic acid, BESC were treated for 24 h or 48 h with 10 μ M zaragozic acid prior to challenge with PLO, and cell viability monitored hourly using the Alamar Blue assay³⁴, which depends on the cellular enzymatic reduction of resazurin to resorufin (Fig. 5C,D). Cellular resilience to PLO was increased by 24 h or 48 h treatment with zaragozic acid (ANOVA, $P < 0.05$, and $P < 0.001$, respectively).

To further explore the impact of mevalonate pathway inhibitors, BESC and HESC were incubated with control serum-free media, or media containing 1 μ M atorvastatin, 10 μ M alendronate, 10 μ M zaragozic acid, or 1 mM methyl- β -cyclodextrin (MBCD) for 48 h, and then challenged with control medium or PLO for 2 h. The inhibitors *per se* did not significantly affect cell viability or the release of LDH in control medium (all values within 10% of control). However, atorvastatin and zaragozic acid reduced the impact of PLO on cell viability, as determined by MTT assay, in BESC (Fig. 6A) and HESC (Fig. 6B). There was also a significant reduction in the leakage of LDH for atorvastatin in BESC, and zaragozic acid in HESC (Fig. 6C,D). To determine whether the protection against PLO by mevalonate pathway inhibitors affected the number of cells, total cellular DNA was measured using the CyQuant assay. Total cellular DNA remained unchanged after challenge with control media or PLO (Fig. 6E,F), indicating that changes in viability were independent of the number of cells. Taken together, these data support a concept that the resilience of stromal cells to PLO is increased by inhibiting specific mevalonate pathway enzymes, particularly FDFT1.

Discussion

We found that inhibiting enzymes in the mevalonate pathway reduced cholesterol abundance in both primary bovine endometrial stromal cells (BESC) and telomerase-immortalized human endometrial stromal cells (HESC). However, surprisingly, inhibiting each mevalonate pathway enzyme did not equally affect cellular resilience to PLO. Inhibiting FDFT1 most consistently increased stromal cell resilience to PLO. We suggest that manipulating the mevalonate pathway may help tissue cells tolerate bacterial infections. We also speculate that topical application of FDFT1 inhibitors could be used therapeutically in the uterus.

Pyolysin is considered to be the only cytolysin secreted by *T. pyogenes*, as determined using an anti-PLO antibody or deletion of the *ply* gene^{10,16,35,36}. To examine mechanisms of cellular resilience, the present study used stromal cells because it is important to use tissue cells that could influence the ability of tissues to tolerate pathogens^{2,4}. As expected, BESC were sensitive to recombinant PLO protein^{16,20}. However, PLO also reduced cellular viability and increased the leakage of LDH into supernatants of HESC. The sensitivity of HESC to PLO was unexpected as *T. pyogenes* only rarely causes disease in humans³⁷. However, the potency of PLO across species and tissues may help explain the wide diversity of species and diseases that are associated with *T. pyogenes* infection. Although a specific cell surface receptor cannot be ruled out, the ability of exogenous cholesterol to inhibit hemolysis in the present study and the sensitivity of HESC to PLO, provides evidence that cholesterol is likely the main target of PLO.

Plasma membranes of cultured fibroblast cells from human foreskin contain half their cellular phospholipid and 90% of their cholesterol and sphingomyelin⁷. In the plasma membrane, cholesterol is distributed amongst an essential pool, a sphingomyelin sequestered pool, and a labile pool of cholesterol that binds to cholesterol-dependent cytolysins¹⁷. We assume that this labile pool of cholesterol is most likely depleted when cells are treated with MBCD¹⁸. In the present study, MBCD consistently reduced cellular cholesterol abundance in BESC and HESC, and this was associated with increased resilience to PLO. Although MBCD has not been used with HESC previously, these observations are consistent with findings for BESC and other cells, and with a range of cholesterol-dependent cytolysins^{6,16,20,33}. Cellular cholesterol abundance depends on cholesterol uptake from LDL, cholesterol efflux, and cholesterol synthesis via the mevalonate pathway^{8,9}. Cholesterol homeostasis is highly regulated in cells and HMGCR is usually considered to be the rate-limiting enzyme in cholesterol synthesis, although this may not apply to all cells^{21,22}. The importance of the mevalonate pathway for cellular cholesterol was verified in the present study, using siRNA or inhibitors to target HMGCR, FDPS, and FDFT1 in endometrial stromal cells, which consistently reduced cellular cholesterol by about half; similar to the effect of MBCD.

The surprising finding in the present study was that despite siRNA and inhibitors reducing cellular cholesterol, the effect on cellular resilience varied. Depleting *FDFT1* mRNA expression or inhibiting *FDFT1* activity




Figure 3. RNA interference of mevalonate pathway enzymes. **(A)** BESC isolated from 3 animals, and **(B)** HESC from 3 independent passages, were transfected with scramble siRNA or siRNA targeting *HMGCR*, *FDPS* or *FDFT1*. The mRNA expression of each cognate gene was measured by qPCR, and data presented as mean (SEM) relative to two reference genes. Data were analyzed by Student's t-test; values differ from scramble, ** $P < 0.01$, *** $P < 0.001$. BESC isolated from 3 animals **(C,E,G)** or HESC from 3 independent passages **(D,F,H)** were incubated for 48 h in serum-free control medium, or media containing scramble siRNA or siRNA targeting *HMGCR*, *FDPS* or *FDFT1*, or cultured with methyl- β -cyclodextrin (MBCD) as a positive control. Cellular cholesterol **(C,D)** was measured and normalized to phospholipid concentrations, and data presented as arbitrary units (AU) with each dot representing an independent animal or cell passage, and a horizontal line indicating the mean. Cells were challenged with control media (■) or media containing PLO (■) using 100 HU/ml for BESC and 200 HU/ml for HESC, for 2 h. Cell viability was quantified by MTT assay **(E,F)**, and supernatants were collected to measure LDH **(G,H)**. Data are presented as mean (SEM), and were analyzed by one-way ANOVA and Dunnett's multiple comparison post-hoc test; values differ from scramble challenged with PLO, * $P < 0.05$, ** $P < 0.01$, *** $P < 0.001$.




Figure 4. Inhibition of mevalonate pathway enzymes. (A,C) BESC from 3 animals and (B,D) HESC from 3 independent passages were incubated in control serum-free medium, or media containing 1 μ M atorvastatin (AT), 10 μ M alendronate (AL), 10 μ M zarogozic acid (ZA) or methyl- β -cyclodextrin (MBCD) for 48 h. (A,B) Cell viability was assessed by MTT assay. Data are expressed as mean (SEM), and were analyzed by one-way ANOVA and Dunnett's multiple comparison post hoc test; values differ from control, * $P < 0.05$. (C,D) Cellular cholesterol was measured and normalized to phospholipid concentrations, and data presented as arbitrary units (AU) with each dot representing an independent animal or cell passage, and a horizontal line indicating the mean. Data were analyzed by one-way ANOVA and Dunnett's multiple comparison post hoc test; values differ from control, * $P < 0.05$, ** $P < 0.01$ and *** $P < 0.001$. (E) The inhibitors or vehicle were incubated with 100 HU/ml PLO for 1 h in DPBS, using 1 mM cholesterol (CHO) as a positive control and DPBS as a negative control (Vehicle), and then incubated with 0.5% (v/v) horse red blood cells for 1 h at 37°C to assess hemolysis. Data are expressed as mean (SEM) relative hemolysis of three independent experiments.




Figure 5. Mevalonate pathway inhibitors modulate the impact of PLO on cell viability. (A) BESC isolated from 4 animals, or (B) HESC from 3 independent passages were incubated for 48 h with control serum-free medium, or media containing the indicated concentrations of atorvastatin, alendronate or zaragozic acid, and then challenged with control media (■) or media containing PLO (■) using 100 HU/ml for BESC and 200 HU/ml for HESC, for 2 h. Cell viability was determined by MTT assay. Data are expressed as mean (SEM), and were analyzed by one-way ANOVA with Dunnett's multiple comparison post hoc test; values differ from PLO treatment with no inhibitor, * $P < 0.05$, ** $P < 0.01$, *** $P < 0.001$. In independent experiments, BESC isolated from 4 animals were incubated for (C) 24 h or (D) 48 h in serum-free media containing vehicle, MBCD or 10 μM zaragozic acid. Cells were then challenged with 100 HU/ml PLO, and cell viability was monitored every hour using the Alamar Blue assay. Data are expressed as mean (SEM) percentage viability of control.

with zaragozic acid most effectively increased cellular resilience to PLO in BESC and HESC. Thus, factors other than the abundance of cholesterol influenced the ability of cells to tolerate PLO. One possible explanation is that off-target effects of the inhibitors could affect cell resilience, although this seems less likely for siRNA. Another possibility is that while inhibition of each enzyme reduces cholesterol synthesis, the abundance of substrates and products will likely vary for each enzyme. For example, inhibiting FDFT1 may lead to increased farnesyl pyrophosphate, whereas inhibiting HMGCR would likely reduce farnesyl pyrophosphate concentrations. In addition, there may be wider impacts on the pathways that regulate cholesterol homeostasis, such as sterol regulatory




Figure 6. Mevalonate pathway inhibitors and cellular resilience to PLO. (A,C,E) BESC from 3 animals, and (B,D,F) HESC from 3 independent passages were incubated with control serum-free media (0), or media containing 1 μ M atorvastatin (AT), 10 μ M alendronate (AL), 10 μ M zaragozic acid (ZA) or methyl- β -cyclodextrin (MBCD) for 48 h, and then challenged with control media (■) or media containing PLO (■) using 100 HU/ml for BESC and 200 HU/ml for HESC, for 2 h. Cell viability was evaluated by MTT assay (A, B), supernatants were collected for LDH assay (C,D), and total cell DNA was determined by CyQuant assay (E,F). Data are expressed as mean (SEM), and were analyzed by one-way ANOVA with Dunnett's multiple comparison post hoc test; values differ from PLO treatment with no inhibitor, * $P < 0.05$, ** $P < 0.01$ and *** $P < 0.001$; ND, below limits of detection.

element binding proteins²¹. Indeed, activation of lipogenic genes by sterol regulatory element binding protein 1 promoted cell survival after toxin attack³⁸.

A limitation of the present study was that BESC show some variation in their susceptibility to PLO, as determined by MTT assay or LDH leakage, and the effects of modulating the mevalonate pathway. Isolation and culture of primary bovine endometrial stromal cells was performed as described previously^{39,40}. Some of the variance may reflect differences amongst the animals from which the cells were isolated, and biological variation is a consistent feature of primary endometrial stromal cells function determined by MTT, LDH and cholesterol assays^{16,20,39}. In addition, as BESC and HESC continue to grow during experiments, the OD₅₇₀ values from the MTT assay also increase when there is a longer duration of the experiment. To provide additional evidence for our claims, future work might use more animals to isolate cells, a wider range of measurements of cell resilience, and use *in vivo* experiments.

The ability to increase tissue cell resilience so that tissues can tolerate pathogens is important for health^{2,4}. Several cellular pathways increase the resilience of cells to cholesterol-dependent cytotoxins, including mitogen-activated protein kinases, the unfolded protein response, autophagy, calcium ion flux, and caspase-1^{38,41–44}. Unfortunately, manipulating these pathways to protect against cholesterol-dependent cytotoxins *in vivo* may be challenging because inhibitors of these pathways are often toxic. However, statins are widely used *in vivo*, and simvastatin enhanced cellular resistance against the cholesterol-dependent cytotoxin pneumolysin in human endothelial cells⁴⁵. Similarly, simvastatin or siRNA targeting HMGCR protected human airway epithelial cells against pneumolysin²⁹. Interestingly, and in accord with our findings, the protective mechanism of simvastatin was not solely mediated by reducing cellular cholesterol²⁹. Furthermore, Statt *et al.* (2015) also found that an inhibitor of FDFT1 or siRNA depleting *FDFT1* mRNA reduced the impact of pneumolysin on cells²⁹. In the present study, zaragozic acid increased the resilience of both BESC and HESC to PLO, and was not toxic to cells at the concentrations we used.

Interestingly, zaragozic acid also reduces BESC inflammatory responses to endotoxin from Gram-negative bacteria⁴⁶; and, reduces serum cholesterol concentrations *in vivo*²⁵. Thus, zaragozic acid might be worthy of further investigation to increase tissue tolerance to bacteria. One therapeutic advantage with uterine infections is that treatments are often topical applications, via intrauterine infusion, which provides high concentration of the pharmaceutical molecule with less risk of toxicity¹¹. We suggest that intrauterine infusion of inhibitors targeting mevalonate pathway enzymes could be used to limit the severity of uterine disease caused by *T. pyogenes* in post-partum animals. An alternative approach would be to infuse MBCD or other cyclodextrins that deplete cholesterol. However, cyclodextrins can also act as cholesterol donors, with the danger of toxicity¹⁹.

In conclusion, the present study provides evidence that manipulating the mevalonate pathway increases the resilience of cells to PLO, to help tissues tolerate bacterial infections. We speculate that inhibitors of the mevalonate pathway could be used to increase cellular resilience to cholesterol-dependent cytotoxins, and limit the severity of disease.

Methods

Pyolysin. The PLO plasmid (pGS59) was a generous gift from Prof BH Jost (University of Arizona, USA), and the DS-PLO mutant plasmid (R219C/G85C) was a kind gift from Prof M Palmer (University of Waterloo, Canada). Proteins were generated as described previously^{16,27,28}. The abundance of proteins was measured by DC assay, and their purity evaluated using SDS-PAGE and Coomassie blue staining, as described previously¹⁶.

A hemolysis assay was used to determine the activity of PLO, as described previously¹⁶. Optical density (OD₄₅₀) was measured using a microplate reader (POLARstar Omega; BMG Labtech, Offenburg, Germany). Hemolytic units were mathematically determined by 4-parameter modelling using the Solver function in Microsoft Excel. Endotoxin concentrations in stock solutions of PLO were all <1 EU/ml, as determined by LAL assay (LAL endotoxin quantitation kit; Thermo Scientific, Hertfordshire, UK). To examine the potential binding to PLO, 1 μM atorvastatin, 10 μM alendronate, 10 μM zaragozic acid, 1 mM methyl-β-cyclodextrin, and 1 mM cholesterol (all Sigma) were incubated with 100 HU/ml PLO for 1 h, prior to conducting the hemolysis assay.

Ethical statement. No live animal experiments were conducted. Uteri were collected from cattle after they were slaughtered and processed as part of the normal work of an abattoir.

Cell culture. Isolation and culture of primary bovine endometrial stromal cells was performed as described previously^{39,40}. Briefly, uteri with no gross evidence of genital disease or microbial infections were collected from cattle after they were slaughtered and processed as part of the normal work of an abattoir. No live animal experiments were used. Stromal cells were isolated by enzymatic digestion of the endometrium, sieving the cell suspension through a 40-μm mesh, and differential adhesion to cell culture plates^{39,40}. Stromal cells were maintained in BESC complete culture medium comprising RPMI-1640 (Gibco, Gaithersburg, MD, USA) with 10% fetal bovine serum, and 50 IU/ml penicillin, 50 μg/ml streptomycin and 2.5 μg/ml amphotericin B (all Sigma). Cells were incubated at 37°C in humidified air with 5% CO₂.

The HESC cells were generously gifted by Dr G Krikun (Yale University School of Medicine, USA; ATCC cell line CRL-4003) and were cultured as described previously²⁶. Briefly, cells were grown in 75 cm² tissue culture flasks containing HESC complete culture medium, comprising a 1:1 mixture of Dulbecco's modified Eagle's medium and Ham's F-12 medium (Gibco), supplemented with 12% fetal bovine serum, 1% insulin-transferrin-selenium-ethanolamine (ITS-X; Gibco), 50 IU/ml penicillin, 50 μg/ml streptomycin, and 2.5 μg/ml amphotericin B. Cells were incubated at 37°C in humidified air with 5% CO₂, and used for up to 30 passages. The identity of HESC was verified at the end of the project by short tandem repeat profiling (LGC Standards, Teddington, UK; FTA barcode STRA1969; exact match to ATCC human cell line CRL-4003).

Species	Gene	Direction	Sequence (5' → 3')
<i>Bos taurus</i>	ACTB	Forward	CAGAAGGACTCGTACGTGGG
		Reverse	TTGGCCTTGGGGTTCAGGG
<i>Bos taurus</i>	RLP19	Forward	TGTTTTCCGGCATCGAGCCCG
		Reverse	ATGCCAACTCCCGCCAGCAGAT
<i>Bos taurus</i>	HMCGR	Forward	TGAGATCCGGAGGATCCGAG
		Reverse	CAGATGGTCAGCGTCACTGT
<i>Bos taurus</i>	FDPS	Forward	ATGACGGGTAAGATCGGCAC
		Reverse	TTCTGCCCATAGTTCCTCTGC
<i>Bos taurus</i>	FDFT1	Forward	GGCACCTGAGGAGTTCTAC
		Reverse	GCATACTGCATGGCGCATT
<i>Homo sapiens</i>	ACTB	Forward	ATGATGATATCGCCGCGCTC
		Reverse	CCACCATCACGCCCTGG
<i>Homo sapiens</i>	RLP19	Forward	GCGAGCTCTTTCCTTTCGCT
		Reverse	TGCTGACGGGAGTTGGCATT
<i>Homo sapiens</i>	HMCGR	Forward	GTAACTGGAGCCAGGCTGA
		Reverse	CCTTGGATCCTCCAGATCTCAC
<i>Homo sapiens</i>	FDPS	Forward	GAGACCGGGCCTTACTTCTG
		Reverse	GGACAGGGGCATCCTTCAAA
<i>Homo sapiens</i>	FDFT1	Forward	GACTCGACAGACTCTAAGGCTC
		Reverse	TGGTCAATAAGTCGCCACG

Table 2. Sequence of target and house-keeping primers.

Cell experiments. To examine cellular resilience, BESC and HESC were seeded at a density of 50,000 cells per well in 24-well tissue culture plates (TPP, Trasadingen, Switzerland), and were cultured for 24 h in their respective complete culture media. Cells were then incubated in serum free media with atorvastatin, alendronate or zaragozic acid, using the concentrations and durations indicated in *Results*, prior to challenge with control medium or medium containing PLO. Methyl- β -cyclodextrin was used to deplete cellular cholesterol, using 0.5 mM for BESC and 1 mM for HESC. Alternatively, BESC and HESC were transfected with scramble siRNA (ON-TARGETplus Non-targeting siRNA #1; Dharmacon; gelifsciences.com) or siRNA targeting *HMCGR*, *FDPS* or *FDFT1*, which were designed using Dharmacon's siDESIGN Center (Table 1). Briefly, duplex complexes were formed by adding 100 pM of siRNA to 500 μ l/well Opti-MEM I medium (Invitrogen, Waltham, MA) in 6-well plates (TPP), and then adding 7.5 μ l Lipofectamine RNAiMAX Reagent (Invitrogen). Following 20 min incubation at room temperature, 250,000 exponentially growing cells were then seeded in 2.5 ml/well of their respective complete media for 48 h, prior to challenge with control medium or medium containing PLO.

Cell viability assays. Cell viability was assessed by the mitochondria-dependent reduction of 3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide (MTT, Sigma) to formazan, as described previously³⁰. Briefly, once supernatants were collected, the remaining cells were incubated for 2 h in 250 μ l/well serum-free culture medium containing 1 mg/ml MTT; the medium was then removed and the cells lysed with 250 μ l/well dimethyl sulfoxide (Sigma); and, optical density (OD₅₇₀) measured using a POLARstar Omega microplate reader.

The kinetics of cell survival was examined by enzymatic reduction of resazurin to resorufin using the Alamar Blue cell viability assay (Pierce, Rockford, IL, USA)³⁴. Fluorescence was measured at 545/590 nm (excitation/emission) using a POLARstar Omega microplate reader, with data expressed as the percentage of control cell survival.

Total cellular DNA concentrations were measured using the CyQuant Cell Proliferation Assay (Thermo Fisher Scientific, Loughborough, UK). Fluorescence was measured at 480/530 nm (excitation/emission), with data expressed as arbitrary fluorescence units.

Leakage of cellular LDH was measured in cell culture supernatants using the Lactate Dehydrogenase Activity Colorimetric Assay Kit (Biovision, California, USA), according to the manufacturer's instructions.

Cholesterol measurement. The BESC and HESC were grown at a density of 100,000 cells/well in 12-well tissue culture plates and treated with siRNA or inhibitors, as described above. After the treatment period, the cells were collected in 200 μ l/well 1X cholesterol assay buffer and stored in Eppendorf tubes at -20°C . When needed, the samples were defrosted at room temperature and sonicated for 10 min in a sonicating water bath. Cellular cholesterol content was measured using the Amplex[®] Red Cholesterol Assay Kit (Invivogen), according to the manufacturer's guidelines. Total cellular phospholipid was measured in the samples prepared for the cholesterol assay using a phospholipid assay kit (Sigma Aldrich, MAK122), according to the manufacturer's guidelines. The cholesterol concentrations were then normalized to the phospholipid concentrations, and the data are expressed as arbitrary units.

Quantitative PCR (qPCR). Cellular RNA was extracted using the RNeasy Mini Kit according to the manufacturer's instructions (Qiagen, Crawley, UK). The RNA was quantified using a Nanodrop ND1000 spectrophotometer (Labtech, Ringmer, UK), and 1 μ g of total RNA added to a genomic DNA elimination reaction, followed

by conversion to cDNA (Quantitect Reverse Transcription Kit, Qiagen), according to the manufacturer's instructions. Quantitative PCR was performed using intron-spanning primers (Table 2) and the IQ5 system (Bio-Rad, Hemel Hempstead, UK). The starting quantity of mRNA was determined using standard curves generated from serial dilutions of pooled reference RNA with Quantifast SYBR green (Qiagen). The target and reference genes were analyzed in triplicate, and mRNA expression normalized to the *ACTB* and *RPL19* reference genes (Table 2) using the IQ5 system (Bio-Rad), with inter-run correlation and run-dependent differences corrected using qBase software on the IQ5 system (Bio-Rad)⁴⁷. The reference genes did not differ in expression with treatment, and were amplified at the same efficiency as the target genes.

Statistical analysis. Data are presented as arithmetic mean and error bars represent SEM. The statistical unit was designated as the animal for BESC, and each independent cell passage for HESC. Statistical analysis was performed using Graphpad Prism 5.0.1 and SPSS 20.0, with $P < 0.05$ considered statistically significant. Comparisons between two treatments were tested using Student's t-test, and amongst several treatments using ANOVA, followed by Dunnett's post hoc multiple comparison test.

Data availability. All data generated or analyzed during this study are included in this published article (and its Supplementary Information files).

References

- Schneider, D. S. & Ayres, J. S. Two ways to survive infection: what resistance and tolerance can teach us about treating infectious diseases. *Nat. Rev. Immunol.* **8**, 889–895 (2008).
- Medzhitov, R., Schneider, D. S. & Soares, M. P. Disease Tolerance as a Defense Strategy. *Science* **335**, 936–941 (2012).
- Raberg, L., Sim, D. & Read, A. F. Disentangling genetic variation for resistance and tolerance to infectious diseases in animals. *Science* **318**, 812–814 (2007).
- Read, A. F., Graham, A. L. & Raberg, L. Animal defenses against infectious agents: is damage control more important than pathogen control. *PLoS Biol.* **6**, e4 (2008).
- Peraro, M. D. & van der Goot, F. G. Pore-forming toxins: ancient, but never really out of fashion. *Nat Rev Micro* **14**, 77–92 (2016).
- Tweten, R. K. Cholesterol-dependent cytolysins, a family of versatile pore-forming toxins. *Infect. Immun.* **73**, 6199–6209 (2005).
- Lange, Y., Swaisgood, M. H., Ramos, B. V. & Steck, T. L. Plasma membranes contain half the phospholipid and 90% of the cholesterol and sphingomyelin in cultured human fibroblasts. *J. Biol. Chem.* **264**, 3786–3793 (1989).
- Bloch, K. The biological synthesis of cholesterol. *Science* **150**, 19–28 (1965).
- Goldstein, J. L. & Brown, M. S. Regulation of the mevalonate pathway. *Nature* **343**, 425–430 (1990).
- Jost, B. H. & Billington, S. J. Arcanobacterium pyogenes: molecular pathogenesis of an animal opportunist. *Antonie van Leeuwenhoek* **88**, 87–102 (2005).
- Sheldon, I. M., Cronin, J., Goetze, L., Donofrio, G. & Schuberth, H. J. Defining postpartum uterine disease and the mechanisms of infection and immunity in the female reproductive tract in cattle. *Biol. Reprod.* **81**, 1025–1032 (2009).
- Bicalho, M. L. S. *et al.* Genetic and functional analysis of the bovine uterine microbiota. Part II: Purulent vaginal discharge versus healthy cows. *J. Dairy Sci.* **100**, 3863–3874 (2017).
- Sheldon, I. M., Noakes, D. E., Rycroft, A. N., Pfeiffer, D. U. & Dobson, H. Influence of uterine bacterial contamination after parturition on ovarian dominant follicle selection and follicle growth and function in cattle. *Reproduction* **123**, 837–845 (2002).
- Westermann, S., Drillich, M., Kaufmann, T. B., Madoz, L. V. & Heuwieser, W. A clinical approach to determine false positive findings of clinical endometritis by vaginoscopy by the use of uterine bacteriology and cytology in dairy cows. *Therio* **74**, 1248–1255 (2010).
- Bonnett, B. N., Martin, S. W., Gannon, V. P., Miller, R. B. & Etherington, W. G. Endometrial biopsy in Holstein-Friesian dairy cows. III. Bacteriological analysis and correlations with histological findings. *Can. J. Vet. Res.* **55**, 168–173 (1991).
- Amos, M. R. *et al.* Differential endometrial cell sensitivity to a cholesterol-dependent cytolysin links *Trueperella pyogenes* to uterine disease in cattle. *Biol. Reprod.* **90**, 54, 1–13 (2014).
- Das, A., Brown, M. S., Anderson, D. D., Goldstein, J. L. & Radhakrishnan, A. Three pools of plasma membrane cholesterol and their relation to cholesterol homeostasis. *eLife* **3**(e02882), 02881–02816 (2014).
- Kilsdonk, E. P. *et al.* Cellular cholesterol efflux mediated by cyclodextrins. *J Biol Chem* **270**, 17250–17256 (1995).
- Christian, A. E., Haynes, M. P., Phillips, M. C. & Rothblat, G. H. Use of cyclodextrins for manipulating cellular cholesterol content. *J. Lipid Res.* **38**, 2264–2272 (1997).
- Preta, G., Lotti, V., Cronin, J. G. & Sheldon, I. M. Protective role of the dynamin inhibitor Dynasore against the cholesterol-dependent cytolysin of *Trueperella pyogenes*. *FASEB J.* **29**, 1516–1528 (2015).
- Goldstein, J. L., DeBose-Boyd, R. A. & Brown, M. S. Protein sensors for membrane sterols. *Cell* **124**, 35–46 (2006).
- Sharpe, L. J. & Brown, A. J. Controlling cholesterol synthesis beyond 3-hydroxy-3-methylglutaryl-CoA reductase (HMGCR). *J Biol Chem* **288**, 18707–18715 (2013).
- Rogers, M. J. From molds and macrophages to mevalonate: a decade of progress in understanding the molecular mode of action of bisphosphonates. *Calcif. Tissue Int.* **75**, 451–461 (2004).
- Bergstrom, J. D. *et al.* Zaragozic acids: a family of fungal metabolites that are picomolar competitive inhibitors of squalene synthase. *Proc. Natl. Acad. Sci. USA* **90**, 80–84 (1993).
- Baxter, A. *et al.* Squalstatin 1, a potent inhibitor of squalene synthase, which lowers serum cholesterol *in vivo*. *J. Biol. Chem.* **267**, 11705–11708 (1992).
- Krikun, G. *et al.* A novel immortalized human endometrial stromal cell line with normal progesterational response. *Endocrinology* **145**, 2291–2296 (2004).
- Billington, S. J., Jost, B. H., Cuevas, W. A., Bright, K. R. & Songer, J. G. The Arcanobacterium (Actinomycetes) pyogenes hemolysin, pyolysin, is a novel member of the thiol-activated cytolysin family. *J. Bacteriol.* **179**, 6100–6106 (1997).
- Pokrajac, L., Baik, C., Harris, J. R., Sarraf, N. S. & Palmer, M. Partial oligomerization of pyolysin induced by a disulfide-tethered mutant. *Biochem. Cell Biol.* **90**, 709–717 (2012).
- Statt, S. *et al.* Statin-conferred enhanced cellular resistance against bacterial pore-forming toxins in airway epithelial cells. *Am. J. Respir. Cell Mol. Biol.* **53**, 689–702 (2015).
- Mosmann, T. Rapid colorimetric assay for cellular growth and survival: application to proliferation and cytotoxicity assays. *J. Immunol. Methods* **65**, 55–63 (1983).
- Park, J. M., Ng, V. H., Maeda, S., Rest, R. F. & Karin, M. Anthrolysin O and other gram-positive cytolysins are toll-like receptor 4 agonists. *J. Exp. Med.* **200**, 1647–1655 (2004).
- Gonzalez, M. R. *et al.* Pore-forming toxins induce multiple cellular responses promoting survival. *Cell. Microbiol.* **13**, 1026–1043 (2011).

33. Giddings, K. S., Johnson, A. E. & Tweten, R. K. Redefining cholesterol's role in the mechanism of the cholesterol-dependent cytolysins. *Proc. Natl. Acad. Sci. USA* **100**, 11315–11220 (2003).
34. Page, B., Page, M. & Noel, C. A new fluorometric assay for cytotoxicity measurements *in-vitro*. *Int. J. Oncol.* **3**, 473–476 (1993).
35. Billington, S. J., Songer, J. G. & Jost, B. H. Molecular characterization of the pore-forming toxin, pyolysin, a major virulence determinant of *Arcanobacterium pyogenes*. *Vet. Microbiol.* **82**, 261–274 (2001).
36. Billington, S. J., Songer, J. G. & Jost, B. H. The variant undecapeptide sequence of the *Arcanobacterium pyogenes* haemolysin, pyolysin, is required for full cytolytic activity. *Microbiology* **148**, 3947–3954 (2002).
37. Ballard, D. O., Upsher, A. E. & Seely, D. D. Infection with *Corynebacterium pyogenes* in man. *Am. J. Clin. Pathol.* **17**, 209–215 (1947).
38. Gurcel, L., Abrami, L., Girardin, S., Tschopp, J. & van der Goot, F. G. Caspase-1 activation of lipid metabolic pathways in response to bacterial pore-forming toxins promotes cell survival. *Cell* **126**, 1135–1145 (2006).
39. Cronin, J. G., Kanamarlapudi, V., Thornton, C. A. & Sheldon, I. M. Signal transducer and activator of transcription-3 licenses Toll-like receptor 4-dependent interleukin (IL)-6 and IL-8 production via IL-6 receptor-positive feedback in endometrial cells. *Mucosal Immunol* **9**, 1125–1136 (2016).
40. Turner, M. L., Cronin, J. C., Healey, G. D. & Sheldon, I. M. Epithelial and stromal cells of bovine endometrium have roles in innate immunity and initiate inflammatory responses to bacterial lipopeptides *in vitro* via Toll-like receptors TLR2, TLR1 and TLR6. *Endocrinology* **155**, 1453–1465 (2014).
41. Bellier, A., Chen, C. S., Kao, C. Y., Cinar, H. N. & Aroian, R. V. Hypoxia and the hypoxic response pathway protect against pore-forming toxins in *C. elegans*. *PLoS Pathog.* **5**, e1000689 (2009).
42. Kao, C. Y. *et al.* Global functional analyses of cellular responses to pore-forming toxins. *PLoS Pathog.* **7**, e1001314 (2011).
43. Huffman, D. L. *et al.* Mitogen-activated protein kinase pathways defend against bacterial pore-forming toxins. *Proc. Natl. Acad. Sci. USA* **101**, 10995–11000 (2004).
44. Bischof, L. J. *et al.* Activation of the unfolded protein response is required for defenses against bacterial pore-forming toxin *in vivo*. *PLoS Pathog.* **4**, e1000176 (2008).
45. Rosch, J. W. *et al.* Statins protect against fulminant pneumococcal infection and cytolysin toxicity in a mouse model of sickle cell disease. *J. Clin. Invest.* **120**, 627–635 (2010).
46. Healey, G. D. *et al.* Mevalonate biosynthesis intermediates are key regulators of innate immunity in bovine endometritis. *J. Immunol.* **196**, 823–831 (2016).
47. Bustin, S. A. *et al.* The MIQE guidelines: minimum information for publication of quantitative real-time PCR experiments. *Clin. Chem.* **55**, 611–622 (2009).

Acknowledgements

This work was funded by the United Kingdom Biotechnology and Biological Sciences Research Council (Grant BB/K006592/1), and SG was funded by a Swansea University Postgraduate Scholarship. We thank Prof BH Jost for access to recombinant PLO, Prof M Palmer for DS-PLO, and Dr G Krikun for HESC. We also thank J Cronin, G Healey, M Amos and S Owens for advice.

Author Contributions

S.G. designed and performed the experiments, collected and analyzed data, and drafted the manuscript; G.P. designed and performed the experiments, and collected and analyzed data; I.M.S. conceived and designed the work, interpreted data, and wrote the manuscript. All authors approved the final version of the manuscript.

Additional Information

Supplementary information accompanies this paper at <https://doi.org/10.1038/s41598-017-17138-y>.

Competing Interests: The authors declare that they have no competing interests.

Publisher's note: Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.



Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons license, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons license and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this license, visit <http://creativecommons.org/licenses/by/4.0/>.

© The Author(s) 2017