

RESEARCH

Open Access



"I know a lot about medicinal plants. I read, I watch, and I search": towards hybrid knowledge systems in the modern era

Julia Prakofjewa^{1,2*} , Luigi Conte^{1,2*} , David Ludwig³ , Povilas Šarka^{1,4} , Pietro Centorrino¹ , Raivo Kalle⁵  and Renata Sõukand^{1,2} 

Abstract

Background Hybrid knowledge systems are central to community negotiations of environmental, social, and epistemic pressures. In multilingual borderland areas, interactions between local ecological knowledge (LEK), formal, and popular knowledge systems remain underexplored, despite their importance for the persistence and transformation of medicinal plant use today.

Methods We conducted 67 semi-structured interviews and participant observation in 21 rural settlements of the Vilnius region (Lithuania), an area bordering Belarus, focusing on the two largest local groups, Lithuanians (LT) and Poles (PL). Detailed Use Reports ($n = 1446$) on medicinal plant use were coded by the origin of knowledge, classified as local, formal, or popular, and the degree of hybridisation was quantified using the Shannon–Wiener diversity index and hybridisation metrics. Sociodemographic variables (age, gender, education, and multilingualism) were tested for associations with hybridisation using Spearman's ρ and Student's t -tests.

Results A total of 139 medicinal taxa were recorded, of which 68 (49%) were shared between the two groups. Overall, recorded medicinal plant knowledge remained primarily grounded in LEK, sustained through intergenerational transmission. Compared with PL, LT interviewees drew on a broader mix of knowledge-origin domains ($H' = 0.97$ vs 0.52) and combined them more often ($HD = 0.195$ vs 0.059). In total, 39 taxa showed hybrid use, predominantly in the LT group. Hybridisation was negatively associated with age but positively correlated with the number of listed plants and their reported uses, while multilingualism showed a near-significant positive trend.

Conclusions The study suggests that medicinal plant knowledge has evolved here through hybridisation, a process whose consequences are context-dependent, offering opportunities for revitalisation but also a risk of displacement. Dialogic exchanges across families, communities, languages, and media expand people's plant repertoire and strengthen community adaptive capacity. Yet when these exchanges lead to excessive standardisation, they risk eroding the diversity of local traditions. Ethnobotanical research must therefore go beyond documenting popular

*Correspondence:

Julia Prakofjewa
yuliya.prakofyeva@unive.it
Luigi Conte
luigi.conte@unive.it

Full list of author information is available at the end of the article



© The Author(s) 2026. **Open Access** This article is licensed under a Creative Commons Attribution-NonCommercial-NoDerivatives 4.0 International License, which permits any non-commercial use, sharing, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if you modified the licensed material. You do not have permission under this licence to share adapted material derived from this article or parts of it. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by-nc-nd/4.0/>.

and formal knowledge sources to interrogate how linguistic and sociopolitical contexts condition the emergence of hybrid knowledge systems, privileging certain forms while rendering others transformed or marginalised.

Keywords Degree of hybridisation, Dzūkija, Formal knowledge, Knowledge diversity, Knowledge transmission, Knowledge systems, Medicinal plants, South-eastern Lithuania, Local ecological knowledge

Background

Contemporary societies face what has been called a *polycrisis*: environmental decline, political turbulence, and uncertainty about knowledge systems reinforce one another [13]. In this context, the ways in which local ecological knowledge (LEK) circulates and persists are being reshaped [16]. Beyond globalisation and climate shocks, the rise of digital media and shifts in governance introduce new channels, authorities and metrics of credibility for local knowledges [7]. In various marginal contexts, such shifts change *how* people value and recognise plant expertise, which remains vital for the health and resilience of local communities [28]. Medicinal plant knowledge is especially exposed to these dynamics because it is rooted in embodied practice, intergenerational memory, and locally embedded systems of care [72]. How such knowledge is remembered, adapted, or forgotten cannot be separated from broader regional trajectories of economy and politics [17].

Hybridisation has been identified as a key driver of both change and continuity in local knowledge systems [21]. In this study, knowledge hybridisation is conceptualised as a dynamic process through which elements from different knowledge domains, such as local, scientific, and popular knowledge, are combined, reinterpreted, and recontextualised in practice. Consistent with this, Yanou et al. [74] document a mixed picture in which certain LEK elements decline, whereas others persist by incorporating outside knowledge. However, Sharifian et al. [62] highlighted that little is known about LEK hybridisation processes. Where hybrid forms are documented, such as across different types of pastoral mobility and specific knowledge domains, the evidence is fragmentary, and patterns are inconsistent. Despite growing interest in knowledge hybridisation, systematic analyses of how local, formal, and popular epistemologies interact are still scarce [10, 35, 67], especially via digitally mediated, informal channels in multilingual, post-socialist regions.

One necessary consequence is that knowledge hybridisation, in response to historical and contemporary pressures, has become a key strategy through which local communities negotiate complex epistemological landscapes [78]. Often, practitioners have adapted known remedies to meet emerging conditions while also developing new medicines in response to novel challenges: “*new practices for new problems*” [75]. Rather than signalling cultural decline [38], such a blend of local, formal, and popular knowledge systems often reflects deliberate,

situated, and context-specific strategies for navigating shifting social-ecological dynamics [76].

Rather than treating knowledge traditions as historically autonomous, we emphasise their longstanding entanglement [45]. With the growing authority of educational, medical, and regulatory institutions, plant knowledge became shaped by more standardised frameworks. In 19th- and 20th-century Europe, people continued to rely on longstanding vernacular traditions of healing, yet plant use was increasingly redirected through formal schooling, professional medicine, and state-led classification systems [48]. By the mid-1900s, written sources such as herbals, school textbooks, technical manuals, and popular literature had become the dominant repositories of botanical knowledge [36]. Regional differences in the use of medicinal plants have gradually weakened due to the spread of formal public education and the adoption of binomial nomenclature, which has made it easier to create standardised botanical knowledge frameworks [31, 54].

The use of medicinal plants in the Vilnius region and surrounding areas has been well-documented by Lithuanian scholars [32, 33, 50, 58, 66, 73]. However, much less is known about how these traditions have hybridised across ethnic groups under contemporary conditions.

After WWII in the Vilnius region, as in the wider western borderlands of the former Soviet Union, knowledge homogenisation processes were reinforced by centralised institutions, including the public education system, state healthcare, and official pharmacies [5, 41, 42]. In the studied region, the collection of medicinal plants was placed under state supervision. Authorities organised seasonal campaigns, assigned quotas to local communities, and prescribed which species were considered medicinal in the official pharmacopoeias. These measures did not only structure the work of foragers but also limited standard practices [9, 11] and reduced the diversity of local plant knowledge [58].

Building on recent debates [12, 35, 38], we approach hybridisation as a complex and context-sensitive process whose consequences are not intrinsically beneficial or detrimental. While mutual learning and cross-fertilisation (e.g., through dialogic knowledge co-production) may lead to expansion or reconfiguration of knowledge content, practices and repertoires [68], this does not necessarily imply an increase in epistemic diversity in the strict sense. At the same time, hybridisation can facilitate epistemic homogenisation [49] when one knowledge

system comes to dominate and redefine the standards of credibility and legitimacy [63]. These ambivalences are not unique to the present but resonate with longer historical trajectories, as plant-related knowledge has long been produced, transmitted, and validated through shifting configurations of authority across generations.

This study advances a nuanced understanding of knowledge hybridisation by examining how LEK, formal, and popular epistemologies intersect in shaping contemporary medicinal plant use. Drawing on fieldwork in the Lithuanian-Belarusian borderland, a historically multi-layered region, we analyse how hybrid forms of medicinal plant knowledge take shape among the two largest local ethnic groups, Lithuanians and Poles. Specifically, we aim to (1) document existing medicinal plant knowledge, (2) compare knowledge content, sources, and transmission pathways across ethnic groups, and (3) assess the degree and character of hybridisation in relation to key sociodemographic variables.

Materials and methods

Study site

Our fieldwork took place in the Vilnius region (Lithuania), close to the Belarusian border (Fig. 1). It was conducted as a part of a broader investigation of ethnobotanical and ethnomedicinal knowledge in the western borderlands of the former Soviet Union [41, 42, 55–57].

This area marks a transition zone between the central Lithuanian lowlands and the western Belarusian uplands, with altitudes varying from about 120 to 240 m and in places rising to nearly 300 m above sea level (locally reaching up to ~294 m). The environment is characterised by sandy, low-fertility soils, extensive peatlands and wetlands [39]. The climate is humid continental (Köppen

Dfb), with a mean annual temperature of ~7.2 °C and mean annual precipitation ranging from 685 mm (Vilnius) to 701 mm (Šalčininkai) [19]. Forest cover in the region is extensive, with pine and mixed stands occupying more than half of the territory. According to Juknelienė and Mozgeris [26], the largest expansion of new forests in the second half of the twentieth century took place here. At the same time, the Šalčininkai municipality underwent considerable deforestation, illustrating how contrasting land-use processes can coexist within the same landscape. These shifts in forest cover are directly relevant for ethnobotanical practices, altering the accessibility and abundance of medicinal species gathered in forests, meadows, and wetland margins.

Historically, this territory was subject to numerous shifts in political affiliation over the last millennium [25]. The Vilnius region is defined as *Vilnija* or *Vilniaus kraštas* in Lithuanian or *Wileńszczyzna* (*Vilenszczyzna*) in Polish and Belarusian historiographies [69]. Within Lithuanian ethnographic discourse, however, this same area is more often associated with *Dzūkija*, a cultural-geographical region institutionalised during the Soviet period [69]. Its borders were largely drawn to coincide with those of the former Vilnius region [8]. Historically, however, the area extended beyond present-day Lithuania to include parts of what are now Belarus and Poland. The prominent “peninsula” of Lithuanian territory, which Belarus nearly encircles, is a distinctive feature of this landscape.

The studied settlements were predominantly rural and dispersed, with a relatively low population density. Proximity to Vilnius, the capital of Lithuania, added another layer of socio-ecological complexity to the study area. In the nineteenth century, Vilnius was a major centre



Fig. 1 Characteristic views of the study site, illustrating ecological and cultural conditions (from left to right): coniferous forest, rural road through fields, traditional homestead with cattle, vernacular cottage garden, roadside cross as a long-standing Roman-Catholic tradition of sacral markers in the countryside, and mural in a local Polish-language school (“Przez oświatę do wolności i szczęścia” [“Through education to freedom and happiness”]). Photo credits: JP, 2018–2019

for medicinal plant harvesting and the wholesale trade [30, 44]. The study region thus lies at a liminal interface between state-driven modernisation and enduring local traditions [61]. In the 1980s, many residents worked or studied in Vilnius during the week but returned to the villages on weekends and holidays. This rural–urban mobility pattern sustained local ecological practices and supported the preservation of LEK despite growing socioeconomic entanglement with the metropolitan core.

The region has long been characterised by the coexistence of multiple ethnic groups, languages, and cultural traditions, including Lithuanian, Polish, Belarusian, Jewish, Romani, Tatar, and Russian, among others. Although different ethnic groups may rely on overlapping epistemic principles, their long-term coexistence in the region has contributed to plural, historically layered practices and interpretations of human–environment relations [20].

Data collection

A total of 67 deep semi-structured interviews were conducted across 21 rural settlements of the Vilnius region between July 2018 and September 2019. Sampling targeted the two most represented ethnic groups in the area: Lithuanians (LT) and Poles (PL), using convenience sampling supplemented by snowball sampling where possible.

The gender distribution was skewed in favour of women (50 women, 17 men). It reflects common demographic trends in many post-Soviet societies [71], where male life expectancy remains significantly lower than that of females. Study participants ranged in age from 39 to 97 years. The mean age was about 69 years (65 years for LT ($n=30$) and 71 years for PL ($n=37$)).

The majority of interviewees (49 out of 67) spoke two or more languages, indicating that multilingualism was a common characteristic among them. The average number of languages reported by interviewees was 2, with LT showing greater linguistic diversity (mean=3) than PL (mean=2). Five LT and two PL interviewees stated that they were proficient in four languages. Common language combinations included Lithuanian (with regional forms such as *dzūkiškai*), Belarusian (with phrases like “po prostu” [simply], “po swojemu” [in one’s own way], and “pa tutejszemu” [in the local way]), and Russian among LT, as well as Polish and Russian among PL.

In terms of educational level, the largest group had completed post-secondary non-tertiary ($n=25$), followed by those with primary ($n=19$), upper secondary ($n=12$), and lower secondary ($n=10$). Only one interviewee reported an equivalent tertiary education level. All interviewees identified themselves as Roman Catholics.

We conducted interviews in the language(s) preferred by the participants. The first and fourth authors’ competencies made this multilingual approach possible,

encouraging more complete responses and an open exchange of knowledge and experiences. In most cases, interviewees naturally switched between languages to choose the one that best suited their needs, particularly when discussing local and scientific plant names as well as local, formal, and popular uses. The interviews were transcribed with linguistic nuances, ensuring that all culturally embedded elements of speech were accurately captured for further interpretation and analysis.

Interviewees were asked to name and describe the plant species they currently or previously foraged or cultivated for medicinal purposes, the ailments treated with these plants, and the corresponding methods of preparation. In addition, information was collected on the sources of medicinal plant knowledge and on whether plant use referred to past or present practices. When possible, we gathered plants together with interviewees to collect and identify voucher specimens. Most interviewees noted they only knew plants by their local names, not their scientific or common names. In some cases, interviewees could not recall plant names but provided descriptions or pointed to them, allowing reliable identification. Collected specimens and dried samples are housed at the Herbarium of Ca’ Foscari University of Venice in Italy (DZULT01–DZULT137; DDZULT01–DDZULT42). Plant identification was carried out using the Plants of the World Online database [52] and Flora Europaea [70], with family classification following [64].

Data analysis

The data were compiled and categorised according to the medicinal uses of plants. We structured the emic categories into Detailed Use Reports (DUR), each documenting the medicinal use, plant part, preparation method, the Latin and local names, time of use, knowledge origin, and codes for the interviewee’s region and ethnic group.

To evaluate the degree of similarity in medicinal plant knowledge between compared ethnic groups, we calculated the Jaccard Index (JI) following González-Tejero et al. [22], based on the presence/absence of reported plant uses. The index was computed using the formula

$JI = \frac{\gamma}{\alpha + \beta - \gamma}$, where α is the total number of plant uses reported by LT, β is the total number of plant uses reported by PL, and γ is the number of plant uses shared by both groups.

Knowledge origin categories were developed inductively from interviewee narratives and divided into three domains: local (A), formal (B), and popular (C). Within each domain, several subcategories captured specific pathways of knowledge transmission (e.g., intergenerational oral transfer, formal education, social media). The coding scheme is outlined in Table 1.

Table 1 Knowledge origin, description of categories

Subcategory	Description	Examples from interviews
A-Local Knowledge		
A1 Intergenerational family transfer	Transmitted mainly orally from parents, grandparents, or close relatives	"I learnt it from my father ." "My grandmother did it, my mother did it, and now I do it too"
A2 Horizontal community transition	Learnt from elders, peers, or respected local figures, but not relatives	"Our neighbour told me that this helps a lot." "The old women in our village always used this plant for wounds"
A3 Oblique transmission (elders/teachers)	Knowledge received from older, non-kin community members or local experts (e.g., teachers, local healers/herbalists)	"At school, our teacher explained the healing plants to us." "The local herbalist told me this works for coughs"
A4 Historical/ancestral roots	Framed as time-tested or long-standing knowledge with profound historical continuity	"This is traditional knowledge passed down for generations ". "This is how it has always been done ", "This knowledge has been known for centuries ". "This has been known for ages "
A5 Own practice/experimentation	Acquired through individual practice, observation, or trial-and-error	"I figured it out myself by trying different things: "I experimented with this plant"
B-Formal Knowledge		
B1 Formal publications	Scientific, peer-reviewed, expert-validated herbals or monographs	"I read it in a medicinal book ": "It was written in a scientific article I read"
B2 Health professionals	Information or recommendations from certified practitioners	"My doctor recommended this"
B3 School/university instruction	Knowledge obtained through formal education	"We remember this from school "
B4 Digital medical resources	Research-based content on medical websites or portals	"I looked it up on a medical website ": "I googled it and found this information on a special medicinal portal "
C-Popular Knowledge		
C1 Newspapers/magazines	General audience press, local or specialised	"I read this in one local magazine ": "I specifically subscribed to a local newspaper dedicated to traditional medicine"
C2 Radio/TV	Mass-media programs	"I have seen it on a morning TV show "
C3 Social media and popular websites	Facebook groups, Instagram, forums, YouTube	"I saw this information in one popular online forum and decided to try it": "I have seen it on YouTube ."

The **degree of hybridisation** (HD) is defined as the proportion of use reports that draw on more than one knowledge-origin domain (AB, AC, BC, or ABC) relative to the total number of use reports. For each plant species and ethnic group, HD was calculated as:

$$HD = \frac{\text{Number of hybrid use reports}}{\text{Total number of use reports}} \quad (1)$$

To enable cross-ethnic comparison of hybridisation patterns, this metric was calculated separately for LT and PL. HD=1 corresponds to full hybrid knowledge of plant uses, while HD=0 indicates a singular and consistent source.

Each DUR was associated with a binary code: 0 for non-hybrid (citing exactly one knowledge origin: A, B, or C) and 1 for hybrid (citing ≥ 2 origins: AB, AC, BC, or ABC). We also calculated the **internal degree of hybridisation** within each general knowledge origin (A, B, or C), defined as the proportion of use reports that included multiple subcategories (e.g., A1, A3, and A4). These were coded analogously (1=internally hybrid, 0=internally non-hybrid), and the proportion was computed using the definition of HD in Eq. 1. A plant is considered fully hybridised (ABC) only when supported by at least two DUR for that taxon explicitly combining all three knowledge domains.

To assess **the diversity of knowledge origins** within and between studied ethnic groups, per taxon or medicinal use, we computed the Shannon–Wiener diversity index (H') over distinct origin categories [46]:

$$H' = -\sum_{i=1}^S p_i \ln(p_i) \quad (2)$$

where S is the total number of different origin categories (e.g., knowledge origin types), p_i is the proportion of the total number of medicinal uses in the i -th category, and \ln denotes the natural logarithm. This index enabled us to assess the heterogeneity of knowledge systems by capturing the richness and distribution of knowledge across origin types within and between the sampled ethnic groups. We used Spearman's rank correlation coefficient to assess whether sociodemographic variables (age, gender, education, the number of languages spoken) were associated with the degree of hybridisation. This helped us to identify patterns and potential drivers behind the emergence and distribution of hybrid knowledge systems in the study region. The group differences were assessed using Student's t -test (robustness confirmed with Wilcoxon rank-sum tests).

All statistical analyses and visualisations were conducted using R v. 4.5.0 [59] and PAST 5 [27].

Results

Medicinal plant knowledge diversity

We recorded 139 taxa belonging to 58 botanical families used for healing and prophylactic purposes (Table 2). The most represented were Asteraceae (21), Rosaceae (16), Lamiaceae (11), Ericaceae (7), Apiaceae (6), and Poaceae (5). Of these, 118 taxa were documented among the LT and 89 among the PL interviewees. Ultimately, 68 plant species (49%) were common to both ethnic groups. In total, 1446 DUR were collected.

The amount of reported medicinal plant uses differed significantly between the studied ethnic groups. LT interviewees named a higher number of DUR (mean \pm standard deviation (SD)) = 28.0 ± 21.4) than PL interviewees (16.4 ± 16.0); Student's $t = 2.458$, $p = 0.017$, $n = 67$. This finding indicates a broader repertoire of medicinal applications among LT. These quantitative differences are further contextualised by cross-ethnic patterns of overlap (Fig. 2), which illustrate the Jaccard indices and proportional Venn diagrams of medicinal taxa cited by LT and PL interviewees, including those mentioned by at least three people.

Approximately 80% of reported uses involved wild plants, while cultivated species constituted approximately 18%. The LT group mentioned a higher number of taxa overall in both the wild and cultivated categories. In both groups, wild resources were used more extensively for healing and prophylactic purposes than cultivated ones.

The most frequently mentioned taxa used among all interviewees included *Artemisia absinthium* (87 DUR), *Rubus idaeus* (79), *Matricaria chamomilla* (67), *Calendula officinalis* (57), *Achillea millefolium* (52), and *Tilia cordata* (47), with examples of use shown in Fig. 3. LT most frequently cited *Artemisia absinthium* (49 DUR), *Rubus idaeus* (37), *Matricaria chamomilla* (36), *Calendula officinalis* (34), *Achillea millefolium* (31), *Tilia cordata* (26), *Betula* spp. (24), and *Carum carvi* (21). Among PL were recorded as mostly cited: *Rubus idaeus* (42), *Artemisia absinthium* (38), *Matricaria chamomilla* (31), *Vaccinium myrtillus* (28), *Calendula officinalis* (23), *Aloe* (22), *Achillea millefolium* (21) and *Tilia cordata* (21).

A comparison of the taxa mentioned by interviewees revealed a statistically significant gender difference (Spearman's $\rho = -0.295$, $p = 0.015$). Compared to male interviewees (9.9 ± 7.1), female interviewees reported significantly more taxa on average (mean \pm SD = 14.9 ± 9.7). Other variables, however, did not exhibit any correlations. Our analyses indicated no significant relationship between age and medicinal plant knowledge (Spearman's $\rho = -0.112$, $p = 0.368$) or between multilingualism and such knowledge (Spearman's $\rho = 0.170$, $p = 0.168$). Educational level was also non-significant for the number of

Table 2 Medicinal plant knowledge diversity in the Vilnius region (Lithuania), as named and reported by members of the Lithuanian (LT) and Polish (PL) communities. DUR refers to the number of detailed use reports per plant taxon

Family	Latin name; voucher number	Local name(s)	Used part(s)	Preparation	Medicinal use	DUR	
						LT	PL
Acoraceae	<i>Acorus calamus</i> L.; DZULT080	LT: ajerai; ajeras PL: aer, ajeras	leaves	eaten fresh	appetiser	1	1
			roots	alcoholic infusion	bellyache	1	1
				decoction	healthy	1	1
				decoction, bath	toothache	1	1
				tea	hair care	1	1
				topical application	head lice	1	1
				topical application	appetiser	1	1
				topical application	toothache	1	1
Amaranthaceae	<i>Beta vulgaris</i> L.	LT: burokas; PL: burak	leaves	topical application	abscesses	1	1
				topical application	abscesses	1	1
				eaten cooked	wounds	1	1
			roots	juice	constipation	2	2
				juice, to rinse	improve blood quality	1	1
				baked, topical application	tonsillitis	1	1
			bulbs	boiled with milk	abscesses	2	3
		LT: svogūnas; PL: cebula, cybula		syrup, drops	cough	1	1
				topical application	cough	1	1
				topical application	earache	1	1
				topical application	runny nose	1	1
			bulbs	eaten fresh	earache	1	1
Apiaceae	<i>Allium sativum</i> L. <i>Aegopodium podagraria</i> L. <i>Anethum graveolens</i> L.; DZULT063, DDZULT29	PL: čosnyk LT: garšva LT: krop, krapai; PL: krapai	bulbs	eaten fresh	runny nose	1	1
			young leaves	eaten fresh	healthy	1	1
			erial parts	tea	hypertension	1	1
			seeds	boiled with milk	hypertension	1	1
				tea	headache	1	1
			seeds	alcoholic infusion	hypertension	5	1
					cold	1	1
		LT: kmynai, kmynas; PL: kmin, kminek, kmynas, kmin	seeds		cold	1	1
	<i>Carum carvi</i> L.; DDZULT40				fever	1	1
					flu	1	1
					stomach ache	1	1
				decoction	for everything	1	1
				tea	stomach ache	2	2
					cold	1	1

Table 2 (continued)

Family	Latin name; voucher number	Local name(s)	Used part(s)	Preparation	Medicinal use	DUR	
						LT	PL
					cough	2	1
					diarrhoea	2	
					digestion problems	1	
					fever	3	
					flatulence	2	
					from losing weight	1	
					headache	1	
					healthy	1	
					heart problems	1	
					helpful	1	
					hypertension	2	1
					immune boosting	1	
					organism cleansing	1	
					sedative	1	
					stomach ache	2	
				with apple juice	stomach ache	1	
			roots	boiled with milk, topical application	abscesses	1	
	<i>Daucus carota</i> L.	LT: morka, PL: marchewka, morka		juice	gastric ulcer	1	
				topical application	heart rate	1	
			seeds	tea	abscesses	1	
			roots	alcohol mac- eration, topical application	hypertension joint pain	1	
	<i>Ostericum palustre</i> (Besser) Besser	LT: starédub		decoction	urinary bladder	1	
	<i>Petroselinum crispum</i> (Mill.) Fuss; DDZULT08	PL: pietruszka, petražoles	aerial parts				
			leaves	decoction	kidney diseases	1	
			leaves	tea	urinary bladder	1	
			leaves	alcoholic maceration	for everything	2	
Asphodelaceae	<i>Aloe</i> spp. (mainly <i>Aloe arborescens</i> Mill.)	LT: aljošius, aliaš, PL: aliasz, aloesz, aloje, staletnik, alijas, aljošius			gastric ulcer	1	
					sore throat	1	
					stomach ache	2	
			eaten fresh		earache	1	
					for everything	1	

Table 2 (continued)

Family	Latin name; voucher number	Local name(s)	Used part(s)	Preparation	Medicinal use	DUR	
						LT	PL
				topical application	bleeding	1	1
			inflorescences	decoction	for everything	2	
				tea	improve blood quality	1	
			leaves	decoction	internal bleeding	1	
					abscesses	1	
					stomach ache	1	
				topical application	abscesses	1	
					bleeding	11	10
					cuts	5	3
					wounds	4	
			leaves	topical application	external pain	1	
					headache	2	
					joint pain	9	9
					knee ache	3	3
					limb pain	4	
					wounds	1	
			roots	decoction	stomach cancer	1	
			aerial parts	compress	leg strain	1	
			aerial parts	tea	women's diseases	2	
					appetiser	1	1
					bellyache	1	4
					diarrhoea	1	6
					for everything	1	
					helminthic infection	1	
					prophylactics	1	
					stomach ache	4	
				fumigation	fright	1	
				put in pillow	headache	1	
					sedative	1	
				tea	appetiser	2	
					bellyache	6	6
					diarrhoea	12	6
					digestion problems	6	2

Arctium tomentosum Mill.; DZULT116LT: lapušnykai, varnalėša, vilkai, varnalėšos;
PL: lapuch, łopuch*Arnica montana* L.

LT: trontnik

Artemisia abrotanum L.; DZULT077;
DZULT129

LT: dievo medelis, diemedis, božje dreūka

Artemisia absinthium L.; DDZULT27,
DZULT052, DZULT083LT: černobilis, kietis, pelūnas, pelyna, pelynas, pelynas, kartusis
kietis, pelyn, pialynas;
PL: pałyn, pelynas, pialynas, pietun, piolun, polyń

Table 2 (continued)

Family	Latin name; voucher number	Local name(s)	Used part(s)	Preparation	Medicinal use	DUR	
						LT	PL
					diaper rash	1	
					diathesis in children	1	
					eye inflammation	1	
					eye pain	1	
					eye problems	2	
					inflammation	1	
					processes		
				decoction, to rinse	abscesses	1	
					sore throat	2	
					tonsillitis	1	
				oil infusion	stomach ache	1	
				tea	allergy	1	
					bladder inflammation	2	
					blisters	1	
					cold	2	
					cough	2	
					diaper rash	1	
					digestion problems	1	
					for everything	1	
					hair care	1	
					headache	1	
					helpful	1	
					inflammation	5	
					processes		
					kidney diseases	1	
					liver diseases	1	
					organism cleansing	1	
					sore throat	1	
					stomach ache	1	
					urinary bladder	1	
					weightlifting injuries	1	
					women's diseases	2	
					fright	1	
			airial parts	tea	for everything	1	
			airial parts	tea	inflammation	1	
					processes		
	<i>Carduus</i> spp.						
	<i>Gyanus segetum</i> Hill; DDZULT31, DZULT10						
		LT: [forgotten name]					
		LT: ruglagelè, ruglagelès, vosilkos					

Table 2 (continued)

Family	Latin name; voucher number	Local name(s)	Used part(s)	Preparation	Medicinal use	DUR	
						LT	PL
	<i>Echinacea purpurea</i> (L.) Moench	LT: ežiulė	inflorescences	tea	bladder inflammation	1	
	<i>Helianthus annuus</i> L.	PL: padošnuch			heart rate	1	
	<i>Helichysum arenarium</i> (L.) Moench; DZULT134, DDZULT37, DDZULT06	LT: cosankos, šlamutis, snaudės; PL: suchawiejka	flowers	tea	pneumonia	1	
			roots	tea	sedative for children	1	
			aerial parts	decoction	urinary inflammation	1	
					immune boosting	1	
					healthy	1	
					liver diseases	2	
			flowers	tea	helpful	1	
					liver diseases	1	
					improve blood quality	1	
					liver diseases	1	
					sleep disturbance	1	
			roots	tea	stomach ache	1	
			roots	alcohol mac- eration, topical application	bruises	1	
	<i>Inula helenium</i> L.; DZULT078, DZULT097	LT: debesylas, deviasylas, debesylas dialysis; PL: dziewiasif, dziwiasif			tonic	1	
					joint pain	2	
					weightlifting injuries	2	
					back pain	1	
					for everything	2	
					joint pain	2	
					uterine prolapse	1	
					weightlifting injuries	1	
					back pain	1	
					for everything	1	
					joint pain	1	
					bellyache	2	
					external pain	1	
					weightlifting injuries	1	
					healthy	1	
					weightlifting injuries	1	
					cold	1	
	<i>Matricaria chamomilla</i> L.; DDZULT26, DZULT036, DZULT059, DDZULT07	LT: ramunėlės; PL: ramaška, rumianački, romashka, rumianek, ramunėlės, rumianki, rumiański	aerial parts	alcoholic infusion			
				compress	diaper rash	2	
					eye inflammation	1	

Table 2 (continued)

Family	Latin name; voucher number	Local name(s)	Used part(s)	Preparation	Medicinal use	DUR	
						LT	PL
					eye pain		1
					eye problems		2
					healthy		1
					neck pain		1
					sore throat		1
					wounds		1
				decoction	bellyache in children		2
					eye pain		1
					healthy		1
					helpful		1
					improve vision		1
					sore throat		1
					stomach ache		1
				decoction, bath	abscesses		1
					diaper rash		1
					diathesis in children		4
					eye problems		2
					skin diseases		1
				drops	earache		1
					eye inflammation		1
					eye pain		1
				tea	bellyache		1
					cold		1
					digestion problems		1
					earache		1
					fever		1
					flatulence		1
					for everything		1
					healthy		3
					helpful		1
					runny nose		1
					sedative		1
					sore throat		2
					stomach ache		1
				inflorescences	for everything		1
				decoction	inflammation		1
					processes		1
				decoction, bath	abscesses		1

Table 2 (continued)

Family	Latin name; voucher number	Local name(s)	Used part(s)	Preparation	Medicinal use	DUR	
						LT	PL
				decoction, to rinse tea	sore throat bellyache in children digestion problems healthy headache inflammation processes for heart sore throat fright	1 1 1 1 1 1 1 1 1	1 1 2 1 1
	<i>Scorzoneroideis autumnalis</i> (L.) Moench	PL: malačaji	aerial parts	decoction	for heart sore throat fright	1 1 1	1 1 1
	<i>Silybum marianum</i> (L.) Gaertn.; DZULT121	LT: čertapalakas, čiertapaloch; PL: čartapaloch, czartapaloch	aerial parts	decoction, bath	fright	1	1
	<i>Tanacetum vulgare</i> L.; DZULT066	LT: pižma, bitkréslié; PL: pižma	flowers leaves aerial parts	fumigation fumigation decoction decoction	fright fright fright haemorrhoids	1 3 1 1	1 3 1 1
	<i>Taraxacum officinale</i> (L.) Weber ex FH.Wigg.; DZULT094	LT: piené; PL: adywanczyk	inflorescences inflorescences	whisked in the sauna decoction alcohol mac- eration, topical application tea	healthy organism cleansing healthy helminthic infection joint pain	1 1 1 1 1	1 1 1 1 1
	<i>Tussilago farfara</i> L.; DZULT108, DDZULT04	LT: mać-maćécha, šaltapusnis, šalpusniai, padbielas; PL: mać-maćécha	leaves roots sap inflorescences	eaten fresh tea topical application alcoholic infusion	cough for everything liver diseases liver diseases warts joint pain	1 1 1 1 1 1	1 1 1 1 1 1
			leaves	tea decoction	cold cough bellyache cough helpful bronchitis	2 1 1 2 1 1	2 1 1 2 1 1

Table 2 (continued)

Family	Latin name; voucher number	Local name(s)	Used part(s)	Preparation	Medicinal use	DUR	
						LT	PL
Betulaceae	<i>Alnus</i> spp. <i>Betula</i> spp. (including <i>Betula pendula</i> Roth, <i>Betula pubescens</i> Ehrh); DZULT013, DZULT050	LT: alksnis; PL: olcha LT: beržas; PL: brzoža, beržas, biaoža, bierježa			cold	1	
					cough	2	
					sore throat	1	
					joint pain	1	
					wounds	1	
					abscesses	1	
					haemorrhoids	1	
					diuretic	1	
					kidney diseases	1	
					diuretic	1	
Boraginaceae	<i>Corylus avellana</i> L. <i>Symphytum officinale</i> L.; DZULT045, DZULT070	LT: lazdynas LT: kaulžolė, žvakost, žvakostas, starodovas, riebešaknis, PL: žvakost, žywokost			kidney diseases	1	
					diuretic	1	
					kidney diseases	1	
					joint pain	1	
					limb pain	1	
					allergy	1	
					gout	1	
					radiculitis	1	
					rheumatic pains	1	
					for everything	1	
		cold	1				
		healthy	16	11			
		helpful	1				
		radiculitis	1				
		for everything	1				
		tea	1				
		whisked in the sauna	1				
		twigs with leaves	1				
		twigs with leaves	1				
		tea	1				
		tea	1				
		bones cohesion	1				
		adhesion of bones	2				
		topical application	1				
		alcohol maceration, topical application	1				
		alcoholic infusion	1				
		adhesion of bones	1				
		bones cohesion	1				
		gastric ulcer	1				
		joint pain	2	2			

Table 2 (continued)

Family	Latin name; voucher number	Local name(s)	Used part(s)	Preparation	Medicinal use	DUR	
						LT	PL
Brassicaceae	<i>Armoracia rusticana</i> G. Gaertn., B. Mey. & Scherb.; DZULT022 <i>Brassica oleracea</i> L.	PL: chren LT: kopūstas; PL: kapusta	roots	alcoholic infusion	joint pain	limb pain	2
						adhesion of bones	1
						bones cohesion	1
						adhesion of bones	1
						stomach ache	1
						adhesion of bones	1
						adhesion of bones	1
						bones cohesion	1
						adhesion of bones	1
						topical application	1
<i>Capsella bursa-pastoris</i> (L.) Medik.; DZULT024	LT: triskiautė žvakidė, žvaginė; PL: pastušja sumka	leaves	fresh, topical application	abscesses	abscesses	1	
					headache	1	
					joint pain	2	
					knee ache	1	
					mastitis	1	
					stomach ache	1	
					women's diseases	2	
					organism cleansing	1	
					stomach ache	1	
					decoction	1	
Campanulaceae	<i>Campanula</i> sp.; DZULT092	LT: skambalukai	roots	tea	healthy	1	
						whisked in the sauna	1
Cannabaceae	<i>Cannabis sativa</i> L.; DZULT102	LT: kanapė	aerial parts	tea	for heart	1	
						tincture	1
Caprifoliaceae	<i>Valeriana officinalis</i> L.; DZULT054	LT: valerijonas; PL: valerjonas, valerjana, valerjanka, walerjana	aerial parts	decoction	heart problems	1	
						sedative	2
						for heart	1
						hypertension	1
						sedative	3
						to calm the heart	1
						for heart	1
						heart problems	1
						sedative	1
						sedative	3

Table 2 (continued)

Family	Latin name; voucher number	Local name(s)	Used part(s)	Preparation	Medicinal use	DUR	
						LT	PL
Caryophyllaceae	<i>Silene latifolia</i> subsp. <i>alba</i> (Mill.) Greuter & Burdet; DZULT071, DDZULT39	LT: ochvatas, ochvatnikas	whole plant with roots	tea	for heart		1
			aerial parts	decoction, bath	skin diseases		1
			aerial parts	tea	skin diseases		1
	<i>Silene vulgaris</i> (Moench) Garcke; DD-ZULT21, DZULT115	LT: ochvatas, plaukštūnai; PL: ochvatnik	aerial parts	decoction, bath	allergy		1
				tea	allergy		1
					bladder inflammation		1
					kidney diseases		1
			flowers		urinary bladder		1
			flowers	tea	urinary inflammation		1
		LT: žliugė	aerial parts	compress	external pain		1
Celastraceae	<i>Stellaria media</i> (L.) Vill	LT: [forgotten name]	whole plant with roots	tea	for heart		1
Commelinaceae	<i>Callisia fragrans</i> (L.indl.) Woodson	PL: zalty wus, zolotoj us	aerial parts	alcoholic infusion	joint pain		1
					lung diseases		1
			leaves	alcohol maceration, topical application	abscesses		1
Crassulaceae	<i>Kalanchoe</i> spp.	LT: paleistuvė, kalankė	leaves	drops	wounds		1
Cucurbitaceae	<i>Cucumis sativus</i> L.	LT: agurkas	fruits	juice	runny nose		1
Cupressaceae	<i>Juniperus communis</i> L.; DZULT001	LT: ėglis, eglėlynas, įieglis, kadagys, kadugys, jalaviec	fruits	eaten fresh	improve blood quality		1
					bronchitis		1
					lung diseases		1
				tea	stomach ache		1
					urinary inflammation		1
			roots	alcoholic infusion	blood vessels		1
			twigs with spikes	decoction, bath	gout		1
					jaundice		1
					rheumatic pains		1
				tea	jaundice		1
				whisked in the sauna	healthy		5
					promotes bloodstream		1
			twigs with leaves	decoction	sleep disturbance in children		1
	<i>Thuja</i> spp.; DZULT012	LT: tuja		decoction, bath	sleep disturbance in children		1

Table 2 (continued)

Family	Latin name; voucher number	Local name(s)	Used part(s)	Preparation	Medicinal use	DUR	
						LT	PL
Cyperaceae	<i>Eriophorum</i> spp.	LT: [forgotten name]	whole plant with roots	tea	facial nerve inflammation		2
Dennstaedtiaceae	<i>Pteridium aquilinum</i> (L.) Kuhn	LT: paparčiai	fronds	decoction, bath	haemorrhoids		1
Elaeagnaceae	<i>Hippophae rhamnoides</i> L.	LT: šaltalknis; PL: ablapicha	fruits	juice	healthy		1
Equisetaceae	<i>Equisetum arvense</i> L.; DDZULT03, DDZULT19	LT: dirvinis asiūklis; PL: chvošč, chwoszcz polewy	aerial parts	tea decoction	stomach ache kidney diseases		1 4
Ericaceae	<i>Arctostaphylos uva-ursi</i> (L.) Spreng	LT: meškauoogės	leaves	tea	urinary bladder		1
	<i>Calluna vulgaris</i> (L.) Hull; DZULT126	LT: viržis	flowers	tea	urinary inflammation healthy		1 1
	<i>Chimaphila umbellata</i> (L.) W.P.C.Barton	LT: parūšainė, parūšainikas, stanaunykas; PL: paruszeniec	aerial parts	alcoholic infusion	hypertension organism cleansing women's diseases		1 2 1
	<i>Rhododendron tomentosum</i> Harmaja; DDZULT30	LT: baūnas, gailius; PL: bahulnik	flowers	tea	urinary bladder women's diseases		1 1
	<i>Vaccinium myrtillus</i> L.; DZULT100	LT: mėlynės, juodos uogos, PL: čarnicy, czarne, czarnice, čarničnik, chernika, czernicznik, mėlynės	aerial parts	decoction, bath	heart problems helpful tonic		1 1 1
			aerial parts	tea	back pain men diseases weight lifting injuries women's diseases		1 1 2 1
			aerial parts	decoction	weightlifting injuries women's diseases		1 1
			aerial parts	decoction, bath	gout		1
			aerial parts	decoction	joint pain radiculitis rheumatic pains		1 1 1
			aerial parts	decoction	helpful		1
				tea	improve vision stomach ache diabetes diarrhoea		1 1 1 1
					for everything		1

Table 2 (continued)

Family	Latin name; voucher number	Local name(s)	Used part(s)	Preparation	Medicinal use	DUR					
						LT	PL				
Lamiaceae	<i>Lamium album</i> L.; DZULT084	LT: baltosios notrelės	aerial parts	alcoholic infusion, to rinse decoction	periodontal disease for everything stomach ache	1	1				
								tea	aids cough for everything herpes stomach ache tonic	1 1 1 1 1 1	
	<i>Leonurus cardiaca</i> L.; DZULT055	LT: sukatzolė; PL: pustyrnik, serdecznik	aerial parts	tea	tea	menstrual disorders women's diseases for heart	1	2 1			
									tea	prophylactics sedative	1 1
	<i>Melissa officinalis</i> L.; DZULT014, DZULT131, DDZULT01	LT: melisa; PL: melisa	flowers	decoction tincture	decoction tincture	heart problems for heart	1	2			
									decoction, bath	sedative	1
	<i>Mentha</i> spp.; DDZULT11 (including <i>Mentha x piperita</i> L.; DZULT021; DZULT043, DZULT067, DZULT090; <i>Mentha spicata</i> L. DZULT046, DZULT068, DZULT085; DZULT087; DZULT113)	LT: senoviška smukialapė mėta; mėta, mėtos, miata, senoviška/ garbiniuotoji/ tikroji mėta; bulgariška mėta; pilkoji mėta; PL: mięta, mėta, miata	leaves	tea	tea	cold for heart headache sedative helpful sedative sleep disturbance bellyache	1	1 1 4 2 1 2 1			
									tea	sedative	1
									tea	helpful sedative	1 1
									tea	sedative	1
<i>Mentha</i> spp.; DDZULT11 (including <i>Mentha x piperita</i> L.; DZULT021; DZULT043, DZULT067, DZULT090; <i>Mentha spicata</i> L. DZULT046, DZULT068, DZULT085; DZULT087; DZULT113)	LT: senoviška smukialapė mėta; mėta, mėtos, miata, senoviška/ garbiniuotoji/ tikroji mėta; bulgariška mėta; pilkoji mėta; PL: mięta, mėta, miata	aerial parts	tea	tea	diarrhoea flatulence for heart for everything headache healthy nausea	1	1 1 1 1 1 1 1				
								tea	diarrhoea	1	
								tea	flatulence for heart	1 1	
								tea	for everything headache healthy	1 1 1	

Table 2 (continued)

Family	Latin name; voucher number	Local name(s)	Used part(s)	Preparation	Medicinal use	DUR	
						LT	PL
					organism cleansing	1	
					prophylactics	1	
					sedative	6	1
					sleep disturbance	1	
				whisked in the sauna	toothache	1	
					healthy	2	
			leaves	decoction	cough	1	
					helpful	1	
					sleep disturbance	1	
					stomach ache	1	
			tea		bellyache	1	
					cold	2	
					cough	1	
					diarrhoea	1	
					fever	1	
					for everything	2	
					healthy	1	
					helpful	1	
			aerial parts	put in pillow	sleep disturbance in children	1	
<i>Nepeta cataria</i> L.; DZULT076		LT: citrininė katažolė		tea	sedative	1	
<i>Origanum vulgare</i> L.; DZULT049, DDZULT09		LT: raudonėlis; PL: majrunas	aerial parts	tea	cold	1	
<i>Prunella vulgaris</i> L.; DZULT006, DZULT040, DZULT042, DDZULT05, DDZULT42		PL: brunelka, harlanački, gorlanka	flowers	decoction	cystitis	1	
<i>Salvia officinalis</i> L.; DZULT086, DZULT088, DZULT101		LT: senoviškas šalavijas; šalavijas	aerial parts	decoction, to rinse	for everything	1	
					sore throat	2	
					sore throat	2	
					bleeding	1	
					tonsillitis	1	
				tea	appetiser	1	
					cold	2	
					headache	1	
					runny nose	1	
					sore throat	1	
					tonsillitis	1	
			stems	decoction, bath	diathesis in children	1	

Table 2 (continued)

Family	Latin name; voucher number	Local name(s)	Used part(s)	Preparation	Medicinal use	DUR	
						LT	PL
	<i>Thymus pulegioides</i> L.; DZULT007	LT: čibreliai	aerial parts	tea	cough	1	
	<i>Thymus serpyllum</i> L.; DZULT026	LT: čibreliai, čibrelis dekaratyvinis, čibrelis pievinis, čibrelis, čiombaras; PL: čibreliai, czambor, čabijec, čamborek, čambor, čamboryk, czamborek	aerial parts	decoction	pneumonia cold	1	1
				decoction, to rinse	healthy sore throat	1	1
				tea	bronchitis	1	
					cold	8	4
					cough	4	3
					for everything	2	
					headache	1	
					healthy	3	
					helpful	1	
					lung diseases	1	
					organism cleansing	3	
					sedative	2	
					sore throat	1	
Linaceae	<i>Linum usitatissimum</i> L.	PL: linas, lon	seeds	decoction	diarrhoea	1	
Lycopodiaceae	<i>Lycopodium clavatum</i> L.; DDZULT28, DZULT124, DDZULT28	LT: dziaraza, pataisai; PL: dzieraza	spores	topical application	duodenal ulcer burns	1	1
					contact dermatitis	1	
					diaper rash	11	1
					skin diseases	1	
					wounds	2	
Malvaceae	<i>Malva neglecta</i> Wall.	LT: [forgotten name]	aerial parts	alcohol mac- eration, topical application	limb pain	1	
				alcoholic infusion	healthy	1	
				decoction	cold	4	
					cough	2	
					fever	1	
				tea	cold	10	5
					cough	7	4
					fever	2	2
	<i>Tilia cordata</i> Mill.; DZULT031, DDZULT10, DDZULT14	LT: liepa, liepžiedžiai (part used); PL: lipa, liepžiedžiai, lipovy cviet (part used)	inflorescences				

Table 2 (continued)

Family	Latin name; voucher number	Local name(s)	Used part(s)	Preparation	Medicinal use	DUR					
						LT	PL				
Menyanthaceae	<i>Menyanthes trifoliata</i> L.	LT: puplaiškis; PL: bobek	leaves	whisked in the sauna alcohol maceration	lung diseases	1	1				
								tea	lung diseases	1	1
Oleaceae	<i>Fraxinus excelsior</i> L.; DZULT093 <i>Syringa vulgaris</i> L.; DZULT023	LT: uosis PL: alyvos, bez	whole plant with roots bark flowers	decoction, bath alcohol maceration, topical application alcoholic infusion	allergy joint pain rheumatic pains	1 1 1	1 1 1				
								tea	joint pain	1	1
Onagraceae	<i>Epilobium angustifolium</i> L.; DZULT074, DZULT082, DDZULT23, DDZULT24, DDZULT25, DDZULT35	LT: ivan chai; PL: kiprej, iwan-czaj	aerial parts flowers	tea tea	healthy prostate problems	1 1	1 1				
Orobanchaceae	<i>Euphrasia officinalis</i> L.	LT: akšivietė	aerial parts	decoction, bath	improve vision	1	1				
Paeoniaceae	<i>Paeonia</i> spp.; DZULT015, DZULT056, DDZULT32	LT: bijūnas, pivonija; PL: pievonija, pion, pivonai, piwonija, bijūnas, pion	flowers	boiled with milk decoction	fright fright	2 3	2 3				
Papaveraceae	<i>Chelidonium majus</i> L.; DZULT035	LT: čistatiel, geltončius, karpažolė, ugniažolė, čistatiela, ugniažolė; PL: cystacief, czystacief, ugniažolė	aerial parts	decoction, bath	sedative for children sleep disturbance fright evil eye fright heart problems sedative sleep disturbance allergy	1 1 2 1 3 1 1 2 1	1 1 2 1 7 1 1 2 1				
								fumigation	sleep disturbance	1	2
								tea	evil eye	1	1
									fright	3	7
									heart problems	1	1
									sedative	1	1
									sleep disturbance	2	2
									allergy	1	1
									diaper rash	1	1
									diathesis in children	1	1

Table 2 (continued)

Family	Latin name; voucher number	Local name(s)	Used part(s)	Preparation	Medicinal use	DUR	
						LT	PL
				tea	diaper rash	1	
			latex	topical application	warts	1	
					diaper rash	1	
					psoriasis	1	
					warts	12	6
					wounds	1	
			seeds	cloth-wrapped infusion	sleep disturbance in children	1	
		LT: aguona; PL: mak		decoction	sleep disturbance in children	1	
	<i>Papaver rhoeas</i> L.			poppy milk	sedative for children	2	
			cones	tea	lung diseases	1	
			resin	ointment	abscesses	1	
					adhesion of bones	1	
					wounds	1	
				topical application	wounds	1	
			shoots with needles	decoction, bath	gout	1	
	<i>Pinus sylvestris</i> L.; DZULT051, DDZULT02, DDZULT15	LT: pušis, pušų kankorėžėliai (part used); PL: sasna, sosna, pušis			radiculitis	1	
					rheumatic pains	1	
					allergy	1	
				alcoholic infusion	cough	1	
			cones	alcohol infusion	for everything	1	
					bronchitis	1	
					cough	2	
				decoction	bronchitis	1	
			resin	ointment	burns	1	
			leaves	decoction	stomach ache	1	
				tea	diarrhoea	1	
					headache	1	
					stomach ache	2	
				topical application	abscesses	2	1
					bleeding	3	1
					cuts	1	1
					knee ache	1	1
Plantaginaceae	<i>Plantago major</i> L.; DZULT004	LT: gyšlotis, padarožnik; PL: babka, podorožnik, babok, padarožnik					

Table 2 (continued)

Family	Latin name; voucher number	Local name(s)	Used part(s)	Preparation	Medicinal use	DUR	
						LT	PL
Poaceae	<i>Veronica</i> sp.; DDZULT33	LT: veronika	aerial parts	tea	wounds	1	3
	<i>Avena sativa</i> L.	PL: avizos, owios	aerial parts	decoction, bath	fright allergy	1	1
	<i>Elymus repens</i> (L.) Gould	LT: varputis	aerial parts	decoction	diathesis in children sore throat	1	1
	<i>Hordeum vulgare</i> L.	PL: jačmierń	seeds	decoction, bath	kidney stones	1	1
	<i>Secale cereale</i> L.	PL: roż, żyta	aerial parts	decoction	organism cleansing	1	1
			seeds	topical application	diathesis in children	1	1
Polemoniaceae	<i>Triticum aestivum</i> L.	PL: ruń	aerial parts	decoction, bath	helpful	1	1
	<i>Polemonium caeruleum</i> L.	PL: rabinka	aerial parts	tea	erysipelas	1	1
Polygonaceae	<i>Fagopyrum esculentum</i> Moench; DZULT130	LT: grikliai; PL: hrečka	flowers	decoction	diathesis in children	1	1
	<i>Polygonum aviculare</i> L.; DZULT044, DZULT123	LT: takażole; PL: pćiczny gorec	seeds aerial parts	decoction, bath tea	diaper rash inflammation processes	1	1
	<i>Reynoutria japonica</i> Houtt.; DZULT096	LT: rasvėratrollis	flowers	topical application tea	men's diseases urine problems bleeding	1	2
Primulaceae					antioxidant	1	1
	<i>Primula veris</i> L.	LT: rakteliai	flowers	tea	flu	1	1
			twigs with leaves	whisked in the sauna	flu	1	1
			flowers	tea	healthy	1	1
					bronchitis	1	1
					cold	1	1
					flu	1	1
					pneumonia	1	1
Rhamnaceae	<i>Frangula alnus</i> Mill.	PL: kruszyna	bark	decoction	diarrhoea	1	1
Rosaceae	<i>Alchemilla vulgaris</i> L.	LT: rasakila, rsakila	aerial parts	decoction, bath tea	skin diseases organism cleansing	1	1
	<i>Argentina anserina</i> (L.) Rydb.; DZULT072	LT: sidabrażolė pievinė	leaves	decoction	women's diseases	1	1
	<i>Aronia melanocarpa</i> (Michx.) Elliott	PL: Ćornaja řabina	fruits	compress	diarrhoea	1	1
				decoction, bath	diathesis in children	1	1

Table 2 (continued)

Family	Latin name; voucher number	Local name(s)	Used part(s)	Preparation	Medicinal use	DUR	
						LT	PL
	<i>Crataegus</i> spp.; DZULT095	LT: gudobelė, gudabelė; PL: gluk, bajarysznik	flowers	eaten fresh	for heart	1	1
	<i>Filipendula ulmaria</i> (L.) Maxim.; DZULT128, DDZULT141	LT: aspirinas, vingiorykštė, ingražolė	fruits	tincture eaten fresh tincture tea	for heart for heart for heart analgetic	1	1
	<i>Fragaria vesca</i> L.; DZULT025, DZULT037	LT: žemuogė, žemuogės; PL: poziomki, žemuogės	fruits	tea	cough for heart for everything headache heart problems antibiotic fever vitamin source antibiotic headache healthy kidney diseases cold cough fever	1	1
	<i>Fragaria viridis</i> Weston <i>Malus sylvestris</i> (L.) Mill.	PL: klubnika LT: laukinė obelis	leaves	tea	antibiotic headache healthy kidney diseases cold cough fever	1	1
	<i>Potentilla erecta</i> (L.) Raeusch.; DZULT072	LT: sidabražolė miškinė, sidabražolė	calyx flowers	decoction tea	cold cough fever	1	1
	<i>Prunus cerasus</i> L.	PL: višnia	aerial parts roots fruits	tea tea alcoholic infusion tea	diarrhoea diarrhoea hypertension for everything	1	2
	<i>Prunus padus</i> L.; DZULT122	PL: leva, czaromucha	flowers	alcohol mac- eration, topical application tea	joint pain rheumatic pains	1	1
	<i>Prunus domestica</i> L.	LT: sliva	fruits	tea eaten fresh	joint pain constipation	1	1

Table 2 (continued)

Family	Latin name; voucher number	Local name(s)	Used part(s)	Preparation	Medicinal use	DUR	
						LT	PL
	<i>Pyrus pyraeaster</i> (L.) Burgsd	LT: laukinė kriaušė	flowers	tea	cough	1	
	<i>Rosa</i> spp.	PL: szypownik	fruits	tea	fever cold cough for everything helpful for everything	1 1 1 1 1 1	
	<i>Rubus idaeus</i> L.; DZULT028, DDZULT36, DZULT107	LT: avietė, avietės, malina; PL: avietės, malina, maliníniak, malinínik, maliny	leaves stems (with leaves)	tea decoction	cold fever pneumonia cough breathlessness cold inflammation processes bronchitis cold cough inflammation processes flu sore throat healthy	4 1 1 2 1 1 1 1 1 6 3 1 1 1	15
			fruits	whisked in the sauna alcoholic infusion	cold flu healthy cold cold helpful cold flu cold cough flu healthy sore throat	2 1 1 3 1 1 3 1 1 5 3 1 1	2
			fruits	eaten fresh jam juice raw jam tea	healthy cold cold helpful cold flu cold cough flu healthy sore throat	1 4 1 3 1 1 3 1 1 5 3 1 1	1

Table 2 (continued)

Family	Latin name; voucher number	Local name(s)	Used part(s)	Preparation	Medicinal use	DUR	
						LT	PL
	<i>Sorbus aucuparia</i> L.; DZULT009	LT: šermukšnis; PL: rabina, rjabina, šermukšnis	leaves	tea	cold fever flu diuretic fever sweating flu healthy vitamin source vitamin source flu	3 1 1 1 1 1 1 1 1 1	1 1 1 1 1 1 1 1 1 1
Rutaceae	<i>Ruta graveolens</i> L.	LT: rūta		jam	for heart		1
Salicaceae	<i>Populus tremula</i> L.; DZULT136	LT: topolis; PL: topal, topol, topolis	aerial parts buds	oil infusion syrup tea tea alcohol mac- eration, topical application	healthy healthy healthy for heart for everything for heart bleeding	1 1 1 1 1 1 1 1	1 1 1 1 1 1 1 1
				alcoholic infusion	pityriasis rosea wounds cuts gastric ulcer jaundice joint pain liver diseases rheumatic pains wounds diarrhoea antiseptic	1 1 1 1 1 1 1 1 1 1 1 1	1 2 1 1 1 3 1 1 1 1 1 1
	<i>Salix</i> spp.; DZULT008, DZULT011, DZULT061, DZULT133	LT: karklas, žilvitis, verba; PL: wierzba	leaves catkins	decoction topical application eaten fresh	liquefies blood		1

Table 2 (continued)

Family	Latin name; voucher number	Local name(s)	Used part(s)	Preparation	Medicinal use	DUR	
						LT	PL
Sapindaceae	<i>Acer platanoides</i> L.; DZULT029, DZULT062	LT: klijanas, klevas	twigs	fumigation	headache	1	
				tea	headache	1	
				fumigation	liquefies blood	1	
				fumigation	evil eye	2	
Sapindaceae	<i>Aesculus hippocastanum</i> L.; DZULT034, DZULT057	PL: kaštan, kaštonas, kasztan	twigs with leaves	whisked in the sauna	fright	1	
				flowers	healthy	1	
				flowers	varicose veins	2	
				flowers	varicose veins	2	
Sapindaceae			fruits	alcohol mac-eration, topical application	joint pain	4	
				alcoholic infusion	joint pain	4	
				alcohol mac-eration, topical application	toothache	1	
				alcohol mac-eration, topical application	varicose veins	2	
Scrophulariaceae	<i>Verbascum nigrum</i> L., DZULT117	LT: daugiažiedė, šimtažiedė	flowers	alcoholic infusion	joint pain	5	
				compress	joint pain	1	
Solanaceae	<i>Datura stramonium</i> L.; DZULT069, DZULT098, DZULT135	LT: durnadagis, durnapjėnas, durmaropė, durnoj ropė; PL: durnaropė, durnapjėnas	aerial parts	tea	sore throat	1	
				alcohol mac-eration, topical application	external pain	1	
Solanaceae	<i>Solanum tuberosum</i> L.; DZULT032	LT: bulvė; PL: bulba, kartofle, kartoška	leaves	fumigation	fright	2	
				fumigation	fright	3	
				tea	sedative	1	
				eaten fresh	sedative	1	
				tea	fright	1	
				seeds	sedative	1	
				fumigation	fright	1	
				tea	fright	1	
				compress	sedative	1	
				compress	abscesses	1	
Solanaceae			tubers	eye inflammation	eye inflammation	1	
				sore throat	sore throat	1	
				cold	cold	2	
				cough	cough	1	
				sore throat	sore throat	2	
Solanaceae			juice	gastric ulcer	gastric ulcer	1	
				heart rate	heart rate	1	

Table 2 (continued)

Family	Latin name; voucher number	Local name(s)	Used part(s)	Preparation	Medicinal use	DUR	
						LT	PL
Theaceae	<i>Camellia sinensis</i> (L.) Kuntze	LT: juoda arbata	leaves	compress	eye inflammation		1
Thymelaeaceae	<i>Daphne mezereum</i> L.	LT: vilkaugė, vilkaugis	aerial parts	topical application	toothache		1
Urticaceae	<i>Urtica dioica</i> L.; DZULT002, DZULT017	LT: dilgėlė, dilgėlis, dilginės; PL: krapawa, pokrzywa	aerial parts	decoction, bath	hair care		1
				tea	acne		1
					blood coagulation		1
					improve blood quality		1
					organism cleansing		1
					stomach ache		1
				topical application	joint pain		1
				whisked in the sauna	healthy		1
			leaves	decoction	healthy		1
				tea	blood vessels		1
					cough		1
					varicose veins		1
			aerial parts in spring	decoction, bath	vitamin source		1
				eaten cooked	healthy		1
				eaten fresh	vitamin source		1
			aerial parts	decoction	healthy		1
		PL: žyčka		topical application	wounds		1
			fruits	syrup with honey	tonic		1
			flowers	tea	diuretic		1
					heart rate		1
					sweating		1
			fruits	alcoholic infusion	heart problems		1
					hypertension		1
				decoction	hypertension		1
				eaten fresh	healthy		1
				juice	hypertension		4
					hypertension		2
					intestinal pain		3
				raw jam	diabetes		2
					for heart		1
Viburnaceae	<i>Sambucus nigra</i> L.; DZULT081 <i>Viburnum opulus</i> L.; DZULT010	LT: juodasis šėivamedis LT: kalina, putinas; PL: kalina, putinas	fruits				

Table 2 (continued)

Family	Latin name; voucher number	Local name(s)	Used part(s)	Preparation	Medicinal use	DUR	
						LT	PL
Violaceae	<i>Viola tricolor</i> L.	LT: laukiniai broliukai, širdininkai	aerial parts	decoction, bath tea	hypertension	6	1
					hypertension	1	1
					varicose veins	1	1
					healthy	1	1
					hypertension	1	1
					vitamin source	1	1
					diaper rash	1	1
					cancer	1	1
					diaper rash for heart	1	1
					for everything	1	1

taxa cited, as it was essentially the same across different schooling levels (Spearman's $\rho = -0.018$, $p = 0.885$).

Marked differences appeared between LT and PL in several medicinal categories, especially digestive and general/unspecified. LT reported more taxa for general/unspecified uses (55 vs 44) and more digestive remedies (30 vs 24). By contrast, musculoskeletal, neurological, urological, and cardiovascular conditions showed similar proportions in both groups. As shown in Fig. 4 (lighter yellow-green shades), etc categories such as ear, eye, and reproductive health (including pregnancy, childbirth, and family planning) contained the fewest taxa. The greater diversity of plants used for general health in LT suggests a broader application of herbal medicine, whereas PL displayed a more balanced spread across categories.

In total, 135 emic categories of medicinal use were documented (PL: 87; LT: 115), while 27 categories that were reported historically are no longer in use. Many of them referred to illnesses remembered by older generations, such as smallpox, diphtheria, scrofula (“gland disease”), jaundice, dysentery, rickets, and worm infections, among others. During the second half of the twentieth century, vaccination and public health measures led to the decline or near disappearance of specific disease categories. As certain diseases faded from everyday discourse, the plant remedies once used to treat them also declined in local practice. Some ailments, including rheumatism, arthritis, hypertension, anaemia, or diabetes, among others, remained relevant but were reframed over time. In the case of anaemia, traditionally it was described as “*weak blood*”, whereas nowadays it is mostly framed by the biomedical definitions circulated in clinics and health literature.

Knowledge hybridisation patterns

Although the dataset is firmly grounded in LEK (A), the most frequent hybrid knowledge combinations were AC (local + popular, $n = 64$), BC (formal + popular, $n = 52$), and ABC (local + formal + popular, $n = 31$) (Fig. 5a). In contrast, formal-only uses occurred in 38 cases and popular-only in 24. The LT group engaged far more with hybridised knowledge, citing 56 cases compared to 22 in the PL group (Fig. 5b). In PL, ABC-type hybridisation was very low, occurring in only two species (*Vaccinium myrtillus* and *Aloe* spp.), whereas in LT, 29 plant species were fully hybridised. This pattern reinforces earlier findings that PL knowledge and practices are less epistemically hybridised. By contrast, 29 fully hybridised species were recorded in the LT community, indicating greater institutional embedding and more active interactions with both the formal and popular knowledge systems.

Some taxa, such as *Hippophae rhamnoides*, displayed diverging trajectories in terms of knowledge hybridisation. It was fully hybridised in the LT community, but it



Fig. 2 Cross-ethnic comparison. **a**) Jaccard index with a proportional Venn diagram for all the medicinal plant taxa reported among the LT and PL communities; illustrative photograph of *Brassica oleracea* in use; **b**) the Jaccard index and a proportional Venn diagram of medicinal taxa mentioned by at least three interviewees, illustrative photograph of *Aloe* spp. in a domestic setting. Photo credits: JP, 2018–2019




Fig. 3 Medicinal uses of wild plants documented during fieldwork, illustrating different use categories and preparation states: **a**) freshly foraged leaves of *Tussilago farfara* (LT); **b**) demonstration of the medicinal use of *Chelidonium majus* (PL); **c**) powdered spores of *Lycopodium clavatum* (LT); **d**) dried buds of *Pinus sylvestris* (PL); **e**) dried aerial parts of *Matricaria chamomilla* (LT); **f**) foraging of *Achillea millefolium* (LT); **g**) dried aerial parts of *Rhododendron tomentosum* (LT); **h**) dried inflorescences of *Helichrysum arenarium* (PL); **i**) dried aerial parts of *Gymnocarpium dryopteris* (PL). Photo credits: JP, 2018–2019




Fig. 4 Comparison of medicinal plant use across ICPC-2 categories among LT and PL ethnic groups. The colour scale ranges from yellow (low values) to dark purple (high values), indicating the relative number of plant taxa used in each category

was not included in the hybrid categories for PL. By contrast, *Acorus calamus* remained non-hybrid, restricted specifically to LEK. Additionally, for *Cucumis sativus*, *Daucus carota*, or *Euphrasia officinalis*, for example, LT interviewees reported hybridised uses drawing on multiple knowledge sources, while PL interviewees reported only single-source uses.

The results point out contrasts between LT and PL in terms of how knowledge is hybridised and the range of sources drawn upon for medicinal plant uses. The Shannon–Wiener diversity index (H') was 0.97 for LT and 0.52 for PL, indicating greater epistemic diversity among LT interviewees. Moreover, when considering the degree of hybridisation across general knowledge origins domains, the LT community showed a markedly higher degree of hybridisation ($HD = 0.195$) compared to PL ($HD = 0.059$). We found a statistically significant difference in the degree of hybridisation between the LT and PL groups, indicating that the knowledge systems differ meaningfully (Student's $t = 3.450$, $p = 0.002$, $n = 67$). This suggests that, despite a shared plant repertoire, patterns of knowledge integration differed. PL showed more internal hybridisation (0.266) than LT (0.106).

A closer look at the plant level reveals more nuanced differences between LT and PL communities. As shown in Fig. 6, among the LT, several culturally and medicinally versatile species (including *Cucumis sativus*, *Daucus carota*, *Equisetum arvense*, *Euphrasia officinalis*, *Hippophae rhamnoides*, *Malva neglecta*, *Mentha × piperita*, *Primula veris*, and *Stellaria media*) attained full hybridisation ($HD = 1$). This suggests that knowledge about these plants is maintained simultaneously through local

practice, scientific reference, and popular culture. By contrast, only two taxa, *Polygonum aviculare* and *Silene vulgaris*, achieved this score in the PL sample, reflecting a narrower set of plants for which knowledge circulates across all three epistemic spheres.

More specifically, when subcategories are considered, the dominant bases are A1 (intergenerational; 533 DUR) and A4 (ancestral; 223). Notable internal A-combinations include A1 + A3 (intergenerational + oblique; 89), A2 (horizontal local transfer; 64), and A1 + A4 (intergenerational + ancestral; 53). Among cross-source hybrids, the most common are B1 + C1 (formal publications + popular media; 32), A1 + B1 (intergenerational + formal publications; 22), A1 + C1 (intergenerational + newspapers/magazines; 20), and A1 + B1 + C1 (intergenerational + formal publications + newspapers/magazines; 20). However, this pattern differs between groups. PL relied comparatively more on combinations linked to formal and professional channels, especially B1 + B2 (formal publications + health professionals; 10) and A3 + B2 (oblique + health professionals; 9). Their profile is thus dominated by local and community-based continuity, with cross-source hybrids occurring only rarely. By contrast, LT demonstrated broader hybridisation, including a higher share of fully integrated ABC cases (LT = 29 vs. PL = 2). LT knowledge patterns also reveal multiple cross-source combinations, for instance, B1 + C1 (formal publications + newspapers/magazines; 24), A1 + B1 (intergenerational + formal publications; 22), A1 + C1 (intergenerational + newspapers/magazines; 17), A1 + B1 + C1 (intergenerational + formal publications + newspapers/magazines; 19), and B1 + B4 + C1 + C3 (formal publications + digital medical




Fig. 5 Distribution of medicinal plant taxa across knowledge origin categories: **a**) Relative frequency of categories in the whole dataset (A – local, B – formal, C – popular, AB – local and formal, AC – local and popular, ABC – local, formal, and popular); **b**) Comparison of category proportions between Lithuanian (blue) and Polish (orange) interviewees; **c**) Taxa associated with different knowledge categories among Lithuanians; **d**) Taxa associated with different knowledge categories among Poles

resources + newspapers/magazines + social media and popular websites; 15), as shown in Fig. 7.

Patterns of hybridisation differed across the emic categories of medicinal plant use, including both disease-related and preventive/functional domains. High knowledge diversity and integration were observed in vitamin supplementation ($H' = 0.64$, $HD = 1.00$), body cleansing ($H' = 1.08$, $HD = 0.75$), bone adhesion ($H' = 1.05$, $HD = 0.60$), blood quality improvement ($H' = 1.33$, $HD = 0.50$), and urinary bladder diseases ($H' = 1.03$, $HD = 0.50$). In contrast, there was little recorded variety and almost no hybridisation in abscesses, diarrhoea, culture-bound syndromes (such as fright or the evil eye), and eye conditions.

Interviewees who mentioned a greater number of plant taxa tended to show higher levels of knowledge hybridisation, as confirmed by a significant Spearman

correlation ($\rho = 0.652$, $p < 0.001$). Specifically, LT participants cited more taxa (mean \pm SD = 28 ± 21.6) than PL (16.38 ± 15.9). The difference was statistically significant (Student’s $t = 2.455$, $p = 0.017$, $n = 67$). Robustness was confirmed with a Wilcoxon rank-sum test ($p < 0.05$). This suggests that both the diversity and hybridity of plant knowledge studied in the borderland area are more pronounced in the LT group.

Sociodemographic correlates of medicinal plant knowledge hybridisation

A significant negative correlation was observed between hybridisation and the age of interviewees (Spearman’s $\rho = -0.38$, $p = 0.001$), indicating that younger interviewees tend to have higher degrees of hybridisation in medicinal use. In contrast, gender showed no significant association with hybridisation (Spearman’s $\rho = -0.17$, $p = 0.185$).



Fig. 6 Cross-ethnic comparison of hybridisation patterns in medicinal plant taxa between the Lithuanian (blue) and Polish (orange) communities. HD (Degree of Hybridisation) quantifies the proportion of use reports combining multiple knowledge sources and ranges from 0 (all use reports derived from single knowledge origin) to 1 (all use reports classified as hybrid)




Fig. 7 Cross-source hybridisation patterns of medicinal plant knowledge among Lithuanians and Poles. The size of the nodes corresponds to the relative importance of knowledge combinations of subcategories, while the thickness of the connecting lines is proportional to the number of detailed use reports (DUR)

Similarly, although a higher educational level was weakly associated with increased hybridisation (Spearman's $\rho=0.23$, $p=0.199$), this relationship was not statistically significant.

Although the correlation between the number of spoken languages and the degree of knowledge hybridisation was weak (Spearman's $\rho=0.23$), it did not reach statistical significance ($p=0.060$). Still, the direction of the association suggests a potential trend worth exploring. This indicates that multilingual individuals may be more likely to access and integrate diverse knowledge domains, although this trend does not reach formal significance in our sample.

Discussion

Our sample generally shows low levels of hybridisation, suggesting that the LT and PL communities along the Belarusian border continue to rely primarily on their local ecological traditions. Nearly 40% of the reported plant-based remedies came from A1-type (intergenerational) sources. In practice, this meant that people most often learnt about plants from their parents or grandparents, underscoring the vital role of kinship in the transmission and translation of ethnomedicinal knowledge. Earlier studies have shown that vertical transmission is key to preserving continuity in plant use [40, 47]. At the borderland site, our interviewees reported that knowledge of medicinal plants continues to circulate within

families and village networks during everyday conversations. One reason for this persistence lies in the social dynamics of learning. Bond & Gaoue [6] found that prestige and homophily significantly predict the structure of social learning networks related to medicinal plant knowledge. That is, individuals tend to learn from those they perceive as similar (e.g., individuals of the same age, gender, or social background) or prestigious, for example, respected elders or educated individuals.

In both the LT and PL groups, ancestral/historical transmission (A4) ranked second among the origins of knowledge. Many described their plant lore as self-evident: “*what has always been there*,” handed down from one generation to the next. Earlier studies on the medicinal use of fungi in the region reached similar conclusions, demonstrating that people view such knowledge as a continuous tradition over time [57]. Such findings point to a dominance of relational worldview: knowledge is not seen as something new but as something carried forward, sustained through trust in local community traditions.

Even with greater hybridisation, grandparents, neighbours, and elders remained important teachers, demonstrating the enduring strength of oblique transmissions. Such practices reveal the resilience of local knowledge systems and their capacity to coexist with, or resist, global knowledge systems, echoing observations by Vandebroek and Balick [72]. Horizontal (A2) and oblique forms of learning (A3), which involve multiple transmitters and



Fig. 8 Interview with Lithuanian herbalist, Elena Targavičienė: her traditional homestead surrounded by a flower garden (top left); framed certificates and photographs testifying to her public recognition (middle left), including an official state award; devotional images and bunches of dried herbs hanging in her home (middle left, lower); and dried mixtures of medicinal plants stored in large quantities (bottom left). Photo credits: JP, 2018

non-parental figures, help maintain coherence within communities while also facilitating adaptation and cultural change [60]. From this perspective, hybridisation can be seen less as a rupture than as a reframing process, since continuity itself provides the basis for renewal and invention.

In the LT case, by contrast, herbal specialists (A3) appear more strongly as mediators of cross-source knowledge, complementing the broader set of hybrids based on A1 (intergenerational). The role of herbal specialists was exemplified by Elena Targavičienė (Fig. 8), a widely respected herbalist in the examined area. We had the possibility to interview her not long before she passed away. When we returned the following year, her daughter welcomed us and continued the conversation, clarifying several moments on plant use. This sequence illustrates the moment of generational transition: while Targavičienė embodied oblique transmission as a community expert, her daughter represents vertical transmission, preserving memory and ethnobotanical practices. Taken together, these encounters demonstrate that the specialist's authority did not end with her life but persisted in her daughters' memories and within local community, which continues to recall her as a leading expert in plant knowledge with high-level expertise dating back to pre-WWII times. These findings highlight the continuing importance of oblique transmission, where (local) expertise outside the family line actively reshapes modern plant knowledge [37].

Popular knowledge (C) was rarely mentioned as an independent source, whether in the form of herb columns in newspapers, calendars or advice shared on social

media. Interviewees often tied popular sources to what they already knew from family or formal channels: “We saw it in the calendar, but we already did this at home” (Polish, female, 45 years old), or “Facebook only repeats what my grandmother used to say” (Lithuanian, female, 52 years old). Badke et al. [3] observed a similar pattern: plants that have already received formal validation typically receive the most discussion in local newspapers and magazines. A comparable process was observed in the re-narrativisation of *Epilobium angustifolium* in Eastern Europe, where media outlets, newspapers, magazines, and later online platforms amplified rather than originated medicinal narratives [53]. In that case, traditional motifs were blended with selectively cited scientific findings, and their active circulation through social media helped shape the plant's modern valuation.

In our dataset, formal knowledge (B) was comparatively marginal and rarely cited as a single source. When it first appeared, it was frequently combined with LEK (e.g., A1 + B1 intergenerational + formal publications), suggesting that integration occurs primarily through informal, practice-based hybridisation rather than the formal replacement or displacement of community knowledge. These patterns align with Tengö et al.'s [68] multiple-evidence-based approach, which emphasises dialogic involvement as a step towards co-production rather than hierarchy across knowledge systems.

Participants in our interviews sometimes expressed a need to discover scientific confirmation for the LEK they possess, vigorously combing the knowledge from academic publications to verify or support long-standing community practices. This behaviour pattern exemplifies

bottom-up epistemic negotiation, where local knowledge holders seek convergence with academic science, not out of deference but in acknowledgement of its legitimate function [29]. Our findings echo Ziman & Ziman's [79] observation: people often mentioned a doctor's advice, a pharmacy leaflet, or a medicinal book not as something new, but as confirmation of remedies they already used and trusted. Our data extend the argument made by Frazão-Moreira et al. [18] that written texts are themselves a vehicle of LEK. This highlights the inherently hybrid nature of contemporary ethnobotanical knowledge systems.

Health professionals (B2) (doctors, nurses, and local pharmacists) were often described as reliable sources of knowledge about medicinal plants. Yet their authority was not presented as purely formal; it was embedded in relationships and everyday practice. Several people explained that they first heard a recommendation from a doctor or pharmacist and then added, with a kind of reassurance, that the same remedy was "also known in our village" or "something we have always done". This layering creates what we might call reverse legitimisation: biomedical voices end up validating long-standing community knowledge rather than replacing it [77]. Biomedical authority is selectively incorporated more into local knowledge of the PL community. In the PL case, combinations such as B1 + B2 (formal publications + health professionals) and A3 + B2 (oblique + health professionals) indicate that biomedical practitioners and non-kin experts remain important transmitters, even if overall hybridisation is less pronounced. Instead of challenging LEK, health professionals seemed to reinforce it, offering a subtle endorsement that allowed hybrid systems to remain continuous while also adapting.

We found that hybridisation is a selective process, primarily occurring in relation to plant species with disputed or complex medicinal applications, rather than representing a universal trait of medicinal plant knowledge. Accordingly, hybridity should not be viewed as inherently positive or negative; rather, its effects are context-dependent: while local knowledge can be displaced and marginalised through formal and popular knowledge in processes of "epistemic homogenisation" [49], our findings also show how local knowledge can become legitimised and potentially safeguarded through processes of hybridisation.

The highest levels of hybridisation were found in areas such as blood coagulation, diabetes, and anaemia, as well as in the few cancer-related uses reported, where biomedical and local explanatory models sometimes converged. Emic categories, such as vitamin supplementation and detoxification, are relatively new health narratives that have been widely disseminated by formal and popular sources [43] and are becoming increasingly

integrated into LEK through formal health channels and media exposure. By contrast, conditions rooted in culture, such as fear, the evil eye, or weightlifting injuries, were addressed almost exclusively within the LEK domain, with only rare points of contact with other systems. Interviewees sometimes combined traditional and biomedical framing within the same person's repertoire. In one interview, a 44-year-old Lithuanian woman described *Artemisia vulgaris* as a plant that "warms the belly", while also noting that it "supports the gastrointestinal system through antispasmodic effects".

Pieroni et al. [51] reported that during the COVID-19 pandemic in the Vilnius region, media narratives and biomedical discourse amplified the traditional use of medicinal plants for immune support, respiratory health, and overall well-being. In such critical moments, popular and academic perspectives did not displace LEK. Instead, they reframed and reinforced it, creating new domains for hybrid health practices.

Our analysis revealed a positive association between the number of medicinal plant taxa reported by interviewees and the degree of knowledge hybridisation. Interviewees with broader botanical repertoires are more likely to expand, support or update their knowledges through new inputs. This co-occurrence reflects a form of resilience through diversification frameworks [15]. In parallel to ecosystems where multiple species contribute to the same function with varying responses to disturbance, our data suggest that individuals drawing on diverse medicinal plant taxa and multiple knowledge sources embody a biocultural form of response diversity: diversified knowledge portfolios may support adaptive capacity under change. On the plant level, the likelihood of knowledge hybridisation appears to be linked to both the therapeutic domain and the circulation of knowledge. Taxa associated with biomedical discourse and supplementation (e.g., *Symphytum officinale*, *Thymus serpyllum*, *Matricaria chamomilla*) were frequently hybridised, while plants tied to culturally specific or marginal uses (e.g., *Acorus calamus*, *Tilia cordata*) remained within LEK. Widely cultivated or commonly encountered in daily life, these plants (e.g., *Cucumis sativus*, *Daucus carota*, or *Aloe* spp.) tend to attract knowledge through multiple channels, reflecting their everyday accessibility. These patterns underscore how therapeutic relevance, visibility, and institutional embedding influence the degree of hybridisation among groups.

According to our findings, age also significantly influences the degree of hybridisation, even though earlier ethnobotanical research has demonstrated that elders have a more comprehensive and varied understanding of medicinal plants and rely on traditional health care more often than younger people [4, 65]. Younger interviewees demonstrated greater levels of integration with digital

media discourses, whereas older interviewees generally retained medicinal plant knowledge based on historical and intergenerational transmission. Our results also support previous findings that the increasing influence of digital media on how people perceive and use herbal remedies is evident nowadays [1]. This suggests that age predicts both relative epistemic flexibility and receptiveness to hybrid knowledge forms, as well as the depth of knowledge within LEK.

Multilingual participants from both ethnic groups explained that they rely on formal and popular sources available in the different languages they use. Monolingual speakers, by contrast, tended to pass remedies only within their families. This difference is particularly noticeable in the borderlands, where plant names, recipes, and advice circulate in several languages simultaneously [2]. In such environments, language serves as more than just a means of communication, it can either unlock or block access to knowledge [14]. In our case, local knowledge has historically circulated in overlapping Lithuanian, Polish, Belarusian, and Russian registers, giving multilinguals a broader repertoire of sources. Earlier work has shown that when people use more than one language, they are forming their ethnobotanical practices from different traditions. It, in turn, enriches the diversity of plants they keep in use within their local communities [34].

These patterns suggest that hybridisation operates not merely as a measure of *what is known* but as a reflection of how individuals are situated to access, interpret, and integrate diverse epistemic traditions. While minimal hybridisation implies the strength of intergenerational transmission and the integrity of locally rooted knowledge systems, higher degrees of hybridisation suggest adaptability, openness, and epistemic flexibility within the community.

Combining knowledge too quickly or without a foundation in lived practice puts it at risk of losing coherence and reliability. It also comes with a risk of marginalising local knowledge, especially in the younger generation who are increasingly immersed in a rapidly digitalising world. Yet when hybridisation grows out of dialogue and everyday practice, it can add diversity without erasing what makes local traditions specific. From this perspective, knowledge hybridisation is not an event with uniform effects on local knowledge but rather a process shaped in diverse ways by culture, community values, and broader socio-ecological conditions.

Conclusions

Our results underline the need for sociocultural positioning and historical exposure to institutional frameworks, such as the language of education and written sources, in shaping the content and diversity of hybrid knowledge

systems within culturally adjacent yet separate communities. Taken together, our findings show that the consequences of knowledge hybridisation are ambivalent and context-dependent, requiring conscious, situated approaches rather than generic promotion.

The study highlights that, especially in light of environmental and sociopolitical changes, hybridity in LEK cannot be understood as intrinsically beneficial or detrimental. Rather, its effects are contextually shaped: on the one hand, formal and popular knowledge can displace local knowledge, marginalising or eroding community practices; on the other hand, they may also safeguard or revitalise it, for instance, when local knowledge is legitimised through formal science or amplified through digital media. Conscious approaches to hybridity, therefore, require attentiveness to both opportunities (e.g., safeguarding or revitalising local knowledge) and risks (e.g., displacement, marginalisation, or the erosion of autonomy).

Our findings suggest that hybrid systems draw on a wider range of plant knowledge sources and medicinal uses, lending credence to the view that knowledge hybridity might be favourably linked with ethnobotanical diversity. Hybridisation, therefore, appears as both a cultural and cognitive tool that enables people to expand and modify their foundational knowledge of medicinal plants.

From the perspective of its origins, contemporary medicinal plant knowledge emerges as fluid, continually reshaping what is learnt at home, transmitted through formal channels, and circulated via popular media. It is not a static inheritance but a living practice, dynamically maintained through continual acts of integration and adaptation. We foresee that a more comprehensive understanding of the dynamics and complexity of hybrid knowledge systems will support the creation of inclusive, flexible, and resilient biocultural frameworks for conservation and transformative change.

Abbreviations

LT	Lithuanians
PL	Poles
LEK	Local Ecological Knowledge
DUR	Detailed Use Report
ICPC-2	International Classification of Primary Care
WWII	Second World War
COVID-19	Coronavirus Disease 2019
HD	Degree of Hybridisation

Acknowledgements

Our warmest gratitude goes to all of the study participants, who patiently shared their knowledge of medicinal plants and provided outstanding support. We are thankful to our Latvian colleagues, Baiba Prūse, Andra Šimanova, and Ieva Mežaka, for their kind assistance during fieldwork. We also want to thank Pietro Daniel Omodeo, cultural historian of science and professor of historical epistemology at the Department of Philosophy and Cultural Heritage, Ca' Foscari University of Venice, Italy, for his generous feedback, intellectual encouragement, and thoughtful discussions that helped shape the direction of this study.

Author contributions

J.P. and R.S. conceived and designed the research. J.P. and P.Š. collected the data, including conducting interviews and collecting voucher specimens. J.P., L.C., D.L., and R.S. contributed materials/analysis tools. J.P. drafted the manuscript. All authors contributed to the writing and gave final approval for publication.

Funding

This work was supported by the European Research Council (grant agreement N° 714874) under the European Union's Horizon 2020 research and innovation initiative.

Data availability

Data are available upon request from the corresponding authors.

Declarations

Ethics approval and consent to participate

Ethical approval was granted by the Ca' Foscari University of Venice Ethics Committee. We strictly followed the ethical guidelines prescribed by the International Society of Ethnobiology.

Consent for publication

Not applicable.

Competing interests

The authors declare no competing interests.

Author details

¹Department of Environmental Sciences, Informatics and Statistics, Ca' Foscari University of Venice, Venice, Italy

²NICHE Centre for Environmental Humanities, Ca' Foscari University of Venice, Venice, Italy

³Knowledge, Technology and Innovation Group (KTI), Wageningen University, Wageningen, the Netherlands

⁴Botanical Garden of Vilnius University, Vilnius, Lithuania

⁵Estonian Literary Museum, Tartu, Estonia

Received: 30 September 2025 / Accepted: 25 December 2025

Published online: 16 February 2026

References

- Anez, S. G., Burkhart, E. P., Kellogg, J. J. (2025). Ghost pipe then and now: the influence of digital media on the medicinal use of *Monotropa uniflora* in the United States. *Economic Botany*, 1–18.
- Aziz MA, Hassan M, Ullah A, Ullah Z, Sökand R, Pieroni A. Keeping their own and integrating the other: medicinal plant use among Ormurs and Pathans in South Waziristan, Pakistan. *J Ethnobiol Ethnomed*. 2023;19(1):62.
- Badke MR, Budó MDLD, Alvim NAT, Zanetti GD, Heisler EV. Popular knowledge and practices regarding healthcare using medicinal plants. *Texto Contexto Enferm*. 2012;21:363–70.
- Begossi A, Hanazaki N, Tamashiro JY. Medicinal plants in the Atlantic Forest (Brazil): knowledge, use, and conservation. *Hum Ecol*. 2002;30(3):281–99.
- Bexultanova G, Prakofjewa J, Sartori M, Kalle R, Pieroni A, Sökand R. Promotion of wild food plant use diversity in the Soviet Union, 1922–1991. *Plants*. 2022;11(20):2670.
- Bond MO, Gaoue OG. Prestige and homophily predict network structure for social learning of medicinal plant knowledge. *PLoS One*. 2020;15(10):e0239345.
- Brondizio ES, Aumeeruddy-Thomas Y, Bates P, Carino J, Fernández-Llamazares Á, Ferrari MF, et al. Locally based, regionally manifested, and globally relevant: indigenous and local knowledge, values, and practices for nature. *Annu Rev Environ Resour*. 2021;46(1):481–509.
- Budryte D. Taming nationalism? Political community building in the post-Soviet Baltic states. Routledge; 2017.
- Conroy MS. The Soviet pharmaceutical industry and dispensing, 1945–1953. *Eur Asia Stud*. 2004;56(7):963–91.
- Congretel M, Pinton F. Local knowledge, know-how and knowledge mobilized in a globalized world: a new approach of indigenous local ecological knowledge. *People Nat*. 2020;2(3):527–43.
- Conroy MS. The Soviet Pharmaceutical Business during its first two decades (1917–1937), vol. Vol. 202. Peter Lang; 2006.
- Dauber, D., Fink, G. (2011). Hybridisation in social systems: A conceptual perspective. Available at SSRN 1748365.
- Delannoy, L., Leveugle, J. C., Maniatakou, S., Jørgensen, P. S. (2025). More than a buzzword? Mapping interpretations of the 'polycrisis'.
- Droz L, Brugnach M, Pascual U. Multilingualism for pluralising knowledge and decision making about people and nature relationships. *People Nat*. 2023;5(3):874–84.
- Elmqvist T, Folke C, Nyström M, Peterson G, Bengtsson J, Walker B, et al. Response diversity, ecosystem change, and resilience. *Front Ecol Environ*. 2003;1(9):488–94.
- Fernández-Llamazares Á, Lepofsky D, Lertzman K, Armstrong CG, Brondizio ES, Gavin MC, et al. Scientists' warning to humanity on threats to indigenous and local knowledge systems. *J Ethnobiol*. 2021;41(2):144–69.
- Fontefrancesco MF, Pieroni A. Renegotiating situativity: transformations of local herbal knowledge in a Western Alpine valley during the past 40 years. *J Ethnobiol Ethnomed*. 2020;16(1):58.
- Frazão-Moreira A., Carvalho A. M., Martins M. E. (2009). Local ecological knowledge also 'comes from books': cultural change, landscape transformation and conservation of biodiversity in two protected areas in Portugal. *Anthropological Notebooks*, 15(1).
- Galvonaite A, Valiukas D, Kilpys J, Kitrienė Z, Misiūnienė M. Climate atlas of Lithuania. Vilnius: Lithuanian Hydrometeorological Service under Ministry of Environment; 2013.
- Gayá P. (2021). Towards ever more extended epistemologies: pluriversity and decolonisation of knowledges in participatory inquiry. *The SAGE handbook of participatory research and inquiry*, 169–184.
- Gómez-Baggethun E, Reyes-García V. Reinterpreting change in traditional ecological knowledge. *Hum Ecol*. 2013;41(4):643–7.
- González-Tejero MR, Casares-Porcel M, Sánchez-Rojas CP, Ramiro-Gutiérrez JM, Molero-Mesa J, Pieroni A, et al. Medicinal plants in the Mediterranean area: synthesis of the results of the project Rubia. *J Ethnopharmacol*. 2008;116(2):341–57.
- ICPC-2 International Classification of Primary Care 2nd ed. Updated March 2003. Available online: <https://www.who.int/standards/classifications/other-classifications/international-classification-of-primary-care>. Accessed on 1 Feb 2024.
- ISE International Society of Ethnobiology. International Society of Ethnobiology Code of Ethics (with 2008 additions) (2016). Available online: <http://www.ethnobiology.net/what-we-do/core-programs/ise-ethics-program/code-of-ethics/code-in-english>. Accessed on 12 May 2018.
- Januszewska-Jurkiewicz J. Stosunki narodowościowe na Wileńszczyźnie w latach 1920–1939. Katowice: Wydawnictwo Uniwersytetu Śląskiego; 2010.
- Juknelienė D., Mozgeris G. (2015). The spatial pattern of forest cover changes in Lithuania during the second half of the twentieth century. *Žemės ūkio mokslai*, 22(4).
- Hammer Ø, Harper D.A.T., Ryan P.D. (2001). PAST: Paleontological Statistics Software Package for Education and Data Analysis. *Palaeontologia Electronica* 4(1): 9pp. <https://www.nhm.uio.no/english/research/resources/past/>. Accessed on 10 May 2024.
- Hanazaki N. Local and traditional knowledge systems, resistance, and socio-environmental justice. *J Ethnobiol Ethnomed*. 2024;20(1):5.
- Harrison HL, Rybråten S, Aas Ø. Hatching knowledge: a case study on the hybridization of local ecological knowledge and scientific knowledge in small-scale Atlantic salmon (*Salmo salar*) cultivation in Norway. *Hum Ecol*. 2018;46:449–59.
- Henneberg M., Stasiulewicz M. (1993). Herbal ethnopharmacology of Lithuania, Vilnius region. Medicament and food. In *Balansard G.(ed.), Cabalion Pierre (ed.), Fleurentin J.(ed.), Mazars G.(ed.) Médicaments et aliments: approche ethnopharmacologique = Medicines and foods: ethnopharmacological approach. Paris (FRA). (Colloques et Séminaires). International Conference on Ethnomedicine (Vol. 2, No. 11, pp. 243–45).*
- Kalle R, Sökand R. The name to remember: flexibility and contextuality of preliterate folk plant categorization from the 1830s, in Pernau, Livonia, historical region on the eastern coast of the Baltic Sea. *J Ethnopharmacol*. 2021;264:113254.
- Karpavičienė B. Traditional uses of medicinal plants in south-western part of Lithuania. *Plants*. 2022;11(16):2093.

33. Kartanaitė L. (2013). Etnobotaninis augalų panaudojimas Dzūkijoje. Bakalaurinis darbas. – Vilnius University.
34. Kazancı C, Oruç S, Mosulishvili M. Medicinal ethnobotany of wild plants: a cross-cultural comparison around Georgia-Turkey border, the Western Lesser Caucasus. *J Ethnobiol Ethnomed*. 2020;16:1–20.
35. Ladio A. H., Albuquerque U. P. (2014). The concept of hybridization and its contribution to urban ethnobiology. *Ethnobiology and Conservation*, 3.
36. Leonti M. The future is written: impact of scripts on the cognition, selection, knowledge and transmission of medicinal plant use and its implications for ethnobotany and ethnopharmacology. *J Ethnopharmacol*. 2011;134(3):542–55.
37. Łuczaj Ł, Jug-Dujaković M, Dolina K, Jeričević M, Vitasović-Kosić I. Insular pharmacopeias: ethnobotanical characteristics of medicinal plants used on the Adriatic islands. *Front Pharmacol*. 2021;12:623070.
38. Maclean K. Cultural hybridity and the environment: Strategies to celebrate local and indigenous knowledge. Springer; 2015.
39. Manton M, Ruffner C, Kibirkštis G, Brazaitis G, Marozas V, Pukienė R, et al. Fire occurrence in Hemi-Boreal forests: exploring natural and cultural Scots pine fire regimes using dendrochronology in Lithuania. *Land*. 2022;11(2):260.
40. Mattalia G, Stryamets N, Pieroni A, Sökand R. Knowledge transmission patterns at the border: ethnobotany of Hutsuls living in the Carpathian Mountains of Bukovina (SW Ukraine and NE Romania). *J Ethnobiol Ethnomed*. 2020;16(1):1–40.
41. Mattalia G, Belichenko O, Kalle R, Kolosova V, Kuznetsova N, Prakofjewa J, et al. Multifarious trajectories in plant-based ethnoveterinary knowledge in northern and southern eastern Europe. *Front Vet Sci*. 2021;8:710019.
42. Mattalia G, Prakofjewa J, Kalle R, Prüse B, Marozzi M, Stryamets N, et al. Centralization can jeopardize local wild plant-based food security NJAS: Impact in Agricultural and Life. Sciences. 2023;95(1):2191798.
43. Marinello V, Buckton C, Combet E. Harmless? Mixed perception and awareness of vitamin and mineral supplements. *Proc Nutr Soc*. 2016;75(OCE2):E66.
44. Mekas T, Petkeviciute Z. Parallels between ethnopharmaceutical research in Wilno in 1927 and at Wornie in 2007. *Analecta Stud Mater Dziej Nauk*. 2010;19(1–2 (36–37)):83–91.
45. Mignolo W. The darker side of western modernity: global futures, decolonial options. Duke University Press; 2011.
46. Nolan K. A., Callahan J. E. (2005). The Shannon-Weiner species diversity index. *Association for Biology Laboratory Education (ABLE) Proceedings*, 37.
47. Ososki AL, Balick MJ, Daly DC. Medicinal plants and cultural variation across Dominican rural, urban, and transnational landscapes. *Traveling cultures plants: ethnobiol ethnopharmacy hum migrations*. 2007;14:39.
48. Pardo-de-Santayana M, Pieroni A, Puri R. The ethnobotany of Europe, past and present. *Ethnobotany new Europe: people, health wild plant resour*. 2010;14:1–15.
49. Peddi B, Adams NK, Ludwig D, Desein J. The people of Techiman eat Teporo[®]: migrant farming and epistemic pluralism in Forikrom, Ghana. *Agric Hum Values*. 2025. <https://doi.org/10.1007/s10460-025-10729-w>.
50. Petkevicius R. (2011). Nervinių sutrikimų gydymas tradicinėmis priemonėmis XX–XXI a. pr. Aukštaitijoje ir Dzūkijoje (*Traditional Healing of Nervous Disorders in XX–XXI c. Aukštaitija and Dzūkija*). Diss. Vytautas Magnus University)
51. Pieroni A, Vandebroek I, Prakofjewa J, Bussmann RW, Paniagua-Zambrana NY, Maroyi A, et al. Taming the pandemic? The importance of homemade plant-based foods and beverages as community responses to COVID-19. *J Ethnobiol Ethnomed*. 2020;16:1–9.
52. Plants of the World Online (POWO). Facilitated by the Royal Botanic Gardens, Kew. 2023. <http://www.plantsoftheworldonline.org>. Accessed 12 January 2024.
53. Prakofjewa J, Kalle R, Belichenko O, Kolosova V, Sökand R. Re-written narrative: transformation of the image of Ivan-chaj in Eastern Europe. *Heliyon*. 2020. <https://doi.org/10.1016/j.heliyon.2020.e04632>.
54. Prakofjewa J, Anegg M, Kalle R, Simanova A, Prüse B, Pieroni A, et al. Diverse in local, overlapping in official medical botany: critical analysis of medicinal plant records from the historic regions of Livonia and Courland in Northeast Europe, 1829–1895. *Plants*. 2022;11(8):1065.
55. Prakofjewa J, Sartori M, Šarka P, Kalle R, Pieroni A, Sökand R. Boundaries are blurred: wild food plant knowledge circulation across the Polish-Lithuanian-Belarusian borderland. *Biology*. 2023;12(4):571.
56. Prakofjewa J, Sartori M, Šarka P, Kalle R, Pieroni A, Sökand R. Knowledge in motion: temporal dynamics of wild food plant use in the Polish-Lithuanian-Belarusian border region. *J Ethnobiol Ethnomed*. 2024;20(1):65.
57. Prakofjewa J, Sartori M, Kalle R, Łuczaj Ł, Karbarz M, Mattalia G, et al. *But how true that is, I do not know[®]*: the influence of written sources on the medicinal use of fungi across the western borderlands of the former Soviet Union. *IMA Fungus*. 2024;15(1):22.
58. Pranskuniene Z, Dauliute R, Pranskunas A, Bernatoniene J. Ethnopharmaceutical knowledge in Samogitia region of Lithuania: where old traditions overlap with modern medicine. *J Ethnobiol Ethnomed*. 2018;14:1–26.
59. R Core Team. R. A language and environment for statistical computing. Vienna: R Foundation for Statistical Computing; 2021.
60. Reyes-García V, Broesch J, Calvet-Mir L, Fuentes-Peláez N, McDade TW, Parsa S, et al. Cultural transmission of ethnobotanical knowledge and skills: an empirical analysis from an Amerindian society. *Evol Hum Behav*. 2009;30(4):274–85.
61. Sasunkevich O. Informal trade, gender and the border experience: from political borders to social boundaries. Routledge; 2016.
62. Sharifian A, Fernández-Llamazares Á, Wario HT, Molnár Z, Cabeza M. Dynamics of pastoral traditional ecological knowledge: a global state-of-the-art review. *Ecol Soc*. 2022;27(1):14.
63. Shrestha NR, Conway D. Globalization's cultural challenges: Homogenization, hybridization and heightened identity. In: *Globalization's Contradictions*. Routledge; 2006. p. 210–25.
64. Stevens P. F. (2001). Angiosperm Phylogeny Website. Version 13. Angiosperm Phylogeny Website. Version 13. St. Louis: Missouri Botanical Garden. <http://www.mobot.org/MOBOT/research/APweb/>. Accessed 30 July 2020.
65. Srithi K, Trisonthi C, Wangpakapattanawong P, Balslev H. Medicinal plants used in Hmong women's healthcare in northern Thailand. *J Ethnopharmacol*. 2012;139(1):119–35.
66. Šulskienė A. Augalai lietuvių liaudies medicinoje. Magistrinis darbas: Vilnius University; 2006.
67. Tareau MA, Bonnefond A, Palisse M, Odonne G. Phytotherapies in motion: French Guiana as a case study for cross-cultural ethnobotanical hybridization. *J Ethnobiol Ethnomed*. 2020;16(1):54.
68. Tengö M, Brondizio ES, Elmqvist T, Malmer P, Spierenburg M. Connecting diverse knowledge systems for enhanced ecosystem governance: the multiple evidence base approach. *Ambio*. 2014;43:579–91.
69. Teškevičius S. Between national and local memories: the case of Vilnius (Vilnija) region in Lithuania. *J Balt Stud*. 2023;54(3):625–40.
70. Tutin T.G., Burges N.A., Chater A.O., Edmondson J.R., Heywood V.H., Moore D.M., Valentine D.H., Walters S.M., Webb D.A. (Eds.) (1964–1980) *Flora Europaea*; University Press: Cambridge, UK; Volumes 1–5, Available online: <http://www2.bgbm.org/EuroPlusMed/query.asp>. Accessed on 2018.
71. Ubarevičienė R, Van Ham M. Population decline in Lithuania: who lives in declining regions and who leaves? *Reg Stud Reg Sci*. 2017;4(1):57–79.
72. Vandebroek I, Balick MJ. Globalization and loss of plant knowledge: challenging the paradigm. *PLoS One*. 2012;7(5):e37643.
73. Vinslauskaitė, U. (2021). *Natūraliųjų vaistinių medžiagų, vartotų širvintų apylinkėse, etnofarmacinis tyrimas* (Master's thesis, Lithuanian University of Health Sciences (Lithuania)).
74. Yanou MP, Ros-Tonen MA, Reed J, Nakwenda S, Sunderland T. The hybridisation, resilience, and loss of local knowledge and natural resource management in Zambia. *Hum Ecol*. 2024;52(5):1087–105.
75. Xaba T. Marginalized medical practice: The marginalization and transformation of indigenous medicines in South Africa. In: de Sousa Santos B, editor. *Another knowledge is possible: beyond northern epistemologies*. New York: Verso; 2007. p. 317–51.
76. Xu J, Grumbine RE. Integrating local hybrid knowledge and state support for climate change adaptation in the Asian Highlands. *Clim Change*. 2014;124:93–104.
77. Wayland C. Contextualising the politics of knowledge: physicians' attitudes toward medicinal plants. *Med Anthropol Q*. 2003;17(4):483–500.
78. Zent S, Zent E. (2013). Processual perspectives on traditional environmental knowledge. *Understanding cultural transmission in anthropology*, 213–65.
79. Ziman J, Ziman JM. *Reliable knowledge: An exploration of the grounds for belief in science*. Cambridge University Press; 1991.

Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.