

A joint limit theorems for trigonometric polynomials

Rasa ŠLEŽEVIČIENĖ (VU)
e-mail: rasa.slezeviciene@maf.vu.lt

Let $s = \sigma + it$ is a complex variable. The Riemann zeta function $\zeta(s)$, for $\sigma > 1$, is defined by

$$\zeta(s) = \sum_{m=1}^{\infty} \frac{1}{m^s},$$

and by analytic continuation elsewhere.

Denote by $\mathcal{B}(S)$ the class of Borel sets of the space S , and let, for $T > 0$,

$$\nu_T^{\dot{t}}(\dots) = \frac{1}{T} \text{meas}\{t \in [0, T], \dots\},$$

where $\text{meas}\{A\}$ stands for the Lebesgue measure of the set A , and in place of dots some condition satisfied by t is to be written.

Let N is a positive integer and

$$p_j N(t) = \sum_{m \leq N} a_{jm} \exp\{i\lambda_{jm}t\}, \quad j = 1, \dots, n,$$

be arbitrary trigonometric polynomials and $\lambda_{jm} \in \mathbb{R}$, $a_{jm} \in \mathbb{C}$.

Let \mathbb{C} is a complex plane and $\mathbb{C}^n = \underbrace{\mathbb{C} \times \mathbb{C} \times \dots \times \mathbb{C}}_n$. On the probability space

$(\mathbb{C}, \mathcal{B}(\mathbb{C}))$ define the probability measure

$$P_{T, p_N}(A) = \nu_T^{\dot{t}}\left(\left(p_{1N}(tk_1), \dots, p_{nN}(tk_n)\right) \in A\right), \quad A \in \mathcal{B}(\mathbb{C}^n).$$

Here k_1, \dots, k_n are natural numbers.

Theorem 1. *On $(\mathbb{C}^n, \mathcal{B}(\mathbb{C}^n))$ here exists a probability measure P_{p_N} such that the measure P_{T, p_N} converges weakly to P_{p_N} as $T \rightarrow \infty$.*

Proof of Theorem 1. Instead of the mesure P_{T, p_N} we can consider the measure

$$Q_{T, p_N}(A) = \nu_T^{\dot{t}}\left(\left(\text{Rep}_{1N}(tk_1), \text{Imp}_{1N}(tk_1) \dots, \text{Rep}_{nN}(tk_n), \text{Imp}_{nN}(tk_n)\right) \in A\right), \\ A \in \mathcal{B}(\mathbb{R}^{2n}).$$

Let $J_k(x)$ is the Bessel functions, and

$$\begin{aligned} \varphi_{p_N}(\tau_1, \tau'_1, \dots, \tau_n, \tau'_n) &= \sum^* \prod_{j \leq n} \prod_{m \leq N} J_{k_{jm}}(|a_{jm}| \tau_j) J_{k'_{jm}}(|a_{jm}| \tau'_j) \\ &\times \exp \left\{ i \sum_{j \leq n} \sum_{m \leq N} \left(k_{jm} \left(\varphi_{jm} + \frac{\pi}{2} \right) + k'_{jm} \varphi_{jm} \right) \right\}. \end{aligned} \quad (1)$$

Here $\varphi_{jm} = \arg a_{jm}$, $j = 1, \dots, n$, and the symbol \sum^* means the summation which runs over all integers k_{jm} and k'_{jm} , $1 \leq j \leq n$, $1 \leq m \leq N$, satisfying the condition

$$\sum_{j \leq n} \sum_{m \leq N} (k_{jm} + k'_{jm}) \lambda_{jm} k_j = 0.$$

The characteristic function of the measure Q_{T,p_N} is

$$\begin{aligned} \varphi_{T,p_N}(\tau_1, \tau'_1, \dots, \tau_n, \tau'_n) &= \int_{\mathbb{R}^{2n}} \exp \{ i(\tau_1 x_1 + \tau'_1 x'_1 + \dots + \tau_n x_n + \tau'_n x'_n) \} dQ_{T,p_N} \\ &= \frac{1}{T} \int_0^T \exp \{ i(\tau_1 \text{Rep}_{1N}(tk_1) + \tau'_1 \text{Imp}_{1N}(tk_1) + \dots + \tau_n \text{Rep}_{nN}(tk_n) + \tau'_n \text{Imp}_{nN}(tk_n)) \} dt. \end{aligned}$$

It is well known that

$$\begin{aligned} \text{Rep}_{jN}(tk_j) &= \sum_{m \leq N} |a_{jm}| \cos(\varphi_{jm} + tk_j \lambda_{jm}), \\ \text{Imp}_{jN}(tk_j) &= \sum_{m \leq N} |a_{jm}| \sin(\varphi_{jm} + tk_j \lambda_{jm}), \quad j = 1, \dots, n. \end{aligned}$$

Since [1]

$$e^{ix \sin \theta} = \sum_{r=-\infty}^{\infty} J_r(x) e^{ir\theta}$$

and

$$e^{ix \cos \theta} = \sum_{r=-\infty}^{\infty} i^r J_r(x) e^{ir\theta},$$

we have that

$$e^{i\tau_j \text{Rep}_{jN}(tk_j)} = \prod_{m \leq N} \sum_{k_{jm}=-\infty}^{\infty} J_{k_{jm}}(|a_{jm}| \tau_j) e^{ik_{jm}(\lambda_{jm} tk_j + \varphi_{jm} + \frac{\pi}{2})} \quad (2)$$

and

$$e^{i\tau'_j \text{Imp}_{jN}(tk_j)} = \prod_{m \leq N} \sum_{k'_{jm} = -\infty}^{\infty} J_{k'_{jm}}(|a_{jm}| \tau'_j) e^{ik'_{jm}(\lambda_{jm} tk_j + \varphi_{jm})}. \tag{3}$$

Substituting (2) and (3) in (1) we obtain that

$$\begin{aligned} \varphi_{T, p_N}(\tau_1, \tau'_1, \dots, \tau_n, \tau'_n) &= \varphi_{p_N}(\tau_1, \tau'_1, \dots, \tau_n, \tau'_n) \\ &+ \sum^{**} \prod_{j \leq n} \prod_{m \leq N} J_{k_{jm}}(|a_{jm}| \tau_j) J_{k'_{jm}}(|a_{jm}| \tau'_j) \\ &\times \exp \left\{ i \left(\frac{\pi}{2} \sum_{j \leq n} \sum_{m \leq N} k_{jm} + \sum_{j \leq n} \sum_{m \leq N} (k_{jm} + k'_{jm}) \varphi_{jm} \right) \right\} \\ &\times \frac{\exp \left\{ iT \sum_{j \leq n} \sum_{m \leq N} (k_{jm} + k'_{jm}) \lambda_{jm} k_j \right\} - 1}{iT \sum_{j \leq n} \sum_{m \leq N} (k_{jm} + k'_{jm}) \lambda_{jm} k_j}. \end{aligned}$$

Here the symbol \sum^{**} means the summation which runs over all integers k_{jm} and k'_{jm} , $1 \leq j \leq n$, $1 \leq m \leq N$, satisfying the condition

$$\sum_{j \leq n} \sum_{m \leq N} (k_{jm} + k'_{jm}) \lambda_{jm} k_j \neq 0.$$

Let $\varepsilon, c_1, c'_1, \dots, c_n, c'_n$ be arbitrary positive numbers, and

$$A_0 = \{(\tau_1, \tau'_1, \dots, \tau_n, \tau'_n) \in \mathbb{R}^{2n} : |\tau_1| \leq c_1, |\tau'_1| \leq c'_1, \dots, |\tau_n| \leq c_n, |\tau'_n| \leq c'_n\}.$$

Hence, using the properties of Bessel functions we find that for any $\varepsilon > 0$ and for any $(\tau_1, \tau'_1, \dots, \tau_n, \tau'_n) \in A_0$ the inequality

$$|\varphi_{T, p_N}(\tau_1, \tau'_1, \dots, \tau_n, \tau'_n) - \varphi_{p_N}(\tau_1, \tau'_1, \dots, \tau_n, \tau'_n)| < \varepsilon. \tag{4}$$

is satisfied. (4) formula shows, that φ_{T, p_N} converges weakly to φ_{p_N} as $T \rightarrow \infty$ and the convergence is uniform for $\tau_1, \tau'_1, \dots, \tau_n, \tau'_n$ in every finite interval. This and the well known continuity theorem, see, for example [2], prove the theorem.

Let D be a region on \mathbb{C} and $H(D)$ denote the space of analytic on D function equipped with the topology of uniform convergence on compacta and $H^n(D) = \underbrace{H(D) \times H(D) \times \dots \times H(D)}_n$. Let

$$p_N = \sum_{m=1}^N \frac{a(m)}{m^s}.$$

Define a probability measure

$$P_{T,p_N}(A) = \nu_T^\tau \left((p_N(s + i\tau k_1), \dots, p_N(s + i\tau k_n)) \in A \right), \quad A \in \mathcal{B}(H^n(D)).$$

Theorem 2. *There exists a probability measure P_{p_N} on $(H^n(D), \mathcal{B}(H^n(D)))$ such that the measure P_{T,p_N} converges weakly to P_{p_N} as $T \rightarrow \infty$.*

Proof of the Theorem 2. Let p_1, \dots, p_r are the distinct primes which divide the product

$$\prod_{m=1, a(m) \neq 0}^N m,$$

and let

$$\Omega_r = \prod_{j=1}^r \gamma_{p_j}, \quad \gamma_{p_j} = \gamma = \{s \in \mathbb{C} : |s| = 1\}.$$

Let us define the function $h : \Omega_r \rightarrow H^n(D)$ by the formula

$$h(x_1, \dots, x_r) = \left(\sum_{m=1}^N \frac{a(m)}{m^s} \left(\prod_{p_j^{\alpha_j} \parallel m, j \leq r} x_j^{\alpha_j} \right)^{-k_1}, \dots, \sum_{m=1}^N \frac{a(m)}{m^s} \left(\prod_{p_j^{\alpha_j} \parallel m, j \leq r} x_j^{\alpha_j} \right)^{-k_n} \right),$$

$$(x_1, \dots, x_r) \in \Omega_r.$$

The function h is continuous on Ω_r and

$$(p_N(s + ik_1\tau), \dots, p_N(s + ik_n\tau)) = h(p_1^{i\tau}, \dots, p_r^{i\tau}). \quad (5)$$

Now we define the probability measure

$$Q_T(A) = \nu_T^\tau \left((p_1^{i\tau}, \dots, p_r^{i\tau}) \in A \right).$$

on $(\Omega_r, \mathcal{B}(\Omega_r))$. The Fourier transform $g_T(l_1, \dots, l_r)$, $l_j \in \mathbb{Z}$, $j = 1, \dots, r$ of Q_T is

$$g_T(l_1, \dots, l_r) = \int_{\Omega_r} x_1^{l_1}, \dots, x_r^{l_r} dQ_T = \frac{1}{T} \int_0^T \prod_{j=1}^r p_j^{il_j\tau} d\tau$$

$$= \begin{cases} 1, & \text{if } (l_1, \dots, l_r) = (0, \dots, 0), \\ \frac{\exp \left\{ iT \sum_{j=1}^r l_j \ln p_j \right\} - 1}{iT \sum_{j=1}^r l_j \ln p_j}, & \text{if } (l_1, \dots, l_r) \neq (0, \dots, 0). \end{cases}$$

Since the logarithms of prime numbers are linearly independent over the field of rational numbers, we find that

$$g_T(l_1, \dots, l_r) = \begin{cases} 1, & \text{if } (l_1, \dots, l_r) = (0, \dots, 0) \\ 0, & \text{if } (l_1, \dots, l_r) \neq (0, \dots, 0). \end{cases}$$

as $T \rightarrow \infty$. Therefore by Theorem 1.3.19 from [1] the measure Q_T converges weakly to the Haar measure m_{rH} on $(\Omega_r, \mathcal{B}(\Omega_r))$ as $T \rightarrow \infty$. Taking into account the continuity of the function h and the formula (5) and applying Theorem 1.1.16 from [1] we obtain that the probability measure P_{T,p_N} converges weakly to the measure $m_{rH} h^{-1}$ as $T \rightarrow \infty$. The theorem is proved.

Theorems 1 and 2 will be applied to prove joint limit theorems for Dirichlet series.

References

- [1] A. Laurinćikas, *Limit Theorems for the Riemann zeta-function*, Kluwer, Dordrecht (1996).
 [2] P. Billingsley, *Convergence of Probability Measures*, John Wiley, New York (1996).

Daugiamatės ribinės teoremos trigonometriniams polinomams

R. Šleževičienė

Straipsnyje įrodomos dvi daugiamatės ribinės teoremos trigonometriniams polinomams tikimybių matų silpno konvergavimo prasme.