

## A weighted universality theorem for zeta-functions of elliptic curves

Virginija GARBALIAUSKIENĖ (ŠU)

e-mail: virginija@fm.su.lt

Let  $E: y^2 = x^3 + ax + b$ ,  $a, b \in \mathbb{Z}$ , be an elliptic curve. Suppose that the discriminant  $\Delta = -16(a^3 + 27b^2) \neq 0$ . In this case an elliptic curve is non-singular.

Let, for a prime  $p$ ,  $\nu(p)$  be the number of solutions of the congruence

$$y^2 \equiv x^3 + ax + b \pmod{p},$$

and  $\lambda(p) = p - \nu(p)$ . Then the classical result of H. Hasse asserts that

$$|\lambda(p)| \leq 2\sqrt{p}$$

for each prime  $p$ .

Let  $s = \sigma + it$  be a complex variable. To the curve  $E$  we attach the  $L$ -function  $L_E(s)$  defined, for  $\sigma > \frac{3}{2}$ , by

$$L_E(s) = \prod_{p|\Delta} \left(1 - \frac{\lambda(p)}{p^s}\right)^{-1} \prod_{p \nmid \Delta} \left(1 - \frac{\lambda(p)}{p^s} + \frac{1}{p^{2s-1}}\right)^{-1}.$$

Now it is known that  $L_E(s)$  is analytically continuable to an entire function.

The papers [2], [3], [5] are devoted to the universality of the function  $L_E(s)$ . For example, in [2], [5] the following statement is given. Let  $\nu_T(\dots) = \frac{1}{T} \text{meas}\{\tau \in [0, T]: \dots\}$ , were in place of dots a condition satisfied by  $\tau$  is to be written.

**THEOREM A.** *Let  $K$  be a compact subset of the strip  $D = \{s \in \mathbb{C}: 1 < \sigma < \frac{3}{2}\}$  with connected complement, and let  $f(s)$  be a continuous non-vanishing function on  $K$  which is analytic in the interior of  $K$ . Then, for every  $\varepsilon > 0$ ,*

$$\liminf_{T \rightarrow \infty} \nu_T \left( \sup_{s \in K} |L_E(s + i\tau) - f(s)| < \varepsilon \right) > 0.$$

The paper [3] contains a discrete version of Theorem A.

The aim of this note is to obtain a weighted universality theorem for the function  $L_E(s)$ .

Let  $T_0$  be a fixed positive number, and let  $w(\tau)$  be a positive function of bounded variation on  $[T_0, \infty)$ . Set

$$U = U(T, w) = \int_{T_0}^T w(\tau) d\tau,$$

and suppose that  $\lim_{T \rightarrow \infty} U(T, w) = +\infty$ . Moreover, we need some weighted analogue of the Birkhoff-Khinchine theorem. Denote by  $E_\xi$  the mean of the random element  $\xi$ . Let  $X(\tau, \omega)$ ,  $\tau \in \mathbb{R}$ , be an ergodic process defined on a certain probability space,  $E|X(\tau, \omega)| < \infty$ , with sample paths almost surely integrable in the Riemann sense over every finite interval. Suppose that the function  $w(\tau)$  satisfies

$$\frac{1}{U} \int_{T_0}^T w(\tau) X(t + \tau, \omega) d\tau = EX(0, \omega) + o(1 + |t|)^\delta \quad (1)$$

almost surely for all  $t \in \mathbb{R}$  with some  $\delta > 0$  as  $T \rightarrow \infty$ .

Denote by  $I_A$  the indicator function of the set  $A$ .

**THEOREM 1.** *Suppose that condition (1) is satisfied. Let  $K$  and  $f(s)$  be the same as in Theorem A. Then, for every  $\varepsilon > 0$ ,*

$$\liminf_{T \rightarrow \infty} \frac{1}{U} \int_{T_0}^T w(\tau) I_{\{\tau: \sup_{s \in K} |L_E(s+i\tau) - f(s)| < \varepsilon\}} d\tau > 0.$$

Clearly, Theorem A is a partial case of Theorem 1.

The proof of Theorem 1 is based on a limit theorem in the sense of the weak convergence of probability measures in the space of analytic functions for the function  $L_E(s)$ . Let  $G$  be a region in the complex plane. Denote by  $H(G)$  the space of functions analytic on  $G$ , equipped with the topology of uniform convergence on compacta. Let, for  $V > 0$ ,  $D_V = \{s \in \mathbb{C}: 1 < \sigma < \frac{1}{2}, |t| < V\}$ . Denote by  $\mathcal{B}(S)$  the class of Borel sets of the space  $S$ , and define the probability measure

$$P_T(A) = \frac{1}{U} \int_{T_0}^T w(\tau) I_{\{\tau: L_E(s+i\tau) \in A\}} d\tau, \quad A \in \mathcal{B}(H(D_V)).$$

Denote by  $\gamma$  the unit circle  $\gamma = \{s \in \mathbb{C}: |s| = 1\}$  on the complex plane, and let

$$\Omega = \prod_p \gamma_p,$$

where  $\gamma_p = \gamma$  for all primes  $p$ . With the product topology and pointwise multiplication the infinite-dimensional torus  $\Omega$  is a compact topological group. Therefore there exists the probability Haar measure  $m_H$  on  $(\Omega, \mathcal{B}(\Omega))$ . Thus we obtain a probability space  $(\Omega, \mathcal{B}(\Omega), m_H)$ . Let  $\omega(p)$  stand for the projection of  $\omega \in \Omega$  into the coordinate space  $\gamma_p$ , and on the probability space  $(\Omega, \mathcal{B}(\Omega), m_H)$  define the  $H(D_V)$ -valued random element  $L_E(s, \omega)$  by the formula

$$L_E(s, \omega) = \prod_{p|\Delta} \left(1 - \frac{\lambda(p)\omega(p)}{p^s}\right)^{-1} \prod_{p \nmid \Delta} \left(1 - \frac{\lambda(p)\omega(p)}{p^s} + \frac{\omega^2(p)}{p^{2s-1}}\right)^{-1}, \quad \omega \in \Omega. \quad (2)$$

**LEMMA 2.** *Under condition (1) the probability measure  $P_T$  converges weakly to the distribution of the random element  $L_E(s, \omega)$  as  $T \rightarrow \infty$ .*

*Proof.* It is not difficult to see that, for  $\sigma > \frac{3}{2}$ ,

$$L_E(s) = \prod_{p|\Delta} \left(1 - \frac{\lambda(p)}{p^s}\right)^{-1} \prod_{p \nmid \Delta} \left(1 - \frac{\alpha(p)}{p^s}\right)^{-1} \left(1 - \frac{\beta(p)}{p^s}\right)^{-1},$$

where  $\lambda(p) = \alpha(p) + \beta(p)$ , and

$$|\alpha(p)| \leq \sqrt{p}, \quad |\beta(p)| \leq \sqrt{p}.$$

Therefore,  $L_E(s)$  is the Matsumoto zeta-function [7], [4] with parameters  $\alpha = 0$  and  $\beta = \frac{1}{2}$ . Moreover, since by [1] every  $L$ -function attached to a non-singular elliptic curve over the field of rational numbers is the  $L$ -function attached to a certain newform of weight 2 of some congruence subgroup, we have that  $L_E(s)$  is an entire function, and, for  $\sigma > 1$ ,

$$L(\sigma + it) = O(|t|^{c_1}), \quad |t| \geq t_0, \quad c_1 > 0,$$

and [6]

$$\int_0^T |L(\sigma + it)|^2 dt = O(T), \quad T \rightarrow \infty.$$

Therefore, by Theorem 8 of [4] the probability measure

$$\frac{1}{U} \int_{T_0}^T w(\tau) I_{\{\tau: L_E(s+i\tau) \in A\}} d\tau, \quad A \in \mathcal{B}(H(D)),$$

where  $D_1 = \{s \in \mathbb{C}: \sigma > 1\}$ , converges weakly to the distribution of the  $H(D)$ -valued random element defined by (2) as  $T \rightarrow \infty$ . Since the function  $u: H(D_1) \rightarrow H(D)$  defined by the coordinatewise restriction is continuous, hence the lemma follows.

For the proof of Theorem 1 the support of the limit measure in Lemma 2 is needed. We recall that the support of a probability measure  $P$  defined on  $(S, \mathcal{B}(S))$  is a minimal closed set  $S_P \subset S$  such that  $P(S_P) = 1$ . The support  $S_P$  consists of all  $x \in S$  such that for every neighbourhood  $G$  of  $x$  the inequality  $P(G) > 0$  is satisfied.

Denote the limit measure in Lemma 2 by  $P_{L_E}$ , i.e.,  $P_{L_E}$  is the distribution of the random element  $L_E(s, \omega)$ .

LEMMA 3. *The support of the measure  $P_{L_E}$  is the set*

$$S_V = \{g \in H(D_V): g(s) \neq 0 \text{ or } g(s) \equiv 0\}.$$

*Proof.* The proof of the lemma coincides with that of Lemma 8 in [5]. A sketch of the proof is also given in [3], Lemma 5. A more general statement for the Matsumoto zeta-function under some additional condition is contained in [4], Lemma 6.

*Proof of Theorem 1.* Let  $K$  be a compact subset of the strip  $D$  with connected complement. Then there exists  $V > 0$  such that  $K \subset D_V$ .

First we suppose that the function  $f(s)$  has a non-vanishing continuation to the rectangle  $D_V$ . Let  $G$  be the set of functions  $g \in H(D_V)$  such that

$$\sup_{s \in K} |g(s) - f(s)| < \varepsilon.$$

Clearly, the set  $G$  is open. Moreover, by Lemma 3 we have that  $G \subset S_V$ . It is well known that the probability measure  $P_n$  converges weakly to  $P$  ( $P_n$  and  $P$  are given on  $(S, \mathcal{B}(S))$ ) if and only if

$$\liminf_{n \rightarrow \infty} P_n(G) \geq P(G) \quad (3)$$

for all open sets  $G$  of  $S$ . Therefore, in view of Lemmas 2 and 3

$$\liminf_{T \rightarrow \infty} \frac{1}{U} \int_{T_0}^T w(\tau) I_{\{\tau: \sup_{s \in K} |L_E(s+i\tau) - f(s)| < \varepsilon\}} d\tau \geq P(G) > 0,$$

and in this case the theorem is proved.

The general case is reduced to the above proved case. Let  $f(s)$  be the same as in the statement of Theorem 1. Then by the Mergelyan theorem, see, for example, [8], there exists a sequence of polynomials  $\{p_n(s)\}$  such that  $p_n(s) \rightarrow f(s)$  as  $n \rightarrow \infty$  uniformly on  $K$ . Since  $f(s) \neq 0$  on  $K$ , there exists a sufficiently large  $n_0$  such that  $p_{n_0}(s) \neq 0$  on  $K$  and

$$\sup_{s \in K} |f(s) - p_{n_0}(s)| < \frac{\varepsilon}{4}.$$

However, the polynomial  $p_{n_0}(s)$  has only finitely many zeros, and therefore we can find a region  $G$  with connected complement such that  $K \subset G$  and  $p_{n_0}(s) \neq 0$  on  $G$ . Hence, we can choose a continuous version of  $\log p_{n_0}(s)$  on  $G$  which is analytic in the interior of  $G$ . By the Mergelyan theorem again we can find a polynomial  $q_n(s)$  such that

$$\sup_{s \in K} |p_{n_0}(s) - e^{q_n(s)}| < \frac{\varepsilon}{4}.$$

The latter two inequalities show that

$$\sup_{s \in K} |f(s) - e^{q_n(s)}| < \frac{\varepsilon}{2}. \quad (4)$$

Since, clearly,  $e^{q_n(s)} \neq 0$ , the first part of the proof implies

$$\liminf_{T \rightarrow \infty} \frac{1}{U} \int_{T_0}^T w(\tau) I_{\{\tau: \sup_{s \in K} |L_E(s+i\tau) - e^{q_n(s)}| < \frac{\varepsilon}{2}\}} d\tau > 0.$$

This and inequality (3) yield the assertion of the theorem.

For example, we can take  $w(\tau) = \tau^{-1}$ .

## References

1. C. Breuil, B. Conrad, F. Diamond, R. Taylor, On the modularity of elliptic curves over  $\mathbb{Q}$ : wild 3-adic exercises, *J. Amer. Math. Soc.*, **14**, 843–939 (2001).
2. V. Garbaliuskienė, A. Laurinčikas, *Some Analytic Properties for L-functions of Elliptic Curves*, Department of Math. and Inform., Vilnius University, Preprint 2003 – 16 (2003).
3. V. Garbaliuskienė, A. Laurinčikas, *Discrete Value-Distribution of L-functions of Elliptic Curves*, Department of Mathematics and Inform., Vilnius University, Preprint 2004 – 01 (2004).
4. A. Laurinčikas, On the Matsumoto zeta-function, *Acta Arith.*, **84**(1) 1–16 (1998).
5. A. Laurinčikas, K. Matsumoto, J. Steuding, The universality of  $L$ -functions associated to newforms, *Izv. Math.*, **67**, 77–90 (2003).
6. K. Matsumoto, Value-distribution of zeta-functions, *Lecture Notes in Math.*, **1434**, 178–187 (1990).
7. K. Matsumoto, A probabilistic study on the value-distribution of Dirichlet series attached to certain cusp forms. *Nagoya Math. J.*, **116**, 123–138 (1989).
8. J.L. Walsh, Interpolation and approximation by rational functions in the complex domain, *Amer. Math. Soc. Collog. Publ.*, **20** (1960).

## REZIUMĖ

### **V. Garbaliuskienė. Ribinė teorema su svoriu elipsinių kreivių dzeta funkcijoms**

Gauta universalumo teorema su svoriu elipsinės kreivės  $L$ -funkcijai.