

## On zeros of some composite functions

Jovita Rašytė

*Department of Mathematics and Informatics, Vilnius University*  
Naugarduko 24, LT-03225 Vilnius  
E-mail: jovita.ras@gmail.com

**Abstract.** We obtain an estimate of the number of zeros for the function  $F(\zeta(s + imh))$ , where  $\zeta(s)$  is the Riemann zeta-function, and  $F : H(D) \rightarrow H(D)$  is a continuous function,  $D = \{s \in \mathbb{C} : \frac{1}{2} < \sigma < 1\}$ .

**Keywords:** Riemann zeta-function, universality.

The distribution of zeros of zeta and  $L$ -functions is the central problem of analytic number theory, and the results in the field allow to solve many other important problems. For example, the location of non-trivial zeros of the Riemann zeta-function  $\zeta(s)$ ,  $s = \sigma + it$ , has a direct relation to the distribution of prime numbers. The best result in this direction asserts that  $\zeta(s) \neq 0$  in the region

$$\sigma > 1 - \frac{c}{(\log |t|)^{\frac{2}{3}} (\log \log |t|)^{\frac{1}{3}}}, \quad |t| \geq t_0 > 0,$$

where  $c > 0$  is an absolute constant. We remind that the Riemann hypothesis says that all non-trivial zeros of  $\zeta(s)$  lie on the critical line  $\sigma = \frac{1}{2}$ , thus by this hypothesis,  $\zeta(s) \neq 0$  in the half-plane  $\sigma > \frac{1}{2}$ .

There are the zeta-functions for which the Riemann hypothesis is not true. For example, this holds for the Hurwitz-function  $\zeta(s, \alpha)$ ,  $0 < \alpha \leq 1$ , defined, for  $\sigma > 1$ , by

$$\zeta(s, \alpha) = \sum_{m=0}^{\infty} \frac{1}{(m + \alpha)^s},$$

and by analytic continuation elsewhere. If  $\alpha$  is a transcendental number, then [2]  $\zeta(s, \alpha)$  has zeros in the strip  $\frac{1}{2} < \sigma < 1$ . Also, the derivative  $\zeta'(s)$  has zeros in the strip  $0 < \sigma < 1$ .

For the investigation of zero-distribution of zeta-functions, universality theorems can be applied. The first universality theorem for the Riemann zeta-function has been proved by S.M. Voronin in [5]. The last version of this theorem is the following:

**Theorem 1.** *Suppose that  $K$  is a compact subset of the strip  $D = \{s \in \mathbb{C} : \frac{1}{2} < \sigma < 1\}$  with connected complement, and  $f(s)$  is a continuous non-vanishing function on  $K$  which is analytic in the interior of  $K$ . Then, for every  $\varepsilon > 0$ ,*

$$\liminf_{T \rightarrow \infty} \frac{1}{T} \text{meas} \left\{ \tau \in [0, T] : \sup_{s \in K} |\zeta(s + i\tau) - f(s)| < \varepsilon \right\} > 0.$$

Here  $\text{meas}\{A\}$  denotes the Lebesgue measure of a measurable set  $A \subset \mathbb{R}$ . The proof of Theorem 1 is given, for example, in [1].

Also, a discrete version of Theorem 1 is known. Let  $h > 0$  be a fixed number.

**Theorem 2.** *Suppose that  $K$  and  $f(s)$  satisfy the hypotheses of Theorem 1. Then, for every  $\varepsilon > 0$ ,*

$$\liminf_{N \rightarrow \infty} \frac{1}{N+1} \#\left\{0 \leq m \leq N: \sup_{s \in K} |\zeta(s + imh) - f(s)| < \varepsilon\right\} > 0.$$

In [3], certain discrete universality theorems were obtained for the composite function  $F(\zeta(s))$ .

We recall some of them. Denote by  $H(D)$  the space of analytic functions on  $D$  equipped with the topology of uniform convergence on compacte, and set

$$S = \{g \in H(D): g^{-1}(s) \in H(D) \text{ or } g(s) \equiv 0\}.$$

**Theorem 3.** *Suppose that the number  $\exp\{\frac{2\pi k}{h}\}$  is irrational for all  $k \in \mathbb{Z} \setminus \{0\}$ , and that  $F : H(D) \rightarrow H(D)$  is a continuous function such that, for every open set  $G \subset H(D)$ , the set  $(F^{-1}G \cap S)$  is non-empty. Let  $K \subset D$  be a compact subset with connected complement, and let  $f(s)$  be a continuous function on  $K$  which is analytic in the interior of  $K$ . Then, for every  $\varepsilon > 0$ ,*

$$\liminf_{N \rightarrow \infty} \frac{1}{N+1} \#\left\{0 \leq m \leq N: \sup_{s \in K} |F(\zeta(s + imh)) - f(s)| < \varepsilon\right\} > 0.$$

The next theorem is a simplification of Theorem 3.

**Theorem 4.** *Suppose that the number  $h$ , the set  $K$  and the function  $f(s)$  satisfy the hypotheses of Theorem 3, and that  $F : H(D) \rightarrow H(D)$  is a continuous function such that, for every polynomial  $p = p(s)$ , the set  $(F^{-1}\{p\}) \cap S$  is non-empty. Then the assertion of Theorem 3 is true.*

Now let  $V$  be an arbitrary positive number. Define

$$D_V = \left\{s \in \mathbb{C}: \frac{1}{2} < \sigma < 1, |t| < V\right\}$$

and

$$S_V = \{g \in H(D_V): g^{-1}(s) \in H(D_V) \text{ or } g(s) \equiv 0\}.$$

**Theorem 5.** *Suppose that the number  $h$ , the set  $K$  and the function  $f(s)$  satisfy the hypotheses of Theorem 3, and that  $V > 0$  is such that  $K \subset D_V$ . Let  $F : H(D_V) \rightarrow H(D_V)$  be a continuous function such that, for every polynomial  $p = p(s)$ , the set  $(F^{-1}\{p\}) \cap S_V$  is non-empty. Then the assertion of Theorem 3 is true.*

We note that, differently from Theorem 2, the approximated function in Theorems 3–5 is not necessarily non-vanishing.

The aim of his note is to prove the following statement.

**Theorem 6.** *Suppose that the number  $\exp\{\frac{2\pi k}{h}\}$  is irrational for all  $k \in \mathbb{Z} \setminus \{0\}$ , and that the function  $F$  is as in one of Theorems 3–5. Then, for arbitrary  $\sigma_1$  and  $\sigma_2$ ,  $\frac{1}{2} < \sigma_1 < \sigma_2 < 1$ , there exists a constant  $c = c(\sigma_1, \sigma_2) > 0$  such that the function  $F(\zeta(s + imh))$  has a zero in the disc*

$$|s - \hat{\sigma}| \leq \frac{\sigma_2 - \sigma_1}{2}, \quad \hat{\sigma} = \frac{\sigma_1 + \sigma_2}{2},$$

for more than  $cN$  numbers  $m$ ,  $0 \leq m \leq N$ .

First we will remind the Rouché theorem.

**Lemma 1.** *Suppose that  $G$  is a region on the complex plane bounded by a closed continuous contour  $L$ . Let  $f_1(s)$  and  $f_2(s)$  be two analytic functions on  $G$ , and  $f_1(s) \neq 0$  and  $|f_2(s)| < |f_1(s)|$  on  $L$ . Then the functions  $f_1(s)$  and  $f_1(s) + f_2(s)$  have the same number of zeros on  $G$ .*

Proof of the lemma can be found, for example, in [4].

*Proof of Theorem 6.* Let

$$\sigma_0 = \max\left(\left|\sigma_1 - \frac{3}{4}\right|, \left|\sigma_2 - \frac{3}{4}\right|\right),$$

$f(s) = s - \hat{\sigma}$  and  $0 < \varepsilon < \frac{\sigma_2 - \sigma_1}{20}$ . Then, in virtue of Theorems 3–5, there exists a constant  $c = c(\sigma_1, \sigma_2) > 0$  such that, for sufficiently large  $N$ ,

$$\frac{1}{N+1} \#\left\{0 \leq m \leq N: \sup_{|s - \frac{3}{4}| \leq \sigma_0} |F(\zeta(s + imh)) - f(s)| < \varepsilon\right\} > c. \quad (1)$$

The circle  $|s - \hat{\sigma}| = \frac{\sigma_2 - \sigma_1}{2}$  lies in the disc

$$\left|s - \frac{3}{4}\right| \leq \sigma_0.$$

Therefore, for  $m$  satisfying (1), we have that

$$\max_{|s - \hat{\sigma}| = \frac{\sigma_2 - \sigma_1}{2}} |F(\zeta(s + imh)) - (s - \hat{\sigma})| < \frac{\sigma_2 - \sigma_1}{20}.$$

This shows that the functions  $(s - \hat{\sigma})$  and

$$F(\zeta(s + imh)) - (s - \hat{\sigma})$$

satisfy the hypotheses of the Rouché theorem in the disc  $|s - \hat{\sigma}| \leq \frac{\sigma_2 - \sigma_1}{2}$ . However, the function  $s - \hat{\sigma}$  has precisely one zero  $s = \hat{\sigma}$  in that disc. Therefore, by the Rouché theorem, the function  $F(\zeta(s + imh))$  also has one zero in the disc  $|s - \hat{\sigma}| \leq \frac{\sigma_2 - \sigma_1}{2}$ . Since, in view of (1) the number of such  $m$ ,  $0 \leq m \leq N$ , is larger than  $cN$ , this proves the theorem.  $\square$

## References

- [1] A. Laurinčikas. *Limit Theorems for the Riemann Zeta-Function*. Kluwer, Dordrecht, 1996.
- [2] A. Laurinčikas and R. Garunkštis. *The Lerch Zeta-Function*. Kluwer, Dordrecht, 2002.
- [3] A. Laurinčikas and J. Rašytė. On discrete universality of a composite function. Submitted.
- [4] E.C. Titchmarsh. *The Theory of Functions*. Oxford University Press, Oxford, 1939.
- [5] S.M. Voronin. Theorem on the “universality” of the riemann zeta-function. *Math. USSR Izv.*, **9**:443–453, 1975.

## REZIUMĖ

### Apie kai kurių sudėtinių funkcijų nulius

Jovita Rašytė

Tarkime, kad  $\zeta(s)$ ,  $s = \sigma + it$ , yra Rymano dzeta funkcija,  $H(D)$ ,  $D = \{s \in \mathbb{C}: \frac{1}{2} < \sigma < 1\}$ , yra analizinių funkcijų srityje  $D$  erdvė, o  $F : H(D) \rightarrow (D)$  yra tolydi funkcija. Straipsnyje gautas funkcijos  $F(\zeta(s + imh))$  nulių skaičiaus įvertis.

*Raktiniai žodžiai:* Rymano dzeta funkcija, universalumas.