



Weighted Discrete Universality of the Riemann Zeta-Function

Antanas Laurinčikas^a, Darius Šiaučius^b and Gediminas Vadeikis^a

^a*Institute of Mathematics, Faculty of Mathematics and Informatics, Vilnius University*

Naugarduko str. 24, LT-03225 Vilnius, Lithuania

^b*Institute of Regional Development, Šiauliai University*

P. Višinskio str. 25, LT-76351 Šiauliai, Lithuania

E-mail(*corresp.*): darius.siauciunas@su.lt

E-mail: antanas.laurincikas@mif.vu.lt

E-mail: gediminas.vadeikis@mif.vu.lt

Received June 5, 2019; revised October 18, 2019; accepted October 18, 2019

Abstract. It is well known that the Riemann zeta-function is universal in the Voronin sense, i.e., its shifts $\zeta(s + i\tau)$, $\tau \in \mathbb{R}$, approximate a wide class of analytic functions. The universality of $\zeta(s)$ is called discrete if τ take values from a certain discrete set. In the paper, we obtain a weighted discrete universality theorem for $\zeta(s)$ when τ takes values from the arithmetic progression $\{kh : k \in \mathbb{N}\}$ with arbitrary fixed $h > 0$. For this, two types of h are considered.

Keywords: approximation of analytic functions, Mergelyan theorem, Riemann zeta-function, universality, weak convergence.

AMS Subject Classification: 11M06; 41A30.

1 Introduction

The Riemann zeta-function $\zeta(s)$, $s = \sigma + it$,

$$\zeta(s) = \sum_{m=1}^{\infty} \frac{1}{m^s}, \quad \sigma > 1,$$

since Riemann's and even Euler's times surprises mathematicians by the extensive field of applications and denseness of the set of its values. It is well

Copyright © 2020 The Author(s). Published by VGTU Press

This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

known the role of $\zeta(s)$ in the theory of distribution of prime numbers and in other problems of arithmetic, however, we, in this paper, prefer the denseness properties of $\zeta(s)$.

In the second decade of the last century, H. Bohr discovered [4] that the function $\zeta(s)$ takes every non-zero value infinitely many times in the strip $\{s \in \mathbb{C} : 1 < \sigma < 1 + \delta\}$ with any $\delta > 0$. H. Bohr and R. Courant proved [5] that, for fixed σ , $\frac{1}{2} < \sigma \leq 1$, the set

$$\{\zeta(\sigma + it) : t \in \mathbb{R}\} \quad (1.1)$$

is dense in \mathbb{C} . S.M. Voronin significantly generalized the above results. He obtained [25] that the set

$$\{(\zeta(s_1 + i\tau), \dots, \zeta(s_n + i\tau)) : \tau \in \mathbb{R}\}$$

with any fixed numbers s_1, \dots, s_n , $\frac{1}{2} < \text{Res } s_k < 1$, $1 \leq k \leq n$, and $s_k \neq s_m$ for $k \neq m$, and the set

$$\left\{ \left(\zeta(s + i\tau), \zeta'(s + i\tau), \dots, \zeta^{(n-1)}(s + i\tau) \right) : \tau \in \mathbb{R} \right\}$$

with every fixed s , $\frac{1}{2} < \sigma < 1$, are dense in \mathbb{C}^n . However, a much more important merit of Voronin is his so-called universality theorem for the function $\zeta(s)$ [26]. This theorem asserts that a wide class of analytic functions can be approximated by shifts $\zeta(s + i\tau)$, $\tau \in \mathbb{R}$. For a modern version of the Voronin universality theorem, it is convenient to use the following notation. Let $D = \{s \in \mathbb{C} : \frac{1}{2} < \sigma < 1\}$. Denote by \mathcal{K} the class of compact subsets of the strip D with connected complements, and by $H_0(K)$, $K \in \mathcal{K}$, the class of continuous non-vanishing functions on K that are analytic in the interior of K . Then the following theorem is true.

Theorem 1. *Let $K \in \mathcal{K}$ and $f(s) \in H_0(K)$. Then, for every $\varepsilon > 0$,*

$$\liminf_{T \rightarrow \infty} \frac{1}{T} \text{meas} \left\{ \tau \in [0, T] : \sup_{s \in K} |\zeta(s + i\tau) - f(s)| < \varepsilon \right\} > 0.$$

Here $\text{meas} A$ denotes the Lebesgue measure of a measurable set $A \subset \mathbb{R}$. By Theorem 1, the set of shifts $\zeta(s + i\tau)$ approximating a given function from $H_0(K)$ has a positive lower density, thus, it is infinite. Also, Theorem 1 can be considered as an infinite-dimensional version of the Bohr-Courant theorem on denseness of the set (1.1). The proof of Theorem 1 is given in [1] (in slightly different form), and in [9], [11], [24].

Theorem 1 is of continuous type: τ in $\zeta(s + i\tau)$ can take arbitrary real values. Also, a discrete version of Theorem 1 is known when τ takes values from a certain discrete set. Let $h > 0$ be a fixed number.

Theorem 2. *Let $K \in \mathcal{K}$ and $f(s) \in H_0(K)$. Then, for every $\varepsilon > 0$,*

$$\liminf_{N \rightarrow \infty} \frac{1}{N} \# \left\{ 1 \leq k \leq N : \sup_{s \in K} |\zeta(s + ikh) - f(s)| < \varepsilon \right\} > 0.$$

Here $\#A$ denotes the cardinality of the set A . The proof of Theorem 2 can be found in [22] and [1]. For shifts $\zeta(s + ik^\alpha h)$ with fixed α , $0 < \alpha < 1$, Theorem 2 is given in [6]. In [15, 21] and [7, 8, 13, 16], more general shifts of Dirichlet L -functions and Riemann zeta-function, respectively, were considered. We note that discrete universality theorems for zeta-functions sometimes are more convenient for practical applications, an example of this is the paper [3].

In [10], a weighted version of Theorem 1 was proposed. Let $w(t)$ be a function of bounded variation on $[T_0, \infty)$ with some $T_0 > 0$ such that the variation $V_a^b w$ on $[a, b]$ satisfies the inequality $V_a^b w \leq cw(a)$ with a certain constant $c > 0$ for any subinterval $[a, b] \subset [T_0, \infty)$. Let

$$U_T = U(T, w) = \int_{T_0}^T w(t)dt,$$

and let $\lim_{T \rightarrow \infty} U(T, w) = +\infty$. Moreover, let I_A denote the indicator function of the set A . Then we have the following generalization of Theorem 1.

Theorem 3. *Suppose that the function $w(t)$ satisfies the above hypotheses. Let $K \in \mathcal{K}$ and $f(s) \in H_0(K)$. Then, for every $\varepsilon > 0$,*

$$\liminf_{T \rightarrow \infty} \frac{1}{U_T} \int_{T_0}^T w(\tau) I_{\left\{ \tau: \sup_{s \in K} |\zeta(s+i\tau) - f(s)| < \varepsilon \right\}}(\tau) d\tau > 0.$$

To be precise, in [10], Theorem 3 was proved under a certain additional hypothesis on the function $w(t)$ which is a weighted version of the classical Birkhoff-Khinchine ergodic theorem. In [18], this technical hypothesis was removed. A generalization of Theorem 3 for Matsumoto zeta-functions was given in [12]. In [17], a weighted discrete universality theorem with the sequence $\{k^\alpha h\}$, $0 < \alpha < 1$, for the periodic zeta-function was obtained.

The aim of this paper is a weighted discrete universality theorem for the Riemann zeta-function. Let $w(t)$ be a real non-negative function having a continuous derivative on $[\frac{1}{2}, \infty)$ such that

$$\lim_{N \rightarrow \infty} V_N = +\infty, \quad V_N = \sum_{k=1}^N w(k), \quad \int_1^N u|w'(u)|du \ll V_N, \quad N \rightarrow \infty.$$

Denote by W the class of functions $w(t)$ satisfying the above hypotheses. Suppose that h is a fixed positive number.

Theorem 4. *Suppose that $w(t) \in W$. Let $K \in \mathcal{K}$ and $f(s) \in H_0(K)$. Then, for every $\varepsilon > 0$,*

$$\liminf_{N \rightarrow \infty} \frac{1}{V_N} \sum_{k=1}^N w(k) I_{\left\{ k: \sup_{s \in K} |\zeta(s+ikh) - f(s)| < \varepsilon \right\}}(k) > 0.$$

For example, the function $w(t) = \frac{\sin(\log t) + 1}{t}$ is not monotonically decreasing and $w(t) \in W$.

Theorem 4 has the following modification.

Theorem 5. *Suppose that $w(t) \in W$. Let $K \in \mathcal{K}$ and $f(s) \in H_0(K)$. Then the limit*

$$\lim_{N \rightarrow \infty} \frac{1}{V_N} \sum_{k=1}^N w(k) I_{\left\{k: \sup_{s \in K} |\zeta(s+ikh) - f(s)| < \varepsilon\right\}}(k) > 0$$

exists for all but at most countably many $\varepsilon > 0$.

For proving of the above universality theorems, we will apply the probabilistic approach.

2 Limit theorems

We remind that $D = \{s \in \mathbb{C} : \frac{1}{2} < \sigma < 1\}$, and by $H(D)$ denote the space of analytic functions on D endowed with the topology of uniform convergence on compacta. The space $H(D)$ is metrisable. There exists a sequence of compact subsets $\{K_l : l \in \mathbb{N}\} \subset D$ such that $D = \bigcup_{l=1}^{\infty} K_l$, $K_l \subset K_{l+1}$ for all $l \in \mathbb{N}$, and if $K \subset D$ is a compact set, then $K \subset K_l$ for some $l \in \mathbb{N}$. For $g_1, g_2 \in H(D)$, define

$$\rho(g_1, g_2) = \sum_{l=1}^{\infty} 2^{-l} \frac{\sup_{s \in K} |g_1(s) - g_2(s)|}{1 + \sup_{s \in K} |g_1(s) - g_2(s)|}.$$

Then ρ is a metric on $H(D)$ which induces its topology of uniform convergence on compacta.

Denote by $\mathcal{B}(X)$ the Borel σ -field of the space X , and, for $A \in \mathcal{B}(H(D))$, define

$$P_N(A) = P_{N,w,h}(A) = \frac{1}{V_N} \sum_{k=1}^N w(k) I_{\{k: \zeta(s+ikh) \in A\}}(k).$$

In this section, we will consider the weak convergence of $P_{N,w,h}$ as $N \rightarrow \infty$. We say that $h > 0$ is of type 1 if $\exp\left\{\frac{2\pi m}{h}\right\}$ is an irrational number for all $m \in \mathbb{Z} \setminus \{0\}$, and $h > 0$ is of type 2 if h is not of type 1. We will examine separately the cases of types 1 and 2.

As usual, we start with one topological structure. Let $\gamma = \{s \in \mathbb{C} : |s| = 1\}$ and $\Omega = \prod_p \gamma_p$, where $\gamma_p = \gamma$ for all primes p . By the Tikhonov theorem, the torus Ω with the product topology and pointwise multiplication is a compact topological Abelian group. Therefore, on $(\Omega, \mathcal{B}(\Omega))$, the probability Haar measure m_H can be defined, and this gives the probability space $(\Omega, \mathcal{B}(\Omega), m_H)$. Let \mathbb{P} be the set of all prime numbers, and let $\omega(p)$ denote the projection of $\omega \in \Omega$ to the circle γ_p , $p \in \mathbb{P}$. For $A \in \mathcal{B}(\Omega)$, define

$$Q_N(A) = \frac{1}{V_N} \sum_{k=1}^N w(k) I_{\{k: (p^{-ikh}, p \in \mathbb{P}) \in A\}}(k).$$

Lemma 1. *Suppose that $w(t) \in W$ and h is of type 1. Then Q_N converges weakly to the Haar measure m_H as $N \rightarrow \infty$.*

Proof. We apply the Fourier transform method. Let $g_N(\underline{k})$, $\underline{k} = (k_p : k_p \in \mathbb{Z}, p \in \mathbb{P})$, be the Fourier transform of Q_N . Then we have that

$$g_N(\underline{k}) = \int_{\Omega} \prod_p^* \omega^{k_p}(p) dQ_N,$$

where the sign “*” means that only a finite number of integers k_p are distinct from zero. Thus, by the definition of Q_N ,

$$g_N(\underline{k}) = \frac{1}{V_N} \sum_{k=1}^N w(k) \prod_p^* p^{-ik_k h} = \frac{1}{V_N} \sum_{k=1}^N w(k) \exp \left\{ -ikh \sum_p^* k_p \log p \right\}. \tag{2.1}$$

Obviously,

$$g_N(\underline{0}) = 1. \tag{2.2}$$

If $\underline{k} \neq \underline{0}$, then

$$\sum_p^* k_p \log p \neq 0,$$

since the logarithms of prime numbers are linearly independent over the field of rational numbers. Thus,

$$\exp \left\{ -ih \sum_p^* k_p \log p \right\} \neq 1. \tag{2.3}$$

Indeed, if inequality (2.3) is not true, then

$$\sum_p^* k_p \log p = \frac{2\pi r}{h}, \quad \prod_p^* p^{-k_p} = \exp \left\{ \frac{2\pi r}{h} \right\}$$

with some $r \in \mathbb{Z} \setminus \{0\}$. However, the left-hand side of this equality is a rational number, and we arrive to the contradiction that h is of type 1. Thus, (2.3) is true, and we find that, for $u \geq 1$,

$$\begin{aligned} & \sum_{k \leq u} \exp \left\{ -ikh \sum_p^* k_p \log p \right\} \\ &= \frac{\exp \left\{ -ih \sum_p^* k_p \log p \right\} - \exp \left\{ i([u] + 1)h \sum_p^* k_p \log p \right\}}{1 - \exp \left\{ -ih \sum_p^* k_p \log p \right\}} \stackrel{def}{=} \Sigma(u). \end{aligned}$$

Hence, in view of (2.1), for $\underline{k} \neq \underline{0}$,

$$g_N(\underline{k}) = \frac{w(N)\Sigma(N)}{V_N} - \frac{1}{V_N} \int_1^N \Sigma(u)w'(u)du.$$

Since the function $\Sigma(u)$ is bounded by a constant not depending of u , we find that, for $\underline{k} \neq \underline{0}$,

$$\lim_{N \rightarrow \infty} g_N(\underline{k}) = 0.$$

This together with (2.2) proves the lemma. \square

Lemma 1 implies a weighted discrete universality theorem for absolutely convergent Dirichlet series. Let $\theta > \frac{1}{2}$ be a fixed number, and

$$v_n(m) = \exp \left\{ - \left(\frac{m}{n} \right)^\theta \right\}, \quad m, n \in \mathbb{N},$$

$$\zeta_n(s) = \sum_{m=1}^{\infty} \frac{v_n(m)}{m^s}, \quad \zeta_n(s, \omega) = \sum_{m=1}^{\infty} \frac{\omega(m)v_n(m)}{m^s},$$

where

$$\omega(m) = \prod_{\substack{p^\alpha | m \\ p^{\alpha+1} \nmid m}} \omega^\alpha(p), \quad m \in \mathbb{N}.$$

Then the series for $\zeta_n(s)$ and $\zeta_n(s, \omega)$ are absolutely convergent for $\sigma > \frac{1}{2}$ [11]. From this, it follows that the function $u_n : \Omega \rightarrow H(D)$, $u_n(\omega) = \zeta_n(s, \omega)$, is continuous. Let $R_n = m_H u_n^{-1}$, where

$$R_n(A) = m_H u_n^{-1}(A) = m_H(u_n^{-1}A), \quad A \in \mathcal{B}(H(D)).$$

Moreover, let

$$P_{N,n}(A) = \frac{1}{V_N} \sum_{k=1}^N w(k) I_{\{k: \zeta_n(s+ikh) \in A\}}(k), \quad A \in \mathcal{B}(H(D)).$$

It is not difficult to see that $P_{N,n} = Q_N u_n^{-1}$. This, the continuity of u_n and Lemma 1 lead to

Lemma 2. *Suppose that $w(t) \in W$ and h is of type 1. Then $P_{N,n}$ converges weakly to R_n as $N \rightarrow \infty$.*

The weak convergence of $P_{N,n}$ is a starting point for proving the weak convergence for P_N as $N \rightarrow \infty$. The investigation of P_N also requires an approximation of $\zeta(s)$ by $\zeta_n(s)$. Let $l_n(s) = \frac{s}{\theta} \Gamma\left(\frac{s}{\theta}\right) n^s$, where $\Gamma(s)$ is the Euler gamma-function. Then [11], for $\sigma > \frac{1}{2}$, the integral representation

$$\zeta_n(s) = \frac{1}{2\pi i} \int_{\theta-i\infty}^{\theta+i\infty} \zeta(s+z) l_n(z) \frac{dz}{z} \quad (2.4)$$

is true. Using the well-known estimates

$$\int_{1/2}^T |\zeta(\sigma+it)|^2 dt \ll T, \quad \int_{1/2}^T |\zeta'(\sigma+it)|^2 dt \ll T,$$

we find that, for $\frac{1}{2} < \sigma < 1$ and $\tau \in \mathbb{R}$,

$$\int_{1/2}^T |\zeta(\sigma+it+i\tau)|^2 dt \ll T(1+|\tau|)$$

and

$$\int_{1/2}^T |\zeta'(\sigma + it + i\tau)|^2 dt \ll T(1 + |\tau|).$$

These estimates together with Gallagher lemma, see, for example, [20, Lemma 1.4], give, for $\frac{1}{2} < \sigma < 1$ and $\tau \in \mathbb{R}$, the bound

$$\begin{aligned} \sum_{k=1}^N |\zeta(\sigma + ikh + i\tau)|^2 &\ll \int_{1/2}^{(N+1/2)h} |\zeta(\sigma + it + i\tau)|^2 dt \\ &+ \left(\int_{1/2}^{(N+1/2)h} |\zeta(\sigma + it + i\tau)|^2 dt \int_{1/2}^{(N+1/2)h} |\zeta'(\sigma + it + i\tau)|^2 dt \right)^{1/2} \ll N(1 + |\tau|). \end{aligned}$$

Hence, for the same σ and τ ,

$$\begin{aligned} \sum_{k=1}^N w(k) |\zeta(\sigma + ikh + i\tau)|^2 &\ll w(N) \sum_{k=1}^N |\zeta(\sigma + ikh + i\tau)|^2 + (1 + |\tau|) \\ &\times \int_1^N u |w'(u)| du \ll Nw(N)(1 + |\tau|) + V_N(1 + |\tau|) \ll V_N(1 + |\tau|), \end{aligned} \tag{2.5}$$

because

$$Nw(N) = \sum_{k=1}^N w(k) + \int_1^N \left(\sum_{k \leq u} 1 \right) w'(u) du \ll V_N.$$

Let $K \subset D$ be a compact set. Then (2.4), (2.5), the residue theorem and Cauchy integral formula imply the equality

$$\lim_{n \rightarrow \infty} \limsup_{N \rightarrow \infty} \frac{1}{V_N} \sum_{k=1}^N w(k) \sup_{s \in K} |\zeta(s + ikh) - \zeta_n(s + ikh)| = 0. \tag{2.6}$$

Now, (2.6) together with the definition of the metric ρ yields the following lemma.

Lemma 3. *Suppose that $w(t) \in W$. Then the equality*

$$\lim_{n \rightarrow \infty} \limsup_{N \rightarrow \infty} \frac{1}{V_N} \sum_{k=1}^N w(k) \rho(\zeta(s + ikh), \zeta_n(s + ikh)) = 0$$

is true for every fixed $h > 0$.

Now, we are in position to prove a weighted discrete limit theorem for the function $\zeta(s)$. On the probability space $(\Omega, \mathcal{B}(\Omega), m_H)$, define the $H(D)$ -valued random element $\zeta(s, \omega)$ by the Euler product

$$\zeta(s, \omega) = \prod_p \left(1 - \frac{\omega(p)}{p^s} \right)^{-1}.$$

The latter product, for almost all $\omega \in \Omega$, is uniformly convergent on compact subsets of the strip D [11]. Denote by P_ζ the distribution of the random element $\zeta(s, \omega)$, i.e.,

$$P_\zeta(A) = m_H(\omega \in \Omega : \zeta(s, \omega) \in A), \quad A \in \mathcal{B}(H(D)).$$

Theorem 6. *Suppose that $w(t) \in W$ and $h > 0$ is of the type 1. Then P_N converges weakly to P_ζ as $N \rightarrow \infty$. Moreover, the support of P_ζ is the set $S = \{g \in H(D) : g(s) \neq 0 \text{ or } g(s) \equiv 0\}$.*

Proof. We will prove that R_n , as $n \rightarrow \infty$, converges weakly to a certain probability measure P , and that P_N , as $N \rightarrow \infty$, also converges weakly to P .

Let θ_N be a random variable defined on a certain probability space with probability measure μ and having the distribution

$$\mu(\theta_N = kh) = \frac{w(k)}{V_N}, \quad k = 1, \dots, N.$$

Moreover, let $Y_{N,n} = Y_{N,n}(s)$ be an $H(D)$ -valued random element defined by

$$Y_{N,n}(s) = \zeta_n(s + i\theta_N),$$

and let $Y_n = Y_n(s)$ be an $H(D)$ -valued random element with the distribution R_n . Then, by Lemma 2,

$$Y_{N,n} \xrightarrow[N \rightarrow \infty]{\mathcal{D}} Y_n. \quad (2.7)$$

Using the absolute convergence of the series for $\zeta_n(s)$, it can be proved by a method of [11] that the family of probability measures $\{R_n : n \in \mathbb{N}\}$ is tight, i.e., for every $\varepsilon > 0$, there exists a compact set $K = K(\varepsilon) \subset H(D)$ such that

$$R_n(K) > 1 - \varepsilon$$

for all $n \in \mathbb{N}$. Hence, by the Prokhorov theorem [2], this family is relatively compact. Therefore, each sequence of $\{R_n\}$ contains a subsequence $\{R_{n_r}\}$ weakly convergent, as $r \rightarrow \infty$, to a certain probability measure P on $(H(D), \mathcal{B}(H(D)))$. In other words,

$$Y_{n_r} \xrightarrow[r \rightarrow \infty]{\mathcal{D}} P. \quad (2.8)$$

Define one more $H(D)$ -valued random element

$$X_N = X_N(s) = \zeta(s + i\theta_N).$$

Then the application of Lemma 3 gives, for $\varepsilon > 0$,

$$\begin{aligned} & \lim_{n \rightarrow \infty} \limsup_{N \rightarrow \infty} \mu(\rho(X_N(s), Y_{N,n}(s)) \geq \varepsilon) \\ &= \lim_{n \rightarrow \infty} \limsup_{N \rightarrow \infty} \frac{1}{V_N} \sum_{k=1}^N w(k) I_{\{k: \rho(\zeta(s+ikh), \zeta_n(s+ikh)) \geq \varepsilon\}}(k) \\ &\leq \lim_{n \rightarrow \infty} \limsup_{N \rightarrow \infty} \frac{1}{\varepsilon V_N} \sum_{k=1}^N w(k) \rho(\zeta(s+ikh), \zeta_n(s+ikh)) = 0. \end{aligned}$$

This equality, (2.7) and (2.8) show that all hypotheses of Theorem 4.2 of [2] are satisfied, therefore,

$$X_N \xrightarrow[N \rightarrow \infty]{\mathcal{D}} P, \tag{2.9}$$

or P_N converges weakly to P as $N \rightarrow \infty$. Moreover, in virtue of (2.9), the measure P is independent of the sequence Y_{n_r} . Since the family $\{R_n\}$ is relatively compact, from this, we obtain that R_n converges weakly to P as $n \rightarrow \infty$. Thus, P_N , as $N \rightarrow \infty$, converges weakly to the limit measure P of R_n as $n \rightarrow \infty$. However, by the proof of a limit theorem for

$$\frac{1}{T} \text{meas} \{ \tau \in [0, T] : \zeta(s + i\tau) \in A \}, \quad A \in \mathcal{B}(H(D)),$$

it is known [11] that R_n , as $n \rightarrow \infty$, converges weakly to P_ζ , and the support of P_ζ is the set S . Therefore, the same statement is also true for P_N , and the theorem is proved. \square

The case of h of type 2 is a more complicated. We must construct a new probability space different from $(\Omega, \mathcal{B}(\Omega), m_H)$. We will index by h the notation related to h of type 2.

Now suppose that $h > 0$ is of type 2. Then there exists the smallest $m_0 \in \mathbb{N}$ such the number $\exp \left\{ \frac{2\pi m_0}{h} \right\}$ is rational. We put

$$\exp \left\{ \frac{2\pi m_0}{h} \right\} = \frac{a}{b}, \quad a, b \in \mathbb{N}, (a, b) = 1.$$

Define the set

$$\mathbb{P}_0 = \left\{ p \in \mathbb{P} : \frac{a}{b} = \prod_{p \in \mathbb{P}} p^{\alpha_p} \text{ with } \alpha_p \neq 0 \right\}.$$

Denote by Ω_h the closed subgroup of Ω generated by the element $\{p^{-ih} : p \in \mathbb{P}\}$. By Lemma 1 of [14], if h is of type 2, then

$$\Omega_h = \{ \omega \in \Omega : \omega(a) = \omega(b) \}.$$

On $(\Omega_h, \mathcal{B}(\Omega_h))$, the probability Haar measure m_H^h exists, and we obtain the probability space $(\Omega_h, \mathcal{B}(\Omega_h), m_H^h)$. By (3.1) of [14], we have that the characters χ of the group Ω_h are of the form

$$\chi(\omega) = \prod_{p \in \mathbb{P} \setminus \mathbb{P}_0}^* \omega^{k_p}(p) \prod_{p \in \mathbb{P}_0} \omega^{k_p + l\alpha_p}(p), \quad l \in \mathbb{Z}. \tag{2.10}$$

Now, we are ready to prove an analogue of Lemma 1 for h of type 2. For $A \in \mathcal{B}(\Omega_h)$, define

$$Q_{N,h}(A) = \frac{1}{V_N} \sum_{k=1}^N w(k) I_{\{k: (p^{-ikh}, p \in \mathbb{P}) \in A\}}(k).$$

Lemma 4. *Suppose that h is of type 2. Then $Q_{N,h}$ converges weakly the Haar measure m_H^h as $N \rightarrow \infty$.*

Proof. In view of (2.10), we have that the Fourier transform $g_{N,h}(\underline{k})$, $\underline{k} = (k_p : k_p \in \mathbb{Z}, p \in \mathbb{P})$, of $Q_{N,h}$ is of the form

$$\begin{aligned} g_{N,h}(\underline{k}) &= \int_{\Omega_h} \chi(\omega) dQ_{N,h} \\ &= \frac{1}{V_N} \sum_{k=1}^N w(k) \prod_{p \in \mathbb{P} \setminus \mathbb{P}_0}^* p^{-ik_k h} \prod_{p \in \mathbb{P}_0} p^{-ikh(k_p + l\alpha_p)}, \quad l \in \mathbb{Z}. \end{aligned} \quad (2.11)$$

If $k_p = 0$ for all $p \in \mathbb{P} \setminus \mathbb{P}_0$ and $k_p = r\alpha_p$ for all $p \in \mathbb{P}_0$ with some $r \in \mathbb{Z}$ (case 1), then

$$g_{N,h}(\underline{k}) = 1, \quad (2.12)$$

because $\prod_{p \in \mathbb{P}_0} \omega^{d\alpha_p}(p) = 1$ with $d \in \mathbb{Z}$.

Now, suppose that $k_p \neq 0$ for some $p \in \mathbb{P} \setminus \mathbb{P}_0$, or there does not exist $r \in \mathbb{Z}$ such that $k_p = r\alpha_p$ for all $p \in \mathbb{P}_0$ (case 2). In [14], it was obtained that

$$\exp\{-ihA_p(k_p, l\alpha_p)\} \neq 1,$$

where

$$A_p(k_p, l\alpha_p) = \sum_{p \in \mathbb{P} \setminus \mathbb{P}_0}^* k_p \log p + \sum_{p \in \mathbb{P}_0} (k_p + l\alpha_p) \log p, \quad l \in \mathbb{Z}.$$

Hence, we find that, for $u \geq 1$,

$$\begin{aligned} &\sum_{k \leq u} \exp\{-ikhA_p(k_p, l\alpha_p)\} \\ &= \frac{\exp\{-ihA_p(k_p, l\alpha_p)\} - \exp\{-ih([u] + 1)A_p(k_p, l\alpha_p)\}}{1 - \exp\{-ihA_p(k_p, l\alpha_p)\}} \stackrel{\text{def}}{=} \Sigma_h(u). \end{aligned}$$

Therefore, in view of (2.11),

$$g_{N,h}(\underline{k}) = \frac{w(N)\Sigma_h(N)}{V_N} - \frac{1}{V_N} \int_1^N \Sigma_h(u)w'(u)du.$$

Using the properties of the function w , hence we find that

$$g_{N,h}(\underline{k}) = 0.$$

This together with (2.12) shows that

$$\lim_{N \rightarrow \infty} g_{N,h}(\underline{k}) = \begin{cases} 1, & \text{in the case 1,} \\ 0, & \text{in the case 2.} \end{cases}$$

Since the right-hand side of the equality is the Fourier transform of the Haar measure m_H^h , the lemma follows by a continuity theorem for probability measures on compact groups. \square

Now, together with $P_{N,n,h}$, consider

$$\hat{P}_{N,n,h}(A) = \frac{1}{V_N} \sum_{k=1}^N w(k) I_{\{k: \zeta_{n,h}(s+ikh, \omega) \in A\}}(k), \quad A \in \mathcal{B}(H(D)),$$

with $\omega \in \Omega_h$.

Lemma 5. *Suppose that $w(t) \in W$ and h is of type 2. Then $P_{N,n,h}$ and $\hat{P}_{N,n,h}$ both converge weakly to the measure $m_H^h u_{n,h}^{-1}$ as $N \rightarrow \infty$, where $u_{n,h} : \Omega_h \rightarrow H(D)$ is given by $u_{n,h}(\omega) = \zeta_{n,h}(s, \omega)$, $\omega \in \Omega_h$.*

Proof. By proving Lemma 2, in view of Lemma 4, we have that $P_{N,n,h}$ converges weakly to $m_H^h u_{n,h}^{-1}$ as $N \rightarrow \infty$. Similarly, we obtain that if $\hat{u}_{n,h}(\hat{\omega}) : \Omega_h \rightarrow H(D)$ is given by

$$\hat{u}_{n,h}(\hat{\omega}) = \zeta_n(s, \omega\hat{\omega}), \quad \hat{\omega} \in \Omega_h,$$

then $\hat{P}_{N,n,h}$ converges weakly to $m_H^h \hat{u}_{n,h}^{-1}$. However, $\hat{u}_{n,h} = u_{n,h}(u)$, where $u : \Omega_h \rightarrow \Omega_h$ is given by $u(\hat{\omega}) = \omega\hat{\omega}$. This and the invariance of the Haar measure m_H^h show that $m_H^h \hat{u}_{n,h}^{-1} = m_H^h u_{n,h}^{-1}$. \square

For further considerations, we need some elements of the ergodic theory. Let $a_h = (p^{-ih} : p \in \mathbb{P})$. Then a_h is an element of Ω_h . Define the transformation $\varphi_h(\omega)$ of Ω_h by

$$\varphi_h(\omega) = a_h \omega, \quad \omega \in \Omega_h.$$

Then we have that φ_h is a measurable measure preserving transformation on the probability space $(\Omega_h, \mathcal{B}(\Omega_h), m_H^h)$. We recall that a set $A \in \mathcal{B}(\Omega_h)$ is called invariant with respect to φ_h if the sets A and $\varphi_h(A)$ can differ from each other at most by a set of m_H^h -measure zero. The transformation φ_h is called ergodic if the σ -field of invariant sets of Ω_h consists only of the sets having m_H^h -measure 1 or 0.

Lemma 6. *Suppose that h is of type 2. Then the transformation φ_h is ergodic.*

Proof of the lemma is given in [14, Lemma 3].

Let, for $\omega \in \Omega_h$,

$$\zeta_h(s, \omega) = \prod_p \left(1 - \frac{\omega(p)}{p^s} \right)^{-1}.$$

The first application of Lemma 6 is devoted to the discrete mean square of $\zeta_h(s, \omega)$.

Lemma 7. *Suppose that $w(t) \in W$, $h > 0$ is of type 2, $\sigma, \frac{1}{2} < \sigma < 1$, is fixed and $t \in \mathbb{R}$. Then, for almost all $\omega \in \Omega_h$,*

$$\sum_{k=1}^N w(k) |\zeta_h(\sigma + it + ikh, \omega)|^2 \ll V_N(1 + |t|).$$

Proof. We have that $\zeta_h(s, \omega)$ coincides with the restriction of the random element $\zeta(s, \omega)$ to the space $(\Omega_h, \mathcal{B}(\Omega_h), m_H^h)$. First we consider the expectation $\mathbb{E}|\zeta_h(\sigma + it, \omega)|^2$. We write $\zeta_h(s, \omega)$ in the form

$$\zeta_h(\sigma + it, \omega) = \prod_{p \in \mathbb{P}_0} \left(1 - \frac{\omega(p)}{p^{\sigma+it}} \right)^{-1} \prod_{p \in \mathbb{P} \setminus \mathbb{P}_0} \left(1 - \frac{\omega(p)}{p^{\sigma+it}} \right)^{-1} \stackrel{def}{=} X_1 X_2. \quad (2.13)$$

The random elements X_1 and X_2 are independent, moreover, for almost all $\omega \in \Omega_h$,

$$X_2 = \sum'_m \frac{\omega(m)}{m^{\sigma+it}},$$

where the sign $'$ means that the summing runs over $m = 1$ and $m \in \mathbb{N}$ with the canonical representation consisting only of primes $p \in \mathbb{P} \setminus \mathbb{P}_0$. In the series for X_2 , the random variables are orthogonal, therefore,

$$\mathbb{E}|X_2|^2 = \sum'_m \frac{1}{m^{2\sigma}} < \infty.$$

Clearly, $\mathbb{E}|X_1|^2$ is bounded by a constant. Therefore, there exists a finite constant $c > 0$ such that, for $\frac{1}{2} < \sigma < 1$ and $t \in \mathbb{R}$,

$$\mathbb{E}|\zeta_h(\sigma + it, \omega)|^2 = \mathbb{E}|X_1|^2 \mathbb{E}|X_2|^2 \leq c.$$

Then (2.13), Lemma 6, the Birkhoff-Khinchine ergodic theorem, see, for example, [23], and the definition of the transformation φ_h show that, for $\frac{1}{2} < \sigma < 1$ and $|t_0| \leq h$,

$$\begin{aligned} \sum_{k=1}^N |\zeta_h(\sigma + it_0 + ikh, \omega)|^2 &= \sum_{k=1}^N |\zeta_h(\sigma + it_0, \varphi_h^k(\omega))|^2 \\ &= N \mathbb{E}|\zeta_h(\sigma + it_0, \omega)|^2 (1 + o(1)) \ll N \end{aligned}$$

for almost all $\omega \in \Omega_h$ as $N \rightarrow \infty$. Hence, denoting by $[u]$ the integer part of $u \in \mathbb{R}$, for $\frac{1}{2} < \sigma < 1$ and $t \in \mathbb{R}$, we find that

$$\sum_{k=1}^N |\zeta_h(\sigma + it + ikh, \omega)|^2 = \sum_{k=1+[t/h]}^{N+[t/h]} |\zeta_h(\sigma + it_0 + ikh, \omega)|^2 \ll N(1 + |t|)$$

for almost all $\omega \in \Omega_h$. From this, summing by parts, we obtain the estimate of the lemma. \square

Similarly to the proof of Lemma 3, we arrive, by using Lemma 7, to

Lemma 8. *Suppose that $w(t) \in W$ and $h > 0$ is of type 2. Then, for almost all $\omega \in \Omega_h$,*

$$\lim_{N \rightarrow \infty} \limsup_{N \rightarrow \infty} \frac{1}{V_N} \sum_{k=1}^N w(k) \rho(\zeta_h(s + ikh, \omega), \zeta_{n,h}(s + ikh, \omega)) = 0.$$

For $\omega \in \Omega_h$, additionally to the measure $P_{N,h}$, define

$$\hat{P}_{N,h}(A) = \frac{1}{V_N} \sum_{k=1}^N w(k) I_{\{k: \zeta_h(s+ikh, \omega) \in A\}}(k), \quad A \in \mathcal{B}(H(D)).$$

Then, using Lemmas 3, 5 and 8, and repeating the first part of the proof of Theorem 6, we obtain

Lemma 9. *Suppose that $w(t) \in W$ and $h > 0$ is of type 2. Then, on $(H(D), \mathcal{B}(H(D)))$, there exists a probability measure P_h such that $P_{N,h}$ and $\hat{P}_{N,h}$ both converges weakly to P_h as $N \rightarrow \infty$.*

Denote by $P_{\zeta,h}$ the distribution of the random element $\zeta_h(s, \omega)$, $\omega \in \Omega_h$. Then we have the following analogue of Theorem 6.

Theorem 7. *Suppose that $w(t) \in W$ and $h > 0$ is of type 2. Then $P_{N,h}$ converges weakly $P_{\zeta,h}$ as $N \rightarrow \infty$. Moreover, the support of the measure $P_{\zeta,h}$ is the set S .*

Proof. In virtue of Lemma 9, it suffices to identify the measure P in that lemma, and to find the support of the limit measure. For the first problem, we will apply Lemma 6, and the Birkhoff-Khintchine theorem. Let A be a continuity set of P . On the probability space $(\Omega_h, \mathcal{B}(\Omega_h), m_H^h)$, define the random variable ξ by the formula

$$\xi(\omega) = \begin{cases} 1, & \text{if } \zeta_h(s, \omega) \in A, \\ 0, & \text{otherwise.} \end{cases}$$

Then we have that

$$\mathbb{E}\xi = \int_{\Omega_h} \xi(\omega) dm_H^h = P_{\zeta,h}(A). \tag{2.14}$$

Moreover, by Lemma 9,

$$\lim_{N \rightarrow \infty} \hat{P}_N(A) = P_h(A). \tag{2.15}$$

In view of Lemma 6 and the Birkhoff-Khintchine theorem, for almost all $\omega \in \Omega_h$,

$$\lim_{N \rightarrow \infty} \frac{1}{N} \sum_{k=1}^N \xi(\varphi_h^k(\omega)) = \mathbb{E}\xi.$$

Since $w \in W$, from this it follows that, for almost all $\omega \in \Omega_h$,

$$\lim_{N \rightarrow \infty} \frac{1}{V_N} \sum_{k=1}^N w(k) \xi(\varphi_h^k(\omega)) = \mathbb{E}\xi. \tag{2.16}$$

However, by the definition of φ_h ,

$$\frac{1}{V_N} \sum_{k=1}^N w(k) \xi(\varphi_h^k(\omega)) = \frac{1}{V_N} \sum_{k=1}^N w(k) I_{\{k: \zeta_h(s+ikh, \omega) \in A\}}(k) = \hat{P}_{N,h}(A).$$

Therefore, by (2.14) and (2.16),

$$\lim_{N \rightarrow \infty} \hat{P}_{N,h}(A) = P_{\zeta,h}(A).$$

This and (2.15) show that $P_h = P_{\zeta,h}$.

For finding the support of $P_{\zeta,h}$, we use the representation (2.13). For $p \in \mathbb{P} \setminus \mathbb{P}_0$, the random variables $\omega(p)$ are independent. Thus, by the proof of Lemma 6.5.5 from [11], we find that the support of the random element X_2 is the set S . Since the random elements X_1 and X_2 are independent and X_1 is not degenerate at zero, we obtain that the support of $X_1 X_2$ is the set S , i.e., the support of the measure $P_{\zeta,h}$ is the set S . The theorem is proved. \square

3 Proof of universality theorems

Theorems 4 and 5 follow from the limit theorems (Theorems 6 and 7), for $\zeta(s)$ as well as from the Mergelyan theorem [19] on the approximation of analytic functions by polynomials.

Proof. (Of Theorem 4). By the Mergelyan theorem, there exists a polynomial $p(s)$ such that

$$\sup_{s \in K} |f(s) - e^{p(s)}| < \frac{\varepsilon}{2}. \quad (3.1)$$

For brevity, denote the limit measure in Theorems 6 and 7 by \hat{P}_ζ , i.e.,

$$\hat{P}_\zeta = \begin{cases} P_\zeta, & \text{if } h \text{ is of type 1,} \\ P_{\zeta, h}, & \text{if } h \text{ is of type 2,} \end{cases} \quad \hat{P}_N = \begin{cases} P_N, & \text{if } h \text{ is of type 1,} \\ P_{N, h}, & \text{if } h \text{ is of type 2.} \end{cases}$$

Then we have that \hat{P}_N converges weakly to \hat{P}_ζ as $N \rightarrow \infty$. Define the set

$$G_\varepsilon = \left\{ g \in H(D) : \sup_{s \in K} |g(s) - e^{p(s)}| < \frac{\varepsilon}{2} \right\}.$$

Since $e^{p(s)} \neq 0$, and, in view of Theorems 6 and 7, the support of the measure \hat{P}_ζ is the set S , the set G_ε is an open neighbourhood of an element of the support, therefore,

$$\hat{P}_\zeta(G_\varepsilon) > 0. \quad (3.2)$$

Moreover, by the first parts of Theorems 6 and 7, and the equivalent of weak convergence of probability measures in terms of open sets [2, Theorem 2.1], we have that

$$\liminf_{N \rightarrow \infty} \hat{P}_N(G_\varepsilon) \geq \hat{P}_\zeta(G_\varepsilon).$$

This, (3.2) and the definitions of \hat{P}_N and G_ε show that

$$\liminf_{N \rightarrow \infty} \frac{1}{V_N} \sum_{k=1}^N w(k) I_{\left\{ k: \sup_{s \in K} |\zeta(s+ikh) - e^{p(s)}| < \frac{\varepsilon}{2} \right\}}(k) > 0. \quad (3.3)$$

It remains to replace $e^{p(s)}$ by $f(s)$ in the latter inequality. Suppose that k satisfies the inequality

$$\sup_{s \in K} |\zeta(s+ikh) - e^{p(s)}| < \frac{\varepsilon}{2}.$$

Then, in virtue of (3.1), the same k satisfies the inequality

$$\sup_{s \in K} |\zeta(s+ikh) - f(s)| < \varepsilon.$$

Therefore,

$$\left\{ k : \sup_{s \in K} |\zeta(s+ikh) - e^{p(s)}| < \frac{\varepsilon}{2} \right\} \subset \left\{ k : \sup_{s \in K} |\zeta(s+ikh) - f(s)| < \varepsilon \right\}.$$

This inclusion together with (3.3) proves the theorem. \square

Proof. (Of Theorem 5). Define the set

$$\hat{G}_\varepsilon = \left\{ g \in H(D) : \sup_{s \in K} |g(s) - f(s)| < \varepsilon \right\}.$$

Then $\partial \hat{G}_\varepsilon = \left\{ g \in H(D) : \sup_{s \in K} |g(s) - f(s)| = \varepsilon \right\}$ is the boundary of \hat{G}_ε . Hence, $\partial \hat{G}_{\varepsilon_1} \cap \partial \hat{G}_{\varepsilon_2} = \emptyset$ if $\varepsilon_1 \neq \varepsilon_2$, $\varepsilon_1 > 0$, $\varepsilon_2 > 0$. Therefore, the set $\partial \hat{G}_\varepsilon$ can have a positive \hat{P}_ζ -measure for at most countably many $\varepsilon > 0$. This means that the set \hat{G}_ε is a continuity set of the measure \hat{P}_ζ for all but at most countably many $\varepsilon > 0$. Using Theorems 6 and 7, and the equivalent of weak convergence of probability measures in terms of continuity sets [2, Theorem 2.1], we have that

$$\lim_{N \rightarrow \infty} \hat{P}_N(\hat{G}_\varepsilon) = \hat{P}_\zeta(\hat{G}_\varepsilon) \tag{3.4}$$

for all but at most countably many $\varepsilon > 0$. Moreover, (3.1) shows that $G_\varepsilon \subset \hat{G}_\varepsilon$. Therefore, by (3.2), $\hat{P}_\zeta(\hat{G}_\varepsilon) > 0$. This, (3.4) and the definition of the set \hat{G}_ε prove the theorem. \square

Acknowledgements

The research of the first author is funded by the European Social Fund according to the activity “Improvement of researchers’ qualification by implementing world-class R&D projects” of Measure No. 09.3.3-LMT-K-712-01-0037.

References

- [1] B. Bagchi. *The Statistical Behavior and Universality Properties of the Riemann Zeta-Function and Other Allied Dirichlet Series*. PhD Thesis, Indian Statistical Institute, Calcutta, 1981.
- [2] P. Billingsley. *Convergence of Probability Measures*. Wiley, New York, 1968.
- [3] K.M. Bitar, N.N. Khuri and H.C. Ren. Path integrals and Voronin’s theorem on the universality of the Riemann zeta-function. *Ann. Phys.*, **211**(1):172–196, 1991. [https://doi.org/10.1016/0003-4916\(91\)90196-F](https://doi.org/10.1016/0003-4916(91)90196-F).
- [4] H. Bohr. Über das verhalten von $\zeta(s)$ in der halbebene $\sigma > 1$. *Nachr. Akad. Wiss. Göttingen II Math. Phys. Kl.*, pp. 409–428, 1911.
- [5] H. Bohr and R. Courant. Neue anwendungen der theorie der diophantischen approximationen auf die Riemannsche zetafunktion. *Reine Angew. Math.*, **1914**(144):249–274, 1914. <https://doi.org/10.1515/crll.1914.144.249>.
- [6] A. Dubickas and A. Laurinćikas. Distribution modulo 1 and the discrete universality of the Riemann zeta-function. *Abh. Math. Semin. Univ. Hamb.*, **86**(1):79–87, 2016. <https://doi.org/10.1007/s12188-016-0123-8>.
- [7] R. Garunkštis and A. Laurinćikas. Riemann hypothesis and universality of the Riemann zeta-function. *Math. Slovaca*, **68**(4):741–748, 2018. <https://doi.org/10.1515/ms-2017-0141>.

- [8] R. Garunkštis, A. Laurinčikas and R. Macaitienė. Zeros of the Riemann zeta-function and its universality. *Acta Arith.*, **181**:127–142, 2017. <https://doi.org/10.4064/aa8583-5-2017>.
- [9] S.M. Gonek. *Analytic Properties of Zeta and L-Functions*. PhD Thesis, University of Michigan, 1979.
- [10] A. Laurinčikas. On the universality of the Riemann zeta-function. *Lith. Math. J.*, **35**(4):399–402, 1995. <https://doi.org/10.1007/BF02335599>.
- [11] A. Laurinčikas. *Limit Theorems for the Riemann Zeta-Function*. Kluwer, Dordrecht, 1996.
- [12] A. Laurinčikas. On the Matsumoto zeta-function. *Acta Arith.*, **84**:1–16, 1998. <https://doi.org/10.4064/aa-84-1-1-16>.
- [13] A. Laurinčikas. Discrete universality of the Riemann zeta-function and uniform distribution modulo 1. *St. Petersburg Math. J.*, **30**:103–110, 2019. <https://doi.org/10.1090/spmj/1532>.
- [14] A. Laurinčikas, K. Matsumoto and J. Steuding. Discrete universality of L -functions of new forms. II. *Lith. Math. J.*, **56**(2):207–218, 2016. <https://doi.org/10.1007/s10986-016-9314-3>.
- [15] A. Laurinčikas and J. Petuškinaitė. Universality of Dirichlet L -functions and non-trivial zeros of the Riemann zeta-function. *Sb. Math.*, **210**, 2019. <https://doi.org/10.1070/SM9194>.
- [16] R. Macaitienė. On discrete universality of the Riemann zeta-function with respect to uniformly distributed shifts. *Arch. Math.*, **108**(3):271–281, 2017. <https://doi.org/10.1007/s00013-016-0998-8>.
- [17] R. Macaitienė, M. Stoncelis and D. Šiaučiūnas. A weighted discrete universality theorem for periodic zeta-functions. II. *Math. Modell. Analysis*, **22**(6):750–762, 2017. <https://doi.org/10.3846/13926292.2017.1365779>.
- [18] R. Macaitienė, M. Stoncelis and D. Šiaučiūnas. A weighted universality theorem for periodic zeta-functions. *Math. Modell. Analysis*, **22**(1):95–105, 2017. <https://doi.org/10.3846/13926292.2017.1269373>.
- [19] S.N. Mergelyan. Uniform approximations to functions of complex variable. *Usp. Mat. Nauk.*, **7**:31–122, 1952 (in Russian).
- [20] H.L. Montgomery. *Topics in Multiplicative Number Theory*. Lecture Notes Math. Vol. 227, Springer-Verlag, Berlin, 1971. <https://doi.org/10.1007/BFb0060851>.
- [21] Ł. Pańkowski. Joint universality for dependent L -functions. *Ramanujan J.*, **45**(1):181–195, 2018. <https://doi.org/10.1007/s11139-017-9886-5>.
- [22] A. Reich. Werteverteilung von zetafunktionen. *Arch. Math.*, **34**(1):440–451, 1980. <https://doi.org/10.1007/BF01224983>.
- [23] A.N. Shiryayev. *Probability*. Graduate Texts Math. vol. 95, Springer-Verlag, New York, 1984.
- [24] J. Steuding. *Value-Distribution of L-Functions*. Lecture Notes Math. vol. 1877, Springer, Berlin, Heidelberg, 2007.
- [25] S.M. Voronin. On the distribution of nonzero values of the Riemann ζ -function. *Trudy Mat. Inst. Steklov*, **128**:131–150, 1972 (in Russian).
- [26] S.M. Voronin. Theorem on the “universality” of the Riemann zeta-function. *Izv. Akad. Nauk SSSR, Ser. Matem.*, **39**:475–486, 1975 (in Russian).