

Journal of Maps



ISSN: (Print) 1744-5647 (Online) Journal homepage: https://www.tandfonline.com/loi/tjom20

Plants and animals in the oikonyms of Lithuania

Giedrė Beconytė, Julius Donatas Budrevičius, Irena Ciparytė & Andrius Balčiūnas

To cite this article: Giedrė Beconytė, Julius Donatas Budrevičius, Irena Ciparytė & Andrius Balčiūnas (2019) Plants and animals in the oikonyms of Lithuania, Journal of Maps, 15:2, 726-732, DOI: 10.1080/17445647.2019.1663282

To link to this article: https://doi.org/10.1080/17445647.2019.1663282

© 2019 The Author(s). Published by Informa UK Limited, trading as Taylor & Francis Group on behalf of Journal of Maps	→ View supplementary material 🗹
Published online: 16 Sep 2019.	Submit your article to this journal 🗗
Article views: 285	View related articles ☑
View Crossmark data 🗹	Citing articles: 2 View citing articles





Science

3 OPEN ACCESS



Plants and animals in the oikonyms of Lithuania

Giedrė Beconytė D, Julius Donatas Budrevičius, Irena Ciparytė and Andrius Balčiūnas

Department for Cartography and Geoinformatics, Vilnius University, Vilnius, Lithuania

ABSTRACT

Two toponymical maps presented in this paper show 2332 inhabited places of Lithuania that have names (oikonyms) associated with vegetation and animals. The maps and the dataset are the outcomes of a project that is aimed at combining the outcomes of professional onomastic research with an environment that fosters exploration. The oikonyms were extracted from the reference base dataset of Lithuania by means of an automated algorithm. Original cartographic symbols have been designed for depicting categories and species. The multiscale map application, with its exploratory tools, makes it easy to see spatial distribution of geographic names related with particular groups of plants and animals. Analysis of map data enables the assertion that local toponymy reflects a distribution of species that were characteristic to the territory over past centuries. The maps are supplemented by comparative density maps and statistical charts. The reference scale of the main maps is 1: 500,000.

ARTICLE HISTORY

Received 28 June 2019 Accepted 30 August 2019

EYWORDS

Geographical names; toponyms; oikonyms; plants; animals; origin

1. Introduction

Toponymical maps represent names of inhabited places, hydrography, forests, forms of relief, natural and cultural heritage and other features that were given names by people – geographical names or toponyms. The toponyms reflect the characteristics of spatial objects that were important for local people and sometimes remain the only source of information about these characteristics after the named objects ceased to exist. The importance of place names as a carrier of history and heritage of the place has been demonstrated in various contexts (Chloupek, 2018; Hakala, Sjöblom, & Kantola, 2015; Qian, Kang, & Weng, 2016; Tucci, Ronza, & Giordano, 2011). Toponyms may be included in the official databases or remain informal. The names of inhabited places (oikonyms) of Lithuania often associate with nature, specifically the names of plants, mushrooms, fish, birds and mammals that take very diverse and characterful forms as toponyms. Several published onomastic studies focus on the origins of Lithuanian toponyms (Grybauskienė, 2004; Kačinaitė, 2008; Vanagas, 1981; Vanagas, 2004).

Information on origins is presented in dictionaries of place names (Razmukaitė & Pupkis, 2002; Vietovardžių žodynas, 2014). There are interesting studies on particular groups of Lithuanian toponyms associated with living nature: myconyms (Lubienė, 2009, 2012), phytonyms (Genelytė, 2001; Gritėnienė, 2006) and zoonyms (Auksoriūtė, 1995; Gaivenis, 1995).

However, to date there have been no serious attempts to study peculiarities of the spatial distribution of toponyms. The development of spatial data infrastructure since 2003 (Beconytė & Kryžanauskas, 2010; Beconytė, Govorov, Ningal, Paršeliūnas, & Urbanas, 2008) and the launching of a Lithuanian spatial information portal (geoportal.lt) in 2009 led to better availability of the toponymical data, both within spatial datasets and on the historical maps in raster format. In 2014 the National Land Service under the Ministry of Agriculture and the state enterprise 'GIS-Centras' published a toponymical gazetteer service compliant with the requirements of Directive 2007/2/EC of the European Parliament and of the European Council of 14 March 2007 (INSPIRE). For this purpose a joint toponymy dataset was generated from several official sources: reference base datasets, state cadastres, the road information system and the scientific database of the Institute for Lithuanian Language. An internet application facilitating work with toponymical data was developed, available for viewing free of charge by the public (Viliuvienė, 2015). In 2015, a spatial toponymy data set was made publicly available for download on geoportal.lt. It is natural that over that decade the interest of Lithuanian residents in toponyms increased. The interest manifested itself with discussions on social media platforms on the common origins of Lithuanian toponyms.

Several geographers raised the issue of lost toponyms (Kavoliutė, 2014). Over the twentieth century, place names on the territory of Lithuania changed because of historical events, including mass

collectivization, land reclamation, land occupation and Sovietization. Some authors contend that during the period 1959-1989 over 5,500 names of abolished homesteads and hamlets were removed from the official data set of inhabited place names (Trakymas & Vaitekūnas, 2006). More recently, as a result of depopulation, some settlements were merged or abolished. According to the data provided by the Centre of Registers, the names of 240 inhabited places were removed from the national register of Addresses in the period 2010–2015 (Viliuvienė, 2015).

The Lithuanian Society of Geographers is calling for action to preserve the oikonyms as a part of the nation's intangible heritage. The year 2019 has been announced as the Year of Toponyms in Lithuania. The authors of this paper decided to contribute to the toponymical initiatives underway by adding a geographic dimension - with expectations of discovering what the pattern of oikonyms may reveal. As a significant part of Lithuanian oikonyms are immediately associated with the names of plants, mushrooms or animals, that, in turn, may reflect past or present characteristics of the environment, we plotted them on maps in order to explore their distribution over the territory and their possible relationships with local fauna and flora.

Two maps presented in this paper (Main Maps) show 2,332 inhabited places of Lithuania that have names associated with plants (691 oikonyms, or 29.5% of the total number of target oikonyms) or plant communities (584 oikonyms, 25%), fungi (36 oikonyms, 1.5%) and fauna (1,021 oikonym, 44%). The maps have been designed for printing at scale 1:500,000. The digital dataset can be explored online at https://lietuvoskartografija. lt/mapping-oikonyms-in-lithuania.

2. Project goal and related work

The initial goal of the project was to demonstrate the power of cartography for dissemination of knowledge, in our case on national toponyms. For this, maps have to meet three requirements:

- (a) the represented information must be of public interest;
- (b) the data, not necessarily scientific, must be comprehensive, systematically selected and carefully processed; and
- (c) the cartographic design must be original and attractive, and the system of symbols consistent and associative.

The maps presented thus continue the series of 'curious' thematic maps designed with this purpose at the Institute of Geosciences of Vilnius University (Beconyte, Alekna, & Rociute, 2011; Beconytė, Eismontaitė, Kuodytė-Dūdė, & Žemaitienė, 2014; Beconytė,

Maneikaitė, Bružas, & Balčiūnas, 2015; Vasiliauskas & Beconytė, 2016). Two popular map series that inspired our work on toponyms must be mentioned. The maps of the Atlas of True Names (Hormes, 2012; Hormes & Preust, 2008) show the native language meanings of familiar present-day geographic names. The atlas contains about three thousand names of cities, countries, rivers, oceans and mountain ranges of the world, Europe, the British Isles, Canada and the United States. As the meanings are presented as texts, maps can be explored by reading individual names but cannot be used for visual analysis. A series of maps designed by Steven Kay (Kay, 2016) shows the patterns of distribution for some place names in Britain by origin (e.g. Old British, Welsh, Pictish). The data is not consistent. Hexagonal choropleths only show simple statistics but an original design and the clarity of patterns make the maps attractive and memorable. In our approach we combine the individual place names and pictograms that show the taxons of the associated species. Thus the user can quickly identify a pattern and then proceed with analysis of individual names that generate that pattern. The online application developed allows for searching by the place name and by the name of a taxon as well as browsing the available scientific data about the name's origin.

Another goal was to elucidate the spatial distribution and make informed assertions concerning the spatial patterns and the semantic relationships of oikonyms with the real-life characteristics of the environment. We strongly support the statement by Jan Tent about the possibilities of extensive (geographic) research in toponymy (Tent, 2015), and share his observation that, although digital toponymy data sources are widely available and research itself is 'more straightforward to conduct' than intensive (etymological) research, only a small part of toponymic studies that actually exist are by their very nature extensive. Extensive studies often focus on the relationship between toponyms and the natural (landscape) elements (Capra, Ganga, Filzmoser, Gaviano, & Vacca, 2016; Cox, Maehr, & Larkin, 2002; Derungs, Wartmann, Purves, & Mark, 2013; Shi, Ren, Du, & Gao, 2015; Tucci et al., 2011). Moreover, the links between linguistic and biological diversity have been demonstrated (Axelsen & Manrubia, 2014; Fagundez & Izco, 2016a,b; Grant, 2012; Harmon, 1996; Sousa & García-Murillo, 2001; Sutherland, 2003). The term 'toponymic species' coined by Fagundez and Izco (2016b) reflects the relationship between the more persistent place names and local species that change and can disappear over time. As stated by the authors, toponyms are 'stable, spatially-explicit elements that may be used as indicators of bio-cultural diversity'.

Mapping oikonyms that relate to specific species is the first step towards answering the research question: does the distribution of toponyms in Lithuania reflect

Table 1. The types and numbers of oikonyms in the database.

		Example	
Number of toponyms	Number of taxons or types	Oikonym	Associated word
691	75	Pušinėlė	Pušis (a pine; Pinus)
36	4	Gryblaukis	Grybas (a mushroom; Fungus)
584	6	Pašilė	<i>Šilas</i> (a pinewood)
999	95	Gegutė	Gegutė (a cuckoo; Cuculus canorus)
22	2	Žiogai	<i>Žiogas</i> (a grasshopper)
2332	182	J	
	691 36 584 999 22	691 75 36 4 584 6 999 95 22 2	691 75 Pušinėlė 36 4 Gryblaukis 584 6 Pašilė 999 95 Gegutė 22 2 Žiogai

past biodiversity? The same question can be raised concerning the distribution of particular toponymic species at present and in the past. Indeed, if 'place names function as independent variables which can be tested against dependent variables...' (Tent, 2015), the research would be possible and could potentially yield valid results.

In the context of toponymy research, our work, being an extensive study, does not attempt to investigate the original meanings of individual toponyms. Instead, it focuses on toponyms (oikonyms) that associate with the names of plants and animals by way of the same radical elements. It is aimed at inspiring geographic analyses and facilitating such analyses by utilization of the tools provided by the online application developed. Some scholars of onomastics have expressed an objection to this idea, because a significant part of our toponymy data is linked to common names without providing detailed proofs for the assertions concerning their origins. We will touch upon this criticism and make remarks on it further on in the text. On the other hand, other researchers (including the authors of this paper) believe that this project has the potential to integrate with the extant and extensive geographic and onomastic research and to foster more such research in the future.

3. Data and methods

The oikonyms were selected from the Lithuanian reference base dataset at scale 1:10,000 that contains exhaustive data on the presently inhabited sites in the country - over 20,000 of oikonyms.

The scope of the study entails the set of oikonyms in Lithuania that associate with the names of species or higher taxons of biological kingdoms of plants, fungi and animals. In addition, we included the toponyms that originate from two groups of informal, commonly used names:

- (a) generic, such as a beast, a beetle and
- (b) collective, such as a garden, a grove, a meadow.

According to the toponym taxonomy developed by Tent and Blair (2011), the target toponyms are mostly

associative. Descriptive toponyms that indicate an inherent characteristic of the feature, e.g. Paberžynė (literally, '[the place] near the birch grove'), Padidmiškis ('[the place] near the great wood'), Užugiris ('[the place] behind the wood') account for only a small part of the set - about 200 toponyms. The types and numbers of the represented oikonyms are shown in Table 1.

The initial selection of oikonyms from the reference base dataset was partially automated, based upon comparison of radical elements of the oikonyms and a thesaurus containing the names of species. Due to the complexity of the Lithuanian language, the dialectal variation, homonymy and various transformations of common names during the process of optimization, the percent of errors was high – about 20% false positives and about 5% false negatives. This was not acceptable. Therefore, several iterative manual revisions followed. Then the data set was verified against the available publications and scientific toponymy databases that contain explanations of origin: the Geoinformational Database of Lithuanian Toponyms (2013) and the Online Database of Lithuanian Last Names (2018) that helped to track the links to common words in the cases of transonimization.

Two attributes from the national toponymy dataset, an explanation of origin and explanation of formation of the toponym, provided by the Institute for Lithuanian Language, were linked to the collected oikonym data. Unfortunately, only about 12% of the zoo-botanical oikonyms had these attributes' values provided.

In quite a few cases, toponyms that have the same roots as the names of birds, animals or plants originate from them indirectly, for example, through personal names. The toponyms may be directly or indirectly derived from the common names that share the same root, but have no relationship with fauna and flora at all – the associations are only due to distortions or to homonymy. Deeper etymological studies are necessary for revealing the true origin of such names. For example: the name of the village Pušėnai, located in a wooded area of southern Lithuania, most likely originates from the distorted personal name Puišys, not directly from the tree name pušis (a pine tree) as common sense would suggest (Palionis, 2008); the name




Figure 1. The most frequent names of plants in the oikonyms of Lithuania.

Arklėnai, though immediately associated with arklys ('horse') is, most likely derived from this word indirectly, via a personal name that means a man who is strong and sturdy as a horse. The oikonym Bitėnai derives from the personal name Bitenas that originates from bitė ('bee'). Such oikonyms were not included in our database in cases of reliable evidence of transonimization. As the identification of the actual local origin of each individual toponym is a resourceconsuming and sometimes utterly impossible task, we decided to include all the oikonyms that could have been derived from the names of plants, fungi and animals even thought their origin is arguable and incapable of demonstration. Being aware of that, users are encouraged to suggest alternative explanations. We expect to receive readers' negations of the claimed zoo-botanical origin of some names. That would demonstrate the interest of users in the dataset and maps. The web application provides the tools for users to submit their comments regarding individual names. If the dataset is maintained as planned, the number of false attributions will gradually decrease.

Why only oikonyms? The original selection algorithm would work equally well for other Lithuanian toponyms. As noted above, the semantic associations of oikonyms are often not primary, they are rather transferred not only from personal names but also from hydronyms. For example, the oikonym *Palokis* ('[the place] near *Lokys*'), derived from the name of the rivulet *Lokys* ('bear'), were among the oikonyms

included in our database. All the registered hydronyms of Lithuania are stored in the official cadastre. The main reason for postponing the inclusion of hydronyms is the geometry of the features (mainly lines) that requires visualization solutions different from those for the point geometry of oikonyms. Oronyms, names of forests, names of islands and various microtoponyms of Lithuania also very often have semantic associations with fauna and flora, but they are scattered across multiple datasets. If the oikonym project succeeds in raising the level of interest, other groups of toponyms will be added subsequently.

4. The results

The dataset contains 2,332 records of oikonyms that associate with the names of animals, plants and fungi. These oikonyms comprise 11% of the 21,171 residential areas of Lithuania. They are all represented on two maps.

The charts show the numbers of the most frequent (10 or more occurrences) common names of plants (12 species, Figure 1) and animals (20 species, Figure 2) in the oikonyms of Lithuania. Of fungi, only the generic name *Grybas* (Mushroom) has 18 occurrences.

The initial idea was to group and represent all the names by the corresponding biological taxons. However, a simple survey (80 persons were asked to comment on proposed classifications) revealed that whereas biological classification of animals is intuitive




Figure 2. The most frequent names of animals in the oikonyms of Lithuania.

and acceptable to typical users, the grouping of plants requires a different approach. The physical appearance of many common plants differs within the same family (e.g. apple tree, rowan, raspberry and several herbaceous plants belong to the same Rosaceae family). Moreover, a large portion of plant-related names are names of plant communities (e.g. a pinewood) or generic names that represent physical properties (e.g. a tree, a flower). For this reason the plants were informally grouped into relatively large groups according to physical appearance, use and habitat: trees and shrubs (conifers, fruiters and other leafy trees); herbaceous/flowering plants; water plants; crops and vegetables.

Cartographic symbols represent three levels of grouping:

- (i) biological kingdom level, represented by shape: hexagons for animals, circles for plants and fungi;
- (ii) class for animals / group for plants, represented by the colour;
- (iii) family for animals / subgroup for plants, represented by individual pictograms.

Collective and generic names are represented by abstract pictograms (Figure 3).

On the maps, it is easy to see the spatial distribution of names related with particular groups of plants and animals - they are represented by different colours. The maps are supplemented with comprehensive legends, hexagonal grid density maps, and statistical charts. On the background main hydrographic features, forest areas, roads and railways and the five



Figure 3. Examples of representation of concrete, generic and collective names.

ethnographic regions of Lithuania are represented. Hexagonal grid maps with the cells of 15 km² containing up to 19 inhabited places allow for visual comparison of the pattern of distribution of the oikonyms related with plants or animals vs. all oikonyms. The number of target oikonyms in a cell is represented by thickness of the cell contour while the value of the cell background colour shows the total number of named places. From these maps it is obvious that the pattern of distribution of the oikonyms related with animals and plants is different from the overall pattern of settlements. Deeper analysis of the map data allows for asserting that local toponymy reflects the distribution of species that were characteristic to the territory in past centuries. Wooded areas in northwestern and southeastern Lithuania and, specifically, the regions with high landscape fragmentation still contain larger number of and larger diversity of the placenames associated with species.

It must be said that there also exists a collection of historical names of settlements, most of which are no longer extant; however, this collection is still far from exhaustive. In the future, we would consider adding historical oikonyms as well as hydronyms and oronyms to the web application.

5. Discussion and concluding remarks

The presented maps constitute one part of a continuous project. Hydronyms, names of forests' and relief forms' will be investigated and the data used to test various hypotheses concerning the characteristics of a place or landscape that the toponymy reflects, and, how this reflection is transformed over time. The maps not only serve to underpin common beliefs (correlation between landscape diversity and density of oikonyms related with the species; reflection of prevalent species in oikonyms etc.) but they may also reveal some unexpected things. To take some examples, birch and lime trees prevail over the more common pine and spruce trees; there is an abundance of corvids while there is no single occurrence of stork, considered to be a national bird of Lithuania. It would be interesting to compare the numbers and spatial patterns of such oikonyms in different countries.

The biological names that associate with the oikonyms are presented in different ways on the two maps: only biological taxons for the animals, and, both taxons and common terms for the plants and fungi. We expect comments from users concerning the usability of these two maps.

The web application was designed with the intent to test the statement that 'place names give rise to feelings of individual and collective identity attached to the places in question' (Helleland, 2012). We believe that users will study various aspects of the data and possibly contribute to the development of an attribute-rich and linguistically tenable spatial inventory of Lithuanian toponyms.

Software

The spatial data was collected and maintained in the ESRI ArcGIS geodatabase. ArcGIS 10.6 was used for data processing. Adobe Illustrator and Adobe Photoshop graphic design software was used for the design of cartographic signs and charts. ArcGIS 10.6 was used for the final cartographic design.

Acknowledgements

We thank Aistė Pangonytė and Dalia Sviderskienė for valuable remarks and information that helped us to improve the data. Thanks are due to the managers of the national spatial information portal (www.geoportal.lt) for enabling unrestricted access to interoperable reference data and the tools that facilitate working with the toponymy datasets. The names of students of Vilnius University who greatly contributed to the research and the design of the maps are listed on the corresponding maps.

Disclosure statement

No potential conflict of interest was reported by the authors.

ORCID

Giedrė Beconytė http://orcid.org/0000-0003-3036-0544

References

Auksoriūtė A. (1995). Klaipėdos krašto paukščių pavadinimai. Lietuvininku žodis (parengė K. Grigas, Z. Kelmickaitė, B. Kerbelytė, N. Vėlius). Kaunas: Litterae universitatis, 514-520.

Axelsen, J. B., & Manrubia, S. (2014). River density and landscape roughness are universal determinants of linguistic diversity. Proceedings of the Royal Society B: Biological Sciences, 281, 20141179. doi:10.1098/rspb.2013.3029

Beconyte, G., Alekna, V., & Rociute, I. (2011). A map of 21st century conflicts in Europe. Journal of Maps, 7, 1-8. doi:10.4113/jom.2011.1166

Beconytė, G., Eismontaitė, A., Kuodytė-Dūdė, J., & Žemaitienė, J. (2014). Mythical creatures of Europe. Journal of Maps, 10(1), 53-60. doi:10.1080/17445647. 2013.867544

Beconytė, G., Govorov, M., Ningal, T. F., Paršeliūnas, E., & Urbanas, S. (2008). Geographic information e-training initiatives for national spatial data infrastructures. Technological and Economic Development of Economy. Baltic Journal on Sustainability, 14(1), Vilnius: Technika, p. 11-28. doi:10.3846/2029-0187.2008.14.11-28

Beconytė, G., & Kryžanauskas, A. (2010). Geographic communication for sustainable decisions / Geografinės informacijos Komunikavimo Priemonės ir Tvarioji Plėtra. Technological and Economic Development of Economy, 16(4), 603-612. doi:10.3846/tede.2010.37

Beconytė, G., Maneikaitė, E., Bružas, D., & Balčiūnas, A. (2015). A map of astronomy in Europe. Journal of Maps, 11(3), 515-524. doi:10.1080/17445647.2014.944943



- Capra, G. F., Ganga, A., Filzmoser, P., Gaviano, C., & Vacca, S. (2016). Combining place names and scientific knowledge on soil resources through an integrated ethnopedological approach. CATENA, 142, 89-101. doi:10.1016/j. catena.2016.03.003
- Chloupek, B. R. (2018). A GIS approach to cultural and historical toponymic research in Nebraska. Journal of Cultural Geography, 35(1), 23-43. doi:10.1080/08873631. 2017.1317182
- Cox, J. J., Maehr, D. S., & Larkin, J. L. (2002). The biogeography of faunal place names in the United States. Conservation Biology, 16, 1143-1150. doi:10.1046/j.1523-1739.2002.01202.x
- Derungs, C., Wartmann, F., Purves, R. S., & Mark, D. M. (2013). The meanings of the generic parts of toponyms: Use and limitations of gazetteers in studies of landscape terms. In T. Tenbrink, J. Stell, A. Galton, & Z. Wood (Eds.), Spatial information theory. Lecture Notes in Computer Science (Vol. 8116, pp. 261-278). Springer International. doi:10.1007/978-3-319-01790-7_15
- Fagúndez, J., & Izco, J. (2016a). Spatial analysis of heath toponymy in relation to present-day heathland distribution. International Journal of Geographical Information Science, 30(1), 51-60. doi:10.1080/13658816. 2015.1017729
- Fagúndez, J., & Izco, J. (2016b). Diversity patterns of plant place names reveal connections with environmental and social factors. Applied Geography, 74, 23-29. doi:10. 1016/j.apgeog.2016.06.012
- Gaivenis, K. (1995). Kai kurių žuvų pavadinimai. Terminologija 2. Vilnius: Lietuvių kalbos institutas, 86–92.
- Genelytė, A. (2001). Augalų pavadinimai, reiškiami pastoviaisiais žodžių junginiais. Lituanistica, 3(47), 31–38.
- Geoinformational Database of Lithuanian Toponyms. (2013). Institute for Lithuanian Language. Retrieved from http:// lkiis.lki.lt/lietuvos-vietovardziu-geoinformacine-duomenu-
- Grant, C. (2012). Analogies and links between cultural and biological diversity. Journal of Cultural Heritage Management and Sustainable Development, 2, 153-163. doi:10.1108/20441261211273644
- Gritėnienė, A. (2006). Augalų pavadinimų motyvacija šiaurės panevėžiškių patarmėje. Vilnius: Lietuvių kalbos institutas.
- Grybauskienė, V. (2004). Lietuvių mitologiniai vietovardžiai: iš leksemos laumė kilę vietų vardai. Tautosakos Darbai, XXI(XXVIII), 92-102.
- Hakala, U., Sjöblom, P., & Kantola, S.-P. (2015). Toponyms as carriers of heritage: Implications for place branding. Journal of Product & Brand Management, 24(3), 263-275. doi:10.1108/JPBM-05-2014-0612
- Harmon, D. (1996). Losing species, losing languages: Connections between biological and linguistic diversity. Southwest Journal of Linguistics, 15, 89–108.
- Helleland, B. (2012). Place names and identities. In B. Helleland, C.-E. Ore, & S. Wikstrøm (Eds.), Names and identities, Oslo studies in language (Vol. 4(2), pp. 95-116). Oslo: University of Oslo. (ISSN 1890-9639).
- Hormes, S. (2012). The atlas of true names. Lübeck: Kalimedia. Hormes, S., & Preust, S. (2008). Atlas der Wahren Namen – Europa. Etymologische Karte. Lübeck: Kalimedia.
- Kačinaitė, D. (2008). Lietuvių sudurtinių oronimų darybos bruožai. BALTISTICA, XLIII(2), 275-288.

- Kavoliutė, F. (2014). Gyvenamųjų vietovių vardai nematerialusis šalies kultūros paveldas. Geografijos Metraštis, 47, 88-102. ISSN 2335-8610.
- Kay, S. (2016). Toponymy mapping place name patterns in Great Britain with QGIS. Retrieved from https://www. stevefaeembra.com/blog/2016/1/10/toponymy-mappingplace-name-patterns-in-great-britain-with-qgis (2016-01-
- Lubienė, J. (2009). Lietuvių kalbos liaudiški grybų pavadinimai su grybų augimo motyvaciniais požymiais. Res Humanitariae VI. Klaipėda: Klaipėdos universiteto leidykla. 180-203.
- Lubienė, J. (2012). Žvėrių pavadinimai lietuvių kalbos mikonimų motyvacijos sistemoje. Res Humanitariae XII. Klaipėda: Klaipėdos universiteto leidykla. 99–121.
- Online Database of Lithuanian Last Names. (2018). Retrieved from http://pavardes.lki.lt
- Palionis, J. (2008). Dėl keleto retų vietovardžių kilmės. BALTISTICA, XLIII(2), 289–295.
- Qian, S., Kang, M., & Weng, M. (2016). Toponym mapping: A case for distribution of ethnic groups and landscape features in Guangdong, China. Journal of Maps, 12(sup1), 546-550. doi:10.1080/17445647.2016.1201017
- Razmukaitė, M., & Pupkis, A. (eds.). (2002). Lietuvos vietovardžių žodynas. Vilnius: Mokslo ir enciklopedijų leidybos institutas.
- Shi, G., Ren, F., Du, Q., & Gao, N. (2015). Phytotoponyms, geographical features and vegetation coverage in Western Hubei, China. Entropy, 17, 984-1006. doi:10. 3390/e17030984
- Sousa, A., & García-Murillo, P. J. (2001). Can place names be used as indicators of landscape changes? Application to the doñana natural park (Spain). Landscape Ecology, 16, 391-406. doi:10.1023/A:1017585101389
- Sutherland, W. J. (2003). Parallel extinction risk and global distribution of languages and species. Nature, 423, 276-
- Tent, J. (2015). Approaches to research in toponymy. Names, 63(2), 65-74. doi:10.1179/0027773814Z.0000000000103
- Tent, J., & Blair, D. (2011). Motivations for naming: The development of a toponymic typology for Australian pla-59(2), 67-89. doi:10.1179/ cenames. Names, 002777311X12976826704000
- Trakymas, A., & Vaitekūnas, S. (2006). Kaimas. Visuotinė lietuvių enciklopedija, T. IX, p. 133-124. Vilnius: Mokslo ir enciklopedijų leidybos institutas.
- Tucci, M., Ronza, R. W., & Giordano, A. (2011). Fragments from many pasts: Layering the toponymic tapestry of Milan. Journal of Historical Geography, 37(2011), 370-384. doi:10.1016/j.jhg.2011.02.001
- Vanagas, A. (1981). Lietuvių hidronimų etimologinis žodynas. Vilnius: Mokslas.
- Vanagas, A. (2004). Lietuvos miestų vardai. Vilnius: Mokslo ir enciklopedijų leidybos institutas.
- Vasiliauskas, D., & Beconytė, G. (2016). Cartography of crime: Portrait of metropolitan Vilnius. Journal of Maps, 12(5), 1236-1241. doi:10.1080/17445647.2015.1101404
- Vietovardžių žodynas. (2014). Vilnius: Lietuvių kalbos institutas. 2014. Retrieved from http://vietovardziai.lki.lt
- Viliuvienė, R. (2015). Place names in the national geoportal of Lithuania. UNGEGN Information Bulletin 2015, Vol.48. May 2015.