

Lyme disease and heart transplantation: presentation of a clinical case and a literature review

Audrius Aurelijus Pilypas¹,

Giedrutė Raišeliienė²,

Jurgita Valaikienė³

¹ Faculty of Medicine,
Vilnius University,
Vilnius, Lithuania

² Clinic of Cardiac and Vascular Diseases,
Vilnius University Hospital
Santaros Klinikos,
Vilnius, Lithuania

³ Centre of Neurology,
Vilnius University Hospital
Santaros Klinikos,
Faculty of Medicine,
Vilnius University,
Vilnius, Lithuania

Background. Lyme disease, the most common anthrozoosis, is a transmissible natural focal infection affecting various organs and systems. Also known as Lyme borreliosis, it is caused by *Borrelia spirochetes*, which are distributed by ticks of the genus *Ixodes*. Early diagnosis is difficult due to frequent occurrence of atypical symptoms, unnoticed tick bites, the absence of migratory erythematous lesions, and symptoms occurring during the non-tick season. If not diagnosed and treated in time, dissemination of the infection occurs and various complications develop since borrelias damage not only the skin but also the nervous system, joints, and, in rare cases, the heart and eyes.

Materials and methods. This article presents a clinical case of Lyme borreliosis-induced myocarditis, which led to the development of dilated cardiomyopathy and, consequently, urgent cardiac transplantation. According to our data, this is one of the first described cases of this complication in the world.

Results and conclusions. When diagnosed in time and treated properly, the prognosis of Lyme myocarditis is usually good. In most cases, the atrioventricular block disappears within 1–2 weeks of antibiotic treatment and the implantation of a temporary pacemaker is rarely needed. In those rare cases of a chronic *Borrelia burgdorferi* infection, dilated cardiomyopathy may develop; thus if a sudden atrioventricular block occurs, the physician should be vigilant and perform the necessary tests to exclude the diagnosis of Lyme disease.

Keywords: Lyme myocarditis, atrioventricular block, Lyme disease, dilated cardiomyopathy

*Correspondence to: Audrius Aurelijus Pilypas, Faculty of Medicine, Vilnius University, Centre of Cardiology and Angiology, Vilnius University Hospital Santaros Klinikos, Vilnius, Lithuania, Santariskiu St. 2, Vilnius 08661, Lithuania. E-mail: audrius.pilypas@santa.lt

INTRODUCTION

Lyme disease (LD) or Lyme borreliosis is an infectious multisystemic disease caused by bacteria of the *Borrelia* (B) genus of the spirochete family (*B. burgdorferi*, *B. afzelii*, *B. garinii*, etc.). Due to natural and climatic conditions, this disease prevails in the temperate zone of the northern hemisphere, especially in Europe and North America and so the risk of infection in these areas is extremely high (1, 2). According to the Lithuanian Centre for Infectious Diseases and AIDS (ULAC), LD affects 2,000-3,000 people in Lithuania annually (3). Although people who are in close contact with nature are at a greater risk of tick bites; ticks are also found in residential yards and lawns. In addition, a nymph's bite often remains unidentified, unlike that of an adult tick. These factors impede the diagnosis of tick-borne diseases, the most common of which is Lyme disease. No memory of a tick bite and atypical disease symptoms will delay the correct diagnosis, which can lead to very serious consequences. Because no vaccine exists for Lyme borreliosis and infection with the disease does not confer immunity, it is possible to become reinfected. This means that, potentially, anyone can contract LD and so doctors should be very vigilant. This article presents a clinical case of LD-induced myocarditis, which led to the development of dilated cardiomyopathy and, ultimately, urgent cardiac transplantation. According to our data, because the link between LD and advanced dilatative cardiomyopathy has only been established in Europe (4), this is one of the first reported cases of this complication in the world. The purpose of this article is to reveal the course of LD-induced myocarditis and long-term complications as well as to promote medical vigilance. In case of a sudden atrioventricular block of unclear aetiology, dilative cardiomyopathy and heart failure, tests to exclude Lyme Borreliosis need to be conducted.

CASE REPORT

A 45-year-old male in cardiogenic shock was brought to the emergency department by ambulance. At the time of the physical examination, the skin and mucous membranes were pale, the lips appeared to be cyanotic, and the skin was covered with cold sweat. The blood pressure (BP) was un-

measurable, the heart rate (HR) 20 bpm. Vesicular breath sounds with wet basal crackles were audible during the auscultation of the lungs. The anamnesis showed that the patient had been experiencing great weakness and dizziness for the last two months and fainted several times. The patient was taking no medication and had no allergies, history of chronic diseases, or memory of a tick bite. The medical records showed the patient had been treated by a dermatovenerologist for rashes on the inner surfaces of the thighs two years before. The patient was transferred to the Cardiac Intensive Care Unit where a temporary cardiac pacemaker was immediately implanted. Blood test data: white blood cells (WBC) – $8.7 \times 10^9/l$ (norm $4.0-9.8 \times 10^9/l$), neutrophils (NEU) – $7.47 \times 10^9/l$ (norm $1.5-7.0 \times 10^9/l$), haemoglobin (HGB) – 141.30 g/l (norm 128–160 g/l), platelets (PLT) – $132.30 \times 10^9/l$ (norm $140-450 \times 10^9/l$), troponin I – 0.81 ng/l (norm up to 34.2 ng/l), and creatinine – 118 $\mu\text{mol/l}$ (norm 64–104 $\mu\text{mol/l}$); the electrolyte, glucose, C-reactive protein (CRP), acid-base balance, and the urine test results were normal. The chest X-ray showed venous stasis in the lungs and a left ventricular enlargement. The urgent coronarography showed a catheter-induced spasm in the left coronary artery. The transthoracic echocardiographic examination showed good general and local inotropy, no valve disorder, an ejection fraction (EF) of 50%; the electrocardiogram (ECG) showed a third-degree atrioventricular (AV) block (Fig. 1). An examination failed to reveal the exact cause of the disorder and the AV block was treated with a permanent pacemaker. After his condition improved, the patient was discharged from the hospital.

Three years later, the patient contacted the emergency department due to general weakness and intense chest pain radiating to the left arm. These symptoms lasted for about two weeks. According to the data from the examination performed in the emergency department, vesicular breath sounds without crackles were audible during the auscultation; the heart activity was rhythmic, the heart rate 70 bpm, the arterial blood pressure 130/80 mmHg. The ECG showed effective cardiac pacemaker activity. The blood tests showed an increased mass of creatine kinase isoenzyme MB (CK-MB) – 31.33 $\mu\text{g/l}$ (norm $<5.2 \mu\text{g/l}$), while troponin I was normal at 0.323 ng/l (norm $<34.2 \text{ ng/l}$). Under suspicion of a myocardial infarction, the patient was




Fig. 1. The third degree atrioventricular block

hospitalized in the Department of Cardiology. The departmental blood tests showed a significant rise in the creatinine kinase (CK) at 1612 U/L (norm 25–195 U/L) and B-type natriuretic peptide (BNP) at 896.8 ng/l (norm <125 ng/l) but normal values of the CK-MB, troponin I, complete blood count (CBC), coagulogram, D-dimers, CRP, electrolytes, lipidogram, and liver enzymes. The coronary angiography performed due to the suspicion of a myocardial infarction showed no significant constrictions of the coronary blood vessels and so the preliminary diagnosis was rejected. An up to 30% reduction in EF was observed during cardiac echocardiography as well as a I° mitral regurgitation and a II° dilation of the left and right atria. A computer tomography of the resting myocardial perfusion performed using Technetium (99mTc) tetrofosmin with electrocardiography synchronization yielded a scintigraphic image of myocarditis, the ejection fraction at rest being 24%. The echocardiography of the dobutamine load performed on the tenth day of hospitalization showed diffuse contraction disorders characteristic of a non-ischemic cardiomyopathy. During the dobutamine infusion, the left ventricular EF increased from 26% to 40%, with significant signs of mechanical intraventricular asynchronism. After discussing the patient's condition and the laboratory and radiology tests at a case conference, myocarditis was diagnosed, but the cause remained unclear. After excluding many diseases, LD was suspected and a test for *Borrelia burgdorferi* antibodies was performed. Positive IgG and IgM antibody titres were detected using

an enzyme-linked immunosorbent assay (ELISA) test. During the detection of the *Borrelia* antibodies with immunoblot, the IgG titre was found to be positive. In the presence of positive serological test results, Lyme borreliosis was diagnosed. After performing an endomyocardial biopsy, chronic myocarditis with lymphocyte infiltration was also diagnosed, the final diagnosis being Lyme myocarditis. The following treatment was applied: Sol. Ceftriaxon 1 g twice a day intravenously for 28 days. The patient's condition became stable after antibiotic treatment. At the time of his discharge from the hospital, his heart activity was rhythmic, HR 70 bpm, BP 120/80 mmHg, and vesicular breath sounds without crackles were audible in the lungs during auscultation. It was recommended that a cardiologist perform an ultrasound examination and assess his general condition after the course of antibiotics. The examination showed the left ventricular EF to be 35% and the overall condition to have improved.

A year later, the patient's condition worsened again and he was hospitalized at the Department of Cardiology for severe general weakness. A transthoracic echocardiographic examination showed significant disorder of the left and right ventricular myocardial contraction, dilation of the left ventricular cavity, a thrombus at the top of left ventricle, a thrombus on the cardiac pacemaker cord in the right atrium cavity, II° left and III° right atrial dilation, III° tricuspid and mitral regurgitation, right heart cavity overload, systemic venous stasis, and an EF of 10% (Fig. 2). Dilated



Fig. 2. The thrombus at the top of the left ventricle

cardiomyopathy was diagnosed and a biventricular pacemaker implanted. As significant cardiac weakness remained, the patient was added to the list of urgent heart transplant recipients. Three months later, the patient underwent cardiac transplantation.

LITERATURE REVIEW AND DISCUSSION

Lyme disease is a transmissible, natural focal infection that affects various organs and systems. The disease was first described in the US in 1977 (6). In Europe and Asia, Lyme disease is usually caused by the following species of *Borrelia* spirocheta: *B. garinii* or *B. afzelii*, less often by *B. burgdorferi*, and rarely by *B. spielmanii* or *B. bavariensis* (7), in the US, mostly by *B. burgdorferi* and less often by *B. mayonii* (8). The pathogens are spread by ticks of the genus *Ixodes*. A person can be infected by both adult mites and nymphs. After a spirochete enters the human body, Lyme borreliosis occurs in three stages:

- The first, initially localized, stage usually occurs as a migratory erythema along with common signs of a viral infection. Migratory erythema usually occurs seven to 14 days after the bite but the period may vary from three to 30 days (5). The most frequent localization of the erythema is the areas of the armpit, groin, popliteal, and belt. Migratory erythema is hotter when touched; it is not painful, although it may sometimes cause pruritus or a burning sensation (9). In the first stage of the disease, symptoms similar to a viral infection

may occur: weakness, fever, muscle and joint pain, headache, and regional lymphadenopathy (10).

- The second, early disseminated, stage starts weeks or months after the tick bite. Organotropism and symptoms of *Borrelia* damage in the organs occur. Nervous system damage may occur and may include paralysis of the facial nerve, meningitis, radiculopathy, and peripheral neuropathy. Cardiac injury is clinically manifested by an atrioventricular block of varying degree (5). In case of eye damage, iridocyclitis, uveitis, conjunctivitis and optic nerve neuropathy develop (10).

- The third, late disseminated, stage can manifest months or even years after the tick bite. This stage is characterized by intermittent or persistent arthritis of the large joints, especially the knees (12). The following neurological disorders also develop: polyneuropathy, radiculopathy, chronic encephalomyelitis, and chronic atrophic acrodermatitis (13, 14).

Lyme disease very rarely manifests in cardiological disorders. In Europe, only 0.3–4% of all cases of Lyme borreliosis manifest in myocarditis (15). Damage of the heart tissue occurs three times more often in men than in women (16). *Borrelia*s, which enter the heart through the haematogenous pathway, cause transmural inflammations. The myocardial infiltration with inflammatory cells is mainly composed of lymphocytes (17). Although a spirochete-induced inflammation can affect many cardiac sites, studies with mice have shown that *Borrelia*s are most commonly found in the connective heart tissue, especially around

the aorta, in the upper ventricular and atrial epicardium, and in the myocardial interstitium and the endocardium (18). Acute myocarditis usually occurs with an atrioventricular block. In a review of 105 clinical cases conducted by van der Lind, 49% reported a complete AV block, 16% a second degree block, and 12% a first degree block (16). Myocarditis can also occur with supraventricular and ventricular arrhythmias, pericarditis and, in rare cases, pancarditis (19). Chronic Lyme myocarditis can be defined as chronic heart failure, with positive serological test results and Borrelia in the endomyocardial biopsate (19). In the course of chronic Lyme myocarditis, a very rare complication – dilated cardiomyopathy – may develop, but a link between it and LD has only been established in Europe (4, 20, 21).

Diagnosis of Lyme myocarditis

If Lyme myocarditis is suspected due to a sudden high-grade AV block, we recommend beginning with an evaluation of the likelihood of Lyme myocarditis by calculating the Suspicious Index in Lyme Carditis (SILC) (Table) (22). Our patient's SILC was retrospectively rated at 5 points: one point for age, male sex and outdoor activity, each, and: two points for constitutional symptoms.

If 0–2 points are collected, the patient is assigned to the low-risk group and other causes of myocarditis are sought. Patients with 3–6 points and 7–12 points (the moderate and high-risk groups, respectively) are subjected to serological tests: using an ELISA test for a search for anti-LD IgM and IgG antibodies and a Western blot test for confirmation (22). The results of the serological tests must be interpreted in the light of the

patient's clinical condition and of the course of the disease (23). A diagnosis of Lyme myocarditis is very complicated and other tests are needed to confirm the final diagnosis:

- Molecular testing for pathogen DNA detection is recommended only in certain situations, such as atypical LD manifestations in the skin or suspected Lyme arthritis (23).

- An electrocardiogram is required: it usually shows a I–III° atrioventricular block, but ST-segment depression and T wave inversion are also possible (16, 24).

- Radiology tests: cardiac echocardiography and magnetic resonance imaging (MRI) are needed. In these imaging examinations, structural changes, such as the pericardial changes found in pericarditis, can be assessed and left ventricular function can be evaluated in the presence of myocarditis, at which point hypokinetic myocardial zones are visible, and the ejection fraction can also be measured. The MRI examination depicts structures in high detail, so any myocardial oedema caused by the inflammatory process and hyperaemia is clearly visible (25). By using intravenous contrast during the MRI, the late contrast accumulation area, which highlights the area affected by the inflammation even more, can be recorded (26).

- An endomyocardial biopsy is the “gold standard” in the diagnostics of myocarditis, especially in complicated cases, such as progressive heart failure or a manifesting left ventricular dilation. Transmural inflammatory infiltration with lymphocytes is seen in biopsy samples. The detection of BB spirochetes in the inflammatory zone, which is mostly located along the collagen fibres, is also possible (17).

Table. Modified Suspicious Index in Lyme Carditis (SILC)

Variable	Value	Our patient
Age < 50 years	1	1
Male	1	1
Outdoor activity	1	1
Constitutional symptoms ¹	2	2
Tick bite	3	0
Erythema migrans	4	0
Total score	12	5

¹ Pre-syncope, syncope, fever, malaise, arthralgia, dyspnea.

Treatment of Lyme myocarditis and patient monitoring

In the case of a sudden high-grade AV block and suspected Lyme myocarditis, treatment should be started in accordance with the SILC score (see Fig. 3). Low-risk patients with a high-grade AV block should be treated based on the standard high-grade AV block treatment recommendations (22). Patients with suspected Lyme borreliosis (intermediate and high-risk patients), who have manifested syncope, dyspnoea, or chest pain and with detected second or third-degree AV block or first-degree AV block at a PR interval of more than 300 milliseconds, should be hospitalized, their heart rate monitored, and empirical antibiotic therapy initiated (22, 27). An atrioventricular block may progress in some patients with Lyme myocarditis, so it is recommended to always be prepared to implant a temporary pacemaker (16). Because any complete AV block is temporary and will disappear within a week if properly treated, implantation of a permanent pacemaker should be avoided (28).

The antibiotic therapy for the treatment of Lyme myocarditis is selected depending on the patient's condition. If the patient is haemodynamically stable, the first-line drug is Doxycycline, 100 mg orally twice a day for 21 days. If the patient is haemodynamically unstable, then Ceftriaxone, 1g twice a day for 21 days intravenously, should be administered. It is important to note that Azithromycin should not be given to patients with heart failure because it prolongs the QT interval (29).

CONCLUSIONS

When diagnosed in time and treated properly, the prognosis of Lyme myocarditis is usually good. In most cases, the atrioventricular block disappears within 1–2 weeks of antibiotic treatment and the implantation of a temporary pacemaker is rarely needed. In those rare cases of a chronic *Borrelia burgdorferi* infection, dilated cardiomyopathy may develop; thus if a sudden atrioventricular block occurs, the physician should be vigilant and perform the necessary tests to exclude the diagnosis of Lyme disease.

Received 13 March 2019

Accepted 3 September 2019

References

1. Steere AC, Strle F, Wormser GP, Hu LT, Branda JA, Hovius JW, et al. Lyme borreliosis. *Nat Rev Dis Primers*. 2016 Dec 15; 2: 16090.
2. Eldina C, Raffetin A, Bouiller K, Hansmann Y, Roblote F, Raoult D, et al. Review of European and American guidelines for the diagnosis of Lyme borreliosis. *Med Mal Infect*. 2018 Dec; pii: S0399-077X(18)30676-0.
3. Čaplinskas S, Čaplinskienė I, Davidavičienė E, Korabliovienė J, Liausėdienė R, Razmuvienė D, et al. Sergamumo užkrečiamosiomis ligomis Lietuvoje 2017 m. apžvalga. Užkrečiamųjų ligų ir AIDS centras [Internet]. Vilnius; 2018 [cited 2019 Jan 19]. 100 p. Available from: <http://www.ulac.lt/lt/>




Fig. 3. Diagnosis and management of Lyme carditis and a high-degree atrioventricular block. SILC – Suspicious Index in Lyme Carditis

- sergamumo-uzkreciamosiomis-ligomis-apzvalgos. Lithuanian.
- Haddad FA, Nadelman RB. Lyme disease and the heart. *Front Biosci*. 2003 Sep; 8: s769–82.
 - Wormser GP, Dattwyler RJ, Shapiro ED, Halperin JJ, Steere AC, Klemperer MS, et al. The clinical assessment, treatment, and prevention of Lyme disease, human granulocytic anaplasmosis, and babesiosis: clinical practice guidelines by the Infectious Diseases Society of America. *Clin Infect Dis*. 2006 Oct; 43(9): 1089–134.
 - Steere AC, Malawista SE, Snyderman DR, Shope RE, Andiman WA, Ross MR, et al. An epidemic of oligoarticular arthritis in children and adults in three connecticut communities. *Arthritis and Rheum*. 1977 Jan–Feb; 20(1): 7–17.
 - Rizzoli A, Hauffe HC, Carpi G, Vourc'h GI, Neteler M, Rosà R. Lyme borreliosis in Europe. *Euro Surveill*. 2011; 16(27): pii=19906.
 - Hoen AG, Margos G, Bent SJ, Diuk-Wasser MA, Barbour A, Kurtenbach K, et al. Phylogeography of *Borrelia burgdorferi* in the eastern United States reflects multiple independent Lyme disease emergence events. *Proc Natl Acad Sci USA*. 2009 Sep 1; 106(35): 15013–8.
 - Edlow JA. Erythema migrans. *Med Clin North Am*. 2002 Mar; 86(2): 239–60.
 - Nadelman RB, Nowakowski J, Forseter G, Goldberg NS, Bittker S, Cooper D, et al. The clinical spectrum of early lyme borreliosis in patients with culture-confirmed erythema migrans. *Am J Med*. 1996 May; 100(5): 502–8.
 - Mikkilä HO, Seppälä IJ, Viljanen MK, Peltomaa MP, Karma A. The expanding clinical spectrum of ocular lyme borreliosis. *Ophthalmology*. 2000 Mar; 107(3): 581–7.
 - Steere AC, Schoen RT, Taylor E. The clinical evolution of Lyme arthritis. *Ann Intern Med*. 1987 Nov; 107(5): 725–31.
 - Oschmann P, Dorndorf W, Hornig C, Schäfer C, Wellensiek HJ, Pflughaupt KW. Stages and syndromes of neuroborreliosis. *J Neurol*. 1998 May; 245(5): 262–72.
 - Stanek G, Strle F. Lyme borreliosis. *Lancet*. 2003 Nov; 362(9396): 1639–47.
 - Mayer W, Kleber FX, Wilske B, Preac-Mursic V, Maciejewski W, Sigl H, et al. Persistent atrioventricular block in Lyme borreliosis. *Klin Wochenschr*. 1990; 68(8): 431–5.
 - Van der Linde. Lyme carditis: clinical characteristics of 105 cases. *Scand J Infect Dis Suppl*. 1991; 77: 81–4.
 - Muehlenbachs A, Bollweg BC, Schulz TJ, Forrester JD, DeLeon Carnes M, Molins C, et al. Cardiac Troponin of *Borrelia burgdorferi*: An Autopsy Study of Sudden Cardiac Death Associated with Lyme Carditis. *Am J Pathol*. 2016 May; 186(5): 1195–205.
 - Armstrong AL, Barthold SW, Persing DH, Beck DS. Carditis in Lyme disease susceptible and resistant strains of laboratory mice infected with *Borrelia burgdorferi*. *Am J Trop Med Hyg*. 1992 Aug; 47(2): 249–58.
 - Bergler-Klein J, Ullrich R, Glogar D, Stanek G. Lyme borreliosis and cardiomyopathy. *Wien Med Wochenschr*. 1995; 145(7–8): 196–8.
 - Kuchynka P, Palecek T, Havranek S, Vitkova I, Nemecek E, Trckova R, et al. Recent-onset dilated cardiomyopathy associated with *Borrelia burgdorferi* infection. *Herz*. 2015 Sep; 40(6): 892–7.
 - Bartůnek P, Gorican K, Mrázek V, Varejka P, Veiser T, Hercogová J, et al. Lyme borreliosis infection as a cause of dilated cardiomyopathy. *Prague Med Rep*. 2006; 107(2): 213–26.
 - Yeung C, Baranchuk A. Systematic Approach to the Diagnosis and Treatment. *Healthcare*. 2018 Dec; 6(4): 119.
 - Dessau RB, van Dam AP, Fingerle V, Gray J, Hovius JW, Hunfeld KP, et al. To test or not to test? Laboratory support for the diagnosis of Lyme borreliosis: a position paper of ESGBOR, the ESCMID study group for Lyme borreliosis. *Clin Microbiol Infect*. 2018 Feb; 24(2): 118–24.
 - Steere AC, Batsford WP, Weinberg M, Alexander J, Berger HJ, Wolfson S, et al. Lyme carditis: cardiac abnormalities of Lyme disease. *Ann Intern Med*. 1980 Jul; 93(1): 8–16.
 - Chetrit M, Friedrich MG. The unique role of cardiovascular magnetic resonance imaging in acute myocarditis. *F1000Res*. 2018 Jul 30; 7. pii: F1000 Faculty Rev–1153.
 - Röttgen R, Christiani R, Freyhardt P, Gutberlet M, Schultheiss HP, Hamm B, et al. Magnetic resonance imaging findings in acute myocarditis and correlation with immunohistological parameters. *Eur Radiol*. 2011 Jun; 21(6): 1259–66.
 - Wormser GP, Dattwyler RJ, Shapiro ED, Halperin JJ, Steere AC, Klemperer MS, et al. The Clinical Assessment, Treatment, and Prevention of Lyme Disease,

- Human Granulocytic Anaplasmosis, and Babesiosis: Clinical Practice Guidelines by the Infectious Diseases Society of America. *Clin Infect Dis*. 2006 Nov 1; 43(9): 1089–134.
28. McAlister HF, Klementowicz PT, Andrews C, Fisher JD, Feld M, Furman S. Lyme carditis: an important cause of reversible heart block. *Ann Intern Med*. 1989 Mar 1; 110(5): 339–45.
29. Cruickshank M, O’Flynn N, Faust SN. Lyme disease: summary of NICE guidance. *BMJ*. 2018; 361: k1261.

**Audrius Aurelijus Pilypas, Giedrutė Raišlienė,
Jurgita Valaikienė**

**LAIMO LIGA IR ŠIRDIES TRANSPLANTACIJA:
KLINIKINIO ATVEJO PRISTATYMAS IR
LITERATŪROS APŽVALGA**

Santrauka

Tikslas. Laimo liga – transmisinė gamtinė židininė infekcinė liga, pažeidžianti įvairius organus ir sistemas. Ši liga dar vadinama Laimo borelioze, nes ją sukelia *Borrelia spirochetos*, kurias platina *Ixodes* genties erkės. Tai labiausiai išplitusi antropozoonozė. Ankstyva Laimo ligos diagnostika yra sudėtinga dėl dažnai pasireiškiančių atipinių ligos simptomų, nepastebėto erkės

įkandimo, nepasireiškiančios migruojančios eritemos, simptomų atsiradimo ne erkių sezono metu. Laiku nedidžius ir nepradėjus gydyti, įvyksta užkrato diseminacija, vystosi įvairios sunkios komplikacijos, nes borelijos pažeidžia ne tik odą, bet ir nervų sistemą, sąnarius, retais atvejais – širdį, akis.

Medžiaga ir metodai. Šiame straipsnyje pristatomas klinikinis atvejis apie Laimo borelioze sukeltą miokarditą, lėmusį dilatacinės kardiomiopatijos išsivystymą ir dėl to skubiai atliktą širdies transplantaciją. Mūsų turimais duomenimis, tai vienas pirmųjų aprašomų šios komplikacijos atvejų pasaulyje.

Išvados. Laiku diagnozavus ir paskyrus tinkamą gydymą, Laimo miokardito prognozė dažniausiai yra gera. Daugeliu atvejų, gydant antibiotikais, atrioventrikulinio mazgo blokada išnyksta per 1–2 savaites, o laikinojo kardiostimuliacijos implantavimo prireikia retai. Retais lėtinės *Borelia Burgdorferi* infekcijos atvejais gali išsivystyti dilatacinė kardiomiopatija, todėl nustačius staigiai atsiradusią atrioventrikulinio mazgo blokadą, gydytojais turi būti itin budrūs ir atlikti reikiamus tyrimus Laimo ligos diagnozei ekskliuduoti.

Raktažodžiai: Laimo miokarditas, atrioventrikulinio mazgo blokada, Laimo liga, dilatacinė kardiomiopatija