

## Research Article

### Influence of Preoperative Patient Factors on TKA Outcomes

Tomas Sveikata, Paulius Kanopa\*, Karolina Daunaraite, Paulius Mankevicius, Vytautas Tutkus, Giedrius Petryla, Jaunius Kurtinaitis, Sigitas Ryliskis, Narunas Porvaneckas, Valentinas Uvarovas and Igoris Satkauskas

Faculty of Medicine, Vilnius University, Vilnius, Lithuania

#### Abstract

**Background:** Osteoarthritis (OA) is one of the most disabling diseases in developed countries. In later stages of the disease, surgical treatment becomes the main alternative. Total Knee Arthroplasty (TKA) has proved a successful and cost-effective treatment for improving pain, function, mental health and overall quality of life in patients with severe knee arthritis. However up to 20% of patients are not satisfied with the outcome of the surgery. Being able to pre-operatively estimate which patient would have satisfactory results after the surgery is beneficial to guide further clinical decision. The aim of this study was to prospectively evaluate the influence of pre-operative patient factors on patient reported outcome after TKA.

**Methods:** 314 patients who underwent TKA in Vilnius University Hospital between end of 2012 and middle of 2014 year were included in a study. The influence of preoperative patient factors on the TKA outcome according to Western Ontario and McMaster Universities Osteoarthritis Index (WOMAC) change score was analyzed.

**Results:** The preoperative WOMAC pain score was the strongest determinant of the WOMAC pain change score one year after TKA. The preoperative WOMAC function and SF-12 mental health scores were the strongest determinants of the WOMAC function change score one year after TKA.

**Conclusion:** Lower preoperative knee pain, function scores and better mental health function results in greater change.

**Keywords:** Mental health; Osteoarthritis; Total knee arthroplasty; WOMAC

\*Corresponding author: Paulius Kanopa, Faculty of Medicine, Vilnius University, M. K. Čiurlionio st. 21, Vilnius, Lithuania, Tel: +370 68792500; E-mail: paulius.kanopa@gmail.com

**Citation:** Sveikata T, Kanopa P, Daunaraite K, Mankevicius P, Tutkus V, et al. (2021) Influence of Preoperative Patient Factors on TKA Outcomes. J Gerontol Geriatr Med 7: 087.

**Received:** February 22, 2021; **Accepted:** March 05, 2021; **Published:** March 12, 2021

**Copyright:** © 2021 Sveikata T, et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

#### Introduction


Osteoarthritis (OA) is one of the most disabling diseases in developed countries [1]. In later stages of the disease, surgery becomes the main alternative [2,3]. Total Knee Replacement (TKR) has proved a successful and cost-effective treatment for improving pain and function in patients with arthritis [3]. However up to 20% of patients are not satisfied with the surgery [4]. Therefore a traditional indicator of surgical success such as implant survival, may not mirror the patient's postoperative experience, which these days assessed with Patient-Reported Outcome Measures (PROMs) [5]. Patient-Reported Outcome Measures (PROMs) have become a corner-stone of outcome assessment after joint surgery [5]. Numerous PROMs are used to measure the outcome. Western Ontario and McMaster Universities Arthritis Index (WOMAC) and the Short Form-12 (SF-12) are most commonly used. WOMAC is a joint specific instrument for measuring clinical outcome in patients treated for knee osteoarthritis [6]. Various patient-related factors can be associated with poor outcome after TKA, including, age, gender, obesity, education and social support [7-9]. It's important to identify those preoperative risk factors in order to understand why some people fail to benefit from TKA. The aim of this study was to evaluate patient factors and preoperative pain, function according to WOMAC and physical and mental function according to SF-12 that might influence the patient reported results after TKA.

#### Methods

The prospective observational study was performed at Vilnius University Hospital, Vilnius, Lithuania, from October 2012 to April 2014. The study protocol was approved by the Regional Ethical Committee of the Vilnius Regional Bioethics Committee, Vilnius, M.K. Čiurlionio 21/27, LT-03101, Lithuania. The cohort consisted of patients from Vilnius University Hospital, orthopedic department. Patients scheduled for primary TKA were screened for inclusion. All enrolled patients gave a written consent. Flowchart of study cohort is shown in Figure 1. Three hundred fourteen participants were enrolled and two hundred ninety four patients completed the study.

Preoperative patient's data was collected during their preoperative evaluation. Information included sex, age, education, social support, height and weight was measured (body mass index (BMI) was calculated), residence place, co-morbidities. We divided all patients in to two groups according to age (younger than 75 and older). Usually the elderly are lumped together, grouping everyone over the age of 65. But a 65-year-old's experience of life is much different from a 90-year-old's. In general older adult population is divided into three life-stage subgroups: the young-old (approximately 65-74), the middle-old (ages 75-84), and the old-old (over age 85) [10]. We took a cut-off between young-old and middle-old subgroups. BMI is a simple index that is commonly used to classify underweight, overweight and obesity. We used WHO's recommendation for splitting patients in groups according to BMI: I. BMI<30 (patients with normal

weight and over-weight), II. BMI 30-34.9 (obese patients class I), III. BMI $\geq$ 35 (obese patients class II and class III) [11]. Patient's educational level either low (secondary school or vocational training) or high (University). Patients lived in rural or urban territories. Social support was determined by living alone or with somebody. Patients underwent radiographic examination and were grouped according to Ahlback osteoarthritis stages [12]. According to walking ability patients were grouped to Charnley classes [13].



Patients filled up two validated and originally authorized Lithuanian translations of questionnaires: the WOMAC and SF-12; before and one year after TKA. The WOMAC is a disease-specific instrument with 3 dimensions: pain, joint stiffness and physical function. It consists of 24 items (5 for pain, 2 for stiffness and 17 for function) [6]. We transformed WOMAC scores to a 0 to 100 point scale for each domain (with 100 being the best score).

The SF-12 score is a generic health measure of a patient's overall physical and mental wellbeing. Physical and Mental Health Composite Scores (PCS and MCS) are from 0 to 100, where zero indicates the lowest level of health and 100 the highest [14]. All TKA were made using cemented implants without resurfacing patella and all patients underwent similar rehabilitation program. Preoperative and in one year clinical outcome scores were analyzed as well as the change in score between preoperative and latest score.

### Statistical Analysis

Influence of preoperative patient factors on knee pain and function after TKA were analysed. Multiple linear regression was used to estimate the influence of preoperative factors on the outcomes according to WOMAC change score. A p-value of < 0.05 was considered to be statistically significant.

### Results

Preoperative predictors of WOMAC pain and function score change in one year after TKA were calculated. The multiple linear regression models are presented in tables 1 and 2.

Change WOMAC pain			
	Coef	95% CI	p-value
Age	-0.06	-0.21; 0.09	0.466
<b>Sex</b>			
Male	Ref.		
Female	-1.35	-4.50; 1.80	0.400
BMI	-0.04	-0.24; 0.16	0.697
<b>Social Support</b>			
No	Ref.		
Yes	0.46	-2.07; 2.99	0.720
<b>Level of Education</b>			
Higher	Ref.		
Lower	-0.37	-2.80; 2.05	0.762
<b>Residence</b>			
Urban	Ref.		
Rural	0.96	-2.07; 3.99	0.532
preop. WOMAC pain	-0.90	-0.97; -0.83	<0.001
<b>Charnley Class</b>			
A	Ref.		
B	-0.94	-3.73; 1.86	0.511
C	-1.46	-5.34; 2.42	0.459
<b>AHLBACK</b>			
1+2	Ref.		
3+4	1.19	-1.09; 3.47	0.305
<b>SF-12</b>			
preop. PCS	0.10	-0.06; 0.26	0.205
preop. MCS	0.10	-0.01; 0.22	0.080

Table 1: Preoperative predictors of WOMAC pain change score.

The preoperative WOMAC pain score was the strongest determinant of the WOMAC pain change score in one year (p<0.001). The preoperative WOMAC function (p<0.001) and SF-12 mental health scores (p=0.034) were the strongest determinants of the WOMAC function change score in one year after TKA.

### Discussion

Our study shows that preoperative knee pain, function and mental function predicts TKA outcome. We found that lower preoperative WOMAC scores and better mental function according to SF-12 are strong determinants of greater pain and function score change at one year after surgery. Several studies have reported that more preoperative pain and anxiety predicts worse postoperative pain [15-18]. Lingard A et al., concludes that patients with scores in the lowest WOMAC quartile group had the greatest improvement in their scores compared with the other groups [19]. A W-Dahl et al., reported that less pre-operative pain is a risk factor for no pain relief one year after surgery. Found that in addition to preoperative knee pain and pain at other sites, catastrophizing and mental health were the strongest independent predictors of persistent pain after TKA [20].

While many studies have explored how depression and lower mental health scores negatively impact TKA outcomes, the inverse has been seldom studied. Very few studies have looked in to the

effects TKA has on the patient's mental health [21]. The relationship of mental health, arthritis, and TKA is complex and not fully understood. Our findings show that following TKA, mental health status dramatically improves in patients with low MCS [22].

Change WOMAC function			
	Coef	95% CI	p-value
Age	-0.13	-0.29; 0.03	0.113
<b>Sex</b>			
Male	Ref.		
Female	-2.35	-5.70; 0.99	0.167
BMI	-0.05	-0.26; 0.16	0.652
<b>Social Support</b>			
No	Ref.		
Yes	1.74	-0.95; 4.42	0.204
<b>Level of Education</b>			
Higher	Ref.		
Lower	-0.62	-3.24; 2.00	0.642
<b>Residence</b>			
Urban	Ref.		
Rural	1.28	-1.93; 4.49	0.433
preop. WOMAC funct.	-0.93	-1.01; -0.85	<0.001
<b>Charnley Class</b>			
A	Ref.		
B	-1.95	-4.92; 1.03	0.198
C	-1.83	-5.94; 2.28	0.382
<b>AHLBACK</b>			
1+2	Ref.		
3+4	0.62	-1.81; 3.05	0.614
<b>SF-12</b>			
preop. PCS	0.09	-0.08; 0.26	0.302
preop. MCS	0.14	-0.01; 0.26	0.034

Table 2: Preoperative predictors of WOMAC function change score.

According to the results of our study, the quality of life of people after TKA significantly improves. This is due to pain factors that are closely related to mental health. These data correlates with our findings as there is more room for pain and function improvement as preoperative scores are lower. Study strengths are its prospective design, high rate of return to follow-up, all patients were operated in a single institution with the same postoperative and rehabilitation protocol.

## Conclusion

The preoperative WOMAC pain score was the strongest determinant of the WOMAC pain change score one year after TKA. The preoperative WOMAC function and SF-12 mental health scores were the strongest determinants of the WOMAC function change score one year after TKA. Lower preoperative knee pain, function scores and better mental health function results in greater change. Further studies are needed to assess the complex relationship of mental health and TKA. Specifically, it would be beneficial to evaluate the specific drivers of low emotional health and whether it is

the result of pain and dysfunction brought on by arthrosis, a lack of social support, catastrophizing patient characteristics, or something else. Furthermore, it would be significant to study the impact of preoperative mental and emotional interventions on patients with low mental health to see whether mental health recovers and improvement also corresponds to recovered outcomes after TKA [21].

## Declaration of Conflicting Interests

The author(s) declared no potential conflicts of interest with respect to the research, authorship, and/or publication of this article.

## References

1. WHO Scientific Group on the Burden of Musculoskeletal Conditions at the Start of the New Millennium (2003) The burden of musculoskeletal conditions at the start of the new millennium. World Health Organ Tech Rep Ser 919.
2. Lau RL, Perruccio AV, Gandhi R, Mahomed NN (2012) The role of surgeon volume on patient outcome in total knee arthroplasty: A systematic review of the literature. BMC Musculoskelet Disord 13: 250.
3. Ethgen O, Bruyère O, Richy F, Dardennes C, Reginster JY (2004) Health-related quality of life in total hip and total knee arthroplasty. A qualitative and systematic review of the literature. J Bone Joint Surg Am 86: 963-974.
4. Robertsson O, Dunbar M, Pehrsson T, Knutson K, Lidgren L (2000) Patient satisfaction after knee arthroplasty: a report on 27,372 knees operated on between 1981 and 1995 in Sweden. Acta Orthop Scand 71: 262-267.
5. Giesinger JM, Hamilton DF, Jost B, Behrend H, Giesinger K (2015) WOMAC, EQ-5D and Knee Society Score Thresholds for Treatment Success After Total Knee Arthroplasty. J Arthroplasty 30: 2154-2158.
6. American College of Rheumatology (2011) Clinical Research Resources. American College of Rheumatology, Atlanta, USA.
7. Ali A, Sundberg M, Robertsson O, Dahlberg LE, Thorstenson CA, et al. (2014) Dissatisfied patients after total knee arthroplasty. Acta Orthop 85: 229-233.
8. Nuñez M, Lozano L, Nuñez E, Segur JM, Sastre S (2011) Factors influencing health-related quality of life after TKA in patients who are obese. Clin Orthop Relat Res 469: 1148-1153.
9. Barrack RL, Ruh EL, Chen J, Lombardi AV Jr, Berend KR, et al. (2014) Impact of socioeconomic factors on outcome of total knee arthroplasty. Clin Orthop Relat Res 472: 86-97.
10. Little W (2016) Phases of Aging: The Young-Old, Middle-Old, and Old-Old. In: Little W (ed.). Introduction to Sociology.
11. WHO (2016) Global Database on Body Mass Index. WHO, Geneva, Switzerland.
12. Ahlbäck S (1968) Osteoarthritis of the knee. A radiographic investigation. Acta Radiol Diagn (Stockh) 277: 7-72.
13. Charnley J, Halley DK (1975) Rate of wear in total hip replacement. Clin Orthop Relat Res: 170-179.
14. Ware J Jr, Kosinski M, Keller SD (1996) A 12-Item Short-Form Health Survey: construction of scales and preliminary tests of reliability and validity. Med Care 34: 220-233.
15. Brander VA, Stulberg SD, Adams AD, Harden RN, Bruehl S, et al. (2003) Predicting total knee replacement pain: A prospective, observational study. Clin Orthop Relat Res: 27-36.
16. Lingard EA, Riddle DL (2007) Impact of psychological distress on pain and function following knee arthroplasty. J Bone Joint Surg Am 89: 1161-1169.

17. Judge A, Arden NK, Cooper C, Kassim Javaid M, et al. (2012) Predictors of outcomes of total knee replacement surgery. *Rheumatology (Oxford)* 51: 1804-1813.
18. Escobar A, Quintana JM, Bilbao A, Azkárate J, Güenaga JI, et al. (2007) Effect of patient characteristics on reported outcomes after total knee replacement. *Rheumatology (Oxford)* 46: 112-119.
19. Lingard EA, Katz JN, Wright EA, Sledge CB, Kinemax Outcomes Group (2004) Predicting the outcome of total knee arthroplasty. *J Bone Joint Surg Am* 86: 2179-2186.
20. Robertsson O, Ranstam J, Sundberg M, W-Dahl A, Lidgren L (2014) The Swedish Knee Arthroplasty Register: A review. *Bone Jt Res* 3: 217-222.
21. Horst PK, Barrett AA, Huddleston JI 3rd, Maloney WJ, Goodman SB, et al. (2020) Total Knee Arthroplasty Has A Positive Effect on Patients With Low Mental Health Scores. *J Arthroplasty* 35: 112-115.
22. Nicassio PM (2010) Arthritis and psychiatric disorders: disentangling the relationship. *J J Psychosom Res* 68: 183-185.



- Advances In Industrial Biotechnology | ISSN: 2639-5665
- Advances In Microbiology Research | ISSN: 2689-694X
- Archives Of Surgery And Surgical Education | ISSN: 2689-3126
- Archives Of Urology
- Archives Of Zoological Studies | ISSN: 2640-7779
- Current Trends Medical And Biological Engineering
- International Journal Of Case Reports And Therapeutic Studies | ISSN: 2689-310X
- Journal Of Addiction & Addictive Disorders | ISSN: 2578-7276
- Journal Of Agronomy & Agricultural Science | ISSN: 2689-8292
- Journal Of AIDS Clinical Research & STDs | ISSN: 2572-7370
- Journal Of Alcoholism Drug Abuse & Substance Dependence | ISSN: 2572-9594
- Journal Of Allergy Disorders & Therapy | ISSN: 2470-749X
- Journal Of Alternative Complementary & Integrative Medicine | ISSN: 2470-7562
- Journal Of Alzheimers & Neurodegenerative Diseases | ISSN: 2572-9608
- Journal Of Anesthesia & Clinical Care | ISSN: 2378-8879
- Journal Of Angiology & Vascular Surgery | ISSN: 2572-7397
- Journal Of Animal Research & Veterinary Science | ISSN: 2639-3751
- Journal Of Aquaculture & Fisheries | ISSN: 2576-5523
- Journal Of Atmospheric & Earth Sciences | ISSN: 2689-8780
- Journal Of Biotech Research & Biochemistry
- Journal Of Brain & Neuroscience Research
- Journal Of Cancer Biology & Treatment | ISSN: 2470-7546
- Journal Of Cardiology Study & Research | ISSN: 2640-768X
- Journal Of Cell Biology & Cell Metabolism | ISSN: 2381-1943
- Journal Of Clinical Dermatology & Therapy | ISSN: 2378-8771
- Journal Of Clinical Immunology & Immunotherapy | ISSN: 2378-8844
- Journal Of Clinical Studies & Medical Case Reports | ISSN: 2378-8801
- Journal Of Community Medicine & Public Health Care | ISSN: 2381-1978
- Journal Of Cytology & Tissue Biology | ISSN: 2378-9107
- Journal Of Dairy Research & Technology | ISSN: 2688-9315
- Journal Of Dentistry Oral Health & Cosmesis | ISSN: 2473-6783
- Journal Of Diabetes & Metabolic Disorders | ISSN: 2381-201X
- Journal Of Emergency Medicine Trauma & Surgical Care | ISSN: 2378-8798
- Journal Of Environmental Science Current Research | ISSN: 2643-5020
- Journal Of Food Science & Nutrition | ISSN: 2470-1076
- Journal Of Forensic Legal & Investigative Sciences | ISSN: 2473-733X
- Journal Of Gastroenterology & Hepatology Research | ISSN: 2574-2566
- Journal Of Genetics & Genomic Sciences | ISSN: 2574-2485
- Journal Of Gerontology & Geriatric Medicine | ISSN: 2381-8662
- Journal Of Hematology Blood Transfusion & Disorders | ISSN: 2572-2999
- Journal Of Hospice & Palliative Medical Care
- Journal Of Human Endocrinology | ISSN: 2572-9640
- Journal Of Infectious & Non Infectious Diseases | ISSN: 2381-8654
- Journal Of Internal Medicine & Primary Healthcare | ISSN: 2574-2493
- Journal Of Light & Laser Current Trends
- Journal Of Medicine Study & Research | ISSN: 2639-5657
- Journal Of Modern Chemical Sciences
- Journal Of Nanotechnology Nanomedicine & Nanobiotechnology | ISSN: 2381-2044
- Journal Of Neonatology & Clinical Pediatrics | ISSN: 2378-878X
- Journal Of Nephrology & Renal Therapy | ISSN: 2473-7313
- Journal Of Non Invasive Vascular Investigation | ISSN: 2572-7400
- Journal Of Nuclear Medicine Radiology & Radiation Therapy | ISSN: 2572-7419
- Journal Of Obesity & Weight Loss | ISSN: 2473-7372
- Journal Of Ophthalmology & Clinical Research | ISSN: 2378-8887
- Journal Of Orthopedic Research & Physiotherapy | ISSN: 2381-2052
- Journal Of Otolaryngology Head & Neck Surgery | ISSN: 2573-010X
- Journal Of Pathology Clinical & Medical Research
- Journal Of Pharmacology Pharmaceutics & Pharmacovigilance | ISSN: 2639-5649
- Journal Of Physical Medicine Rehabilitation & Disabilities | ISSN: 2381-8670
- Journal Of Plant Science Current Research | ISSN: 2639-3743
- Journal Of Practical & Professional Nursing | ISSN: 2639-5681
- Journal Of Protein Research & Bioinformatics
- Journal Of Psychiatry Depression & Anxiety | ISSN: 2573-0150
- Journal Of Pulmonary Medicine & Respiratory Research | ISSN: 2573-0177
- Journal Of Reproductive Medicine Gynaecology & Obstetrics | ISSN: 2574-2574
- Journal Of Stem Cells Research Development & Therapy | ISSN: 2381-2060
- Journal Of Surgery Current Trends & Innovations | ISSN: 2578-7284
- Journal Of Toxicology Current Research | ISSN: 2639-3735
- Journal Of Translational Science And Research
- Journal Of Vaccines Research & Vaccination | ISSN: 2573-0193
- Journal Of Virology & Antivirals
- Sports Medicine And Injury Care Journal | ISSN: 2689-8829
- Trends In Anatomy & Physiology | ISSN: 2640-7752

Submit Your Manuscript: <https://www.heraldopenaccess.us/submit-manuscript>